

SAT Math · Test II

1

★ Mark for Review

ABC

For the equation, b and c are constants. For any real number t , the point $\left(\frac{-2t+4}{3}, t\right)$ lies on the graph of the equation $bx + 3y = c$. What is the value of c ?

A. -2 B. 4 C. 6 D. 12

2

★ Mark for Review

ABC

The function q is defined by

$$q(x) = a|x - 9|^2 - 87|x - 9| + b,$$

where a and b are constants and $a > b > 87$. In the xy -plane, the graph of $y = q(x)$ contains the points $(639, h)$ and $(-621, k)$, where h and k are constants. What is the value of $9(-639)^{h-k} + 621(9)^{k-h}$?

3

★ Mark for Review

ABC

In the given equation, a and b are positive constants. A solution to the equation is $x = 42$.

$$x(9x - 2a) + b(9x - 2a) - (x + b) = 0$$

What is the value of a ?

4

★ Mark for Review

ABC

The function z is defined by

$$z(w) = (0.829)^{2w}.$$

The value of $z(w)$ decreases by $p\%$ for each increase by 1 in the value of w . Which of the following is closest to the value of p ?

A. 0.171

B. 0.829


C. 17.1

D. 31.3

5

★ Mark for Review

ABC



The graph of line h is shown in the xy -plane. Line k (not shown) is defined by the equation $sx + 40y = t$, where s and t are constants. If line k is graphed in this xy -plane, the result is a graph of two linear equations. This system of two linear equations has no solution. Which of the following is NOT a possible value of t ?

A. -360 B. -9 C. 72 D. 200

6

★ Mark for Review

ABC

The function h is defined by

$$h(x) = -\sqrt{x^2 + bx + c},$$

where b and c are constants. The graph of $y = h(x)$ contains the points $(2, 0)$ and $(0, -\sqrt{358})$. If $h(m) = 0$, what is the greatest possible value of m ?

A. 2

B. 177

C. 179

D. 358

7

★ Mark for Review

ABC

The functions f and g are defined by the equations shown, where a and b are integer constants, $a < b$, and $b < 0$. If $y = f(x)$ and $y = g(x)$ are graphed in the xy -plane, which of the following equations displays, as a constant or coefficient, the y -coordinate of the y -intercept of the graph of the corresponding function?

I. $f(x) = a(2.1)^{x+b}$

II. $g(x) = a(2.1)^x + b$

A. I only

B. II only

C. I and II

D. Neither I nor II

8

★ Mark for Review

ABC

The function f is defined by

$$f(x) = \frac{x^2 + ax + b}{2x + c},$$

where a , b , and c are constants. The graph of $y = f(x)$ does not intersect the line $x = 4$. If $f(5) = f(6) = 0$, what is the value of $a + b + c$?

A. 10**B.** 11**C.** 12**D.** 13

9

★ Mark for Review

ABC

The positive number a is 2,106% of the sum of the positive numbers b and c , and b is 81% of c . What percent of b is a ?

A. 4,706%**B.** 47.06%**C.** 470,600%**D.** 47,060%

10

★ Mark for Review

ABC

Point N lies on a unit circle in the xy -plane and has coordinates $(1, 0)$. Point O is the center of the circle and has coordinates $(0, 0)$. Point M also lies on the unit circle, and the measure of angle $\angle NOM$ is $\frac{266\pi}{3}$ radians. If the coordinates of point M are (a, b) , where a and b are constants, what is the value of a ?

A. $\frac{1}{2}$

B. $\frac{\sqrt{3}}{2}$

C. $-\frac{1}{2}$

D. $-\frac{\sqrt{3}}{2}$

11

★ Mark for Review

ABC

In the given equation, a and b are positive constants. The sum of the solutions to the given equation is $k(6a + b)$, where k is a constant. What is the value of k ?

$$24x^2 - (12a + 2b)x + ab = 0$$

A. $\frac{1}{12}$

B. $-\frac{1}{12}$

C. 12

D. -12

12

★ Mark for Review

ABC

In triangle ABC , the measure of angle B is 90° and BD is an altitude of the triangle. The length of $AB = 15$, and the length of AC is 23 greater than the length of AB . What is the value of $\frac{BC}{BD}$?

A. $\frac{15}{38}$

B. $\frac{15}{23}$

C. $\frac{23}{15}$

D. $\frac{38}{15}$

13

★ Mark for Review

ABC

A right square pyramid has a height of 11 centimeters (cm). A second right square pyramid has a height of 22 centimeters. The area of the bases of each of the two pyramids is 100 cm^2 . Which of the following is closest to the difference in the surface area of the second pyramid and the surface area of the first pyramid, in cm^2 ?

A. 209

B. 220

C. 367

D. 551

14

★ Mark for Review

ABC

A researcher investigated two species of mites: a predator and its prey. At the start of a week, there was an equal number of the two species. At the end of the week:

- The number of prey had increased by 2,700% of the number of prey at the start of the week.
- The number of predators had increased by 180% of the number of predators at the start of the week.

The number of prey at the end of the week was $p\%$ greater than the number of predators at the end of the week. What is the value of p ?

A. 9

B. 90

C. 900

D. 9,000

15

★ Mark for Review

ABC

A beekeeper's initial observation of the population of a certain bee colony was 1,100 bees. The beekeeper set a goal to increase the population to 2,600 bees. The beekeeper uses a model that predicts the population of this bee colony:

- begins at 1,100
- increases by 120 bees per week in the first two weeks
- then increases by 180 bees per week until the goal is reached

According to this model, at the end of week w after the initial observation, where $w > 2$, which of the following functions gives the predicted number of bees still needed to reach the beekeeper's goal?

A. $p(w) = 2600 - 180w$

B. $p(w) = 2480 + 180w$

C. $p(w) = 1620 - 180w$

D. $p(w) = -120 + 180w$

16

★ Mark for Review

ABC

The speed of a vehicle is increasing at a rate of 7.3 meters per second squared. What is this rate, in miles per minute squared, rounded to the nearest tenth? (Use 1 mile = 1,609 meters.)

A. 0.3**B.** 16.3**C.** 195.8**D.** 220.4

17

★ Mark for Review

ABC

Which of the following expressions has a factor of $x + 2b$, where b is a positive integer constant?

A. $3x^2 + 7x + 14b$

B. $3x^2 + 16x + 14b$

C. $3x^2 + 18x + 14b$

D. $3x^2 + 25x + 14b$

18

★ Mark for Review

ABC

In the given system of equations, t is a constant. If the system has no solution, what is the value of t ?

$$16x - 20y = 6y + 8$$

$$ty = 8x + \frac{1}{5}$$

19

★ Mark for Review

ABC

In the given equation, k is a positive constant. The product of the solutions to the equation is 72. What is the value of k ?

$$\frac{5}{9}(5x + 9)(x + \sqrt{5k + 9})(x - \sqrt{5k + 9}) = 0$$

20

★ Mark for Review

ABC

In the xy -plane, a unit circle with center at the origin O contains point A with coordinates $(1, 0)$ and point B with coordinates $\left(\frac{5}{\sqrt{34}}, -\frac{3}{\sqrt{34}}\right)$. If the measure of angle AOB is p radians, what is the value of $\frac{\cos p}{\sin p}$?

21

★ Mark for Review

ABC

A pyramid with square base area of 36 m^2 has a total surface area of $36 + 12\sqrt{493}\text{ m}^2$. Find the height of the pyramid.

22

★ Mark for Review

ABC

The height of a right circular cylinder is 36 inches, and the circumference of its base is 360 inches. Which expression represents the total surface area, in square inches, of the cylinder?

A. $(36)(360) + 2\pi \left(\frac{180}{\pi}\right)^2$

B. $(36)(360) + \pi \left(\frac{180}{\pi}\right)^2$


C. $\pi \left(\frac{180}{\pi}\right)^2 (36)$

D. $(36)(360)$

23

★ Mark for Review

ABC



The parallel line segment divides the other two sides of the triangle into segments of lengths m and n , and p and q , respectively. If $\frac{p}{q} = k$ and $\frac{m}{n} = k$, what is the value of k ?

24

★ Mark for Review


ABC

A gear ratio $r : s$ is the ratio of the number of teeth of two connected gears. The ratio of the number of revolutions per minute (rpm) of two gear wheels is $s : r$. In the diagram below, Gear A is turned by a motor. The turning of Gear A causes Gears B and C to turn as well.

A gear ratio $r:s$ is the ratio of the number of teeth of two connected gears. The ratio of the number of revolutions per minute (rpm) of two gear wheels is $s:r$. In the diagram below, Gear A is turned by a motor. The turning of Gear A causes Gears B and C to turn as well.

If Gear A is rotated by the motor at a rate of 100 rpm, what is the number of revolutions per minute for Gear C?

A) 50
B) 110
C) 200
D) 1,000



The diagram shows three interlocking gears. Gear A is on the left, Gear B is in the middle, and Gear C is on the right. Gear A has 20 teeth, Gear B has 60 teeth, and Gear C has 10 teeth. Gear A meshes with Gear B, and Gear B meshes with Gear C.

If Gear A is rotated by the motor at a rate of 100 rpm, what is the number of revolutions per minute for Gear C?

A. 50**B.** 110**C.** 200**D.** 1,000