

Base-two primitive permutation groups and their Saxl graphs

Hong Yi Huang

43rd Australasian Combinatorics Conference

13 December 2021



Let $G \leq \text{Sym}(\Omega)$ be a permutation group.

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G .

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Question: For a subset $\Delta \subseteq \Omega$, when do we have

$$G_{(\Delta)} := \bigcap_{\alpha \in \Delta} G_\alpha = 1?$$

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Question: For a subset $\Delta \subseteq \Omega$, when do we have

$$G_{(\Delta)} := \bigcap_{\alpha \in \Delta} G_\alpha = 1?$$

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$ and $\Delta = \{1, \dots, n-1\}$.

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Question: For a subset $\Delta \subseteq \Omega$, when do we have

$$G_{(\Delta)} := \bigcap_{\alpha \in \Delta} G_\alpha = 1?$$

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$ and $\Delta = \{1, \dots, n-1\}$.
- $G = \text{GL}(V)$, $\Omega = V$ and Δ contains a basis of V .

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

From now on, we will assume Ω is finite.

Base size $b(G)$: the minimal size of a base for G .

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

From now on, we will assume Ω is finite.

Base size $b(G)$: the minimal size of a base for G .

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 1$.

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

From now on, we will assume Ω is finite.

Base size $b(G)$: the minimal size of a base for G .

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 1$.
- $G = \text{GL}(V)$, $\Omega = V$: $b(G) = \dim(V)$.

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Question: Can we classify the finite transitive groups G with $b(G) = 2$?

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Question: Can we classify the finite transitive groups G with $b(G) = 2$?

Primitive group: G_α is a maximal subgroup of G .

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Question: Can we classify the finite transitive groups G with $b(G) = 2$?

Primitive group: G_α is a maximal subgroup of G .

Problem. Classify the finite primitive groups G with $b(G) = 2$.

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Question: Can we classify the finite transitive groups G with $b(G) = 2$?

Primitive group: G_α is a maximal subgroup of G .

Problem. Classify the finite primitive groups G with $b(G) = 2$.

Example

Consider the action of $G = D_{2n}$ on $\{1, \dots, n\}$. Then

- $\{1, 2\}$ is a base, so $b(G) = 2$;

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Question: Can we classify the finite transitive groups G with $b(G) = 2$?

Primitive group: G_α is a maximal subgroup of G .

Problem. Classify the finite primitive groups G with $b(G) = 2$.

Example

Consider the action of $G = D_{2n}$ on $\{1, \dots, n\}$. Then

- $\{1, 2\}$ is a base, so $b(G) = 2$;
- G is primitive iff n is a prime.

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem

Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem

Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem

Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Almost simple: $\text{Inn}(T) \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem

Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Almost simple: $\text{Inn}(T) \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem

Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Almost simple: $\text{Inn}(T) \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- CFSG is used. Partial results (**Burness et al.**)

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem

Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Almost simple: $\text{Inn}(T) \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- CFSG is used. Partial results (**Burness et al.**)

Diagonal and twisted wreath types: Partial results (**Fawcett, 2013/21**)

Some results

The **O’Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem

Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Almost simple: $\text{Inn}(T) \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- CFSG is used. Partial results (**Burness et al.**)

Diagonal and twisted wreath types: Partial results (**Fawcett, 2013/21**)

Product type: In progress

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Examples

- $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$:


Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Examples

- $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 


Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Examples

- $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 
- $G = C_4 = \langle (1234) \rangle$, $\Omega = \{1, 2, 3, 4\}$:


Saxl graphs


Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Examples

• $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 

• $G = C_4 = \langle (1234) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 


Saxl graphs


Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Examples

• $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 

• $G = C_4 = \langle (1234) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 

• $G = D_{10} = \langle (12345), (25)(34) \rangle$, $\Omega = \{1, 2, 3, 4, 5\}$:


Saxl graphs


Definition (Burness & Giudici, 2020)


Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Examples

• $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 

• $G = C_4 = \langle (1234) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 

• $G = D_{10} = \langle (12345), (25)(34) \rangle$, $\Omega = \{1, 2, 3, 4, 5\}$: $\Sigma(G) =$ 

Another example

Let $G = \mathrm{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 .

- $G_\alpha = D_{2(q-1)}$;
- α and β form a base iff they share a common 1-space.

Another example

Let $G = \mathrm{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 .

- $G_\alpha = D_{2(q-1)}$;
- α and β form a base iff they share a common 1-space.

Hence, $\Sigma(G) \cong J(q+1, 2)$ is a **Johnson graph**: vertices 2-subsets of $\{1, \dots, q+1\}$ and two vertices are adjacent if they are not disjoint.


Another example

Let $G = \text{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 .

- $G_\alpha = D_{2(q-1)}$;
- α and β form a base iff they share a common 1-space.

Hence, $\Sigma(G) \cong J(q+1, 2)$ is a **Johnson graph**: vertices 2-subsets of $\{1, \dots, q+1\}$ and two vertices are adjacent if they are not disjoint.

For example, when $q = 4$ we have the complement of the Petersen graph.



First observations

From now on, we will assume $b(G) = 2$ and G is transitive.

First observations

From now on, we will assume $b(G) = 2$ and G is transitive. Then

- $\Sigma(G)$ is G -vertex-transitive;

First observations

From now on, we will assume $b(G) = 2$ and G is transitive. Then

- $\Sigma(G)$ is G -vertex-transitive;
- $K < G \implies \Sigma(G)$ is a subgraph of $\Sigma(K)$;

First observations

From now on, we will assume $b(G) = 2$ and G is transitive. Then

- $\Sigma(G)$ is G -vertex-transitive;
- $K < G \implies \Sigma(G)$ is a subgraph of $\Sigma(K)$;
- $\Sigma(G)$ has valency $r|G_\alpha|$, where r is the number of regular G_α -orbits;

First observations

From now on, we will assume $b(G) = 2$ and G is transitive. Then

- $\Sigma(G)$ is G -vertex-transitive;
- $K < G \implies \Sigma(G)$ is a subgraph of $\Sigma(K)$;
- $\Sigma(G)$ has valency $r|G_\alpha|$, where r is the number of regular G_α -orbits;
- $\Sigma(G)$ is the union of the regular **orbital graphs** of G ;

First observations

From now on, we will assume $b(G) = 2$ and G is transitive. Then

- $\Sigma(G)$ is G -vertex-transitive;
- $K < G \implies \Sigma(G)$ is a subgraph of $\Sigma(K)$;
- $\Sigma(G)$ has valency $r|G_\alpha|$, where r is the number of regular G_α -orbits;
- $\Sigma(G)$ is the union of the regular **orbital graphs** of G ;
- G is primitive $\implies \Sigma(G)$ is connected.

First observations

From now on, we will assume $b(G) = 2$ and G is transitive. Then

- $\Sigma(G)$ is G -vertex-transitive;
- $K < G \implies \Sigma(G)$ is a subgraph of $\Sigma(K)$;
- $\Sigma(G)$ has valency $r|G_\alpha|$, where r is the number of regular G_α -orbits;
- $\Sigma(G)$ is the union of the regular **orbital graphs** of G ;
- G is primitive $\implies \Sigma(G)$ is connected.

Question: What is the diameter of $\Sigma(G)$ if G is primitive?

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓
- “Most” almost simple groups with alternating or sporadic socles

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓
- “Most” almost simple groups with alternating or sporadic socles
- **Chen & Du, 2021; Burness & H, 2021:** $\text{soc}(G) = L_2(q)$ ✓

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓
- “Most” almost simple groups with alternating or sporadic socles
- **Chen & Du, 2021; Burness & H, 2021:** $\text{soc}(G) = L_2(q)$ ✓
- **Burness & H, 2021:** almost simple groups with soluble stabilisers ✓

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓
- “Most” almost simple groups with alternating or sporadic socles
- **Chen & Du, 2021; Burness & H, 2021:** $\text{soc}(G) = L_2(q)$ ✓
- **Burness & H, 2021:** almost simple groups with soluble stabilisers ✓
- **Lee & Popiel, 2021:** some affine groups

Soluble stabilisers

$\mathcal{S} := \{\text{almost simple primitive groups with soluble stabilisers}\}$

Soluble stabilisers

$\mathcal{S} := \{\text{almost simple primitive groups with soluble stabilisers}\}$

- **Li & Zhang, 2011:** \mathcal{S} is completely known ✓

Soluble stabilisers

$\mathcal{S} := \{\text{almost simple primitive groups with soluble stabilisers}\}$

- **Li & Zhang, 2011:** \mathcal{S} is completely known ✓

$\mathcal{B} := \{G \in \mathcal{S} \mid b(G) = 2\}$

Soluble stabilisers

$\mathcal{S} := \{\text{almost simple primitive groups with soluble stabilisers}\}$

- **Li & Zhang, 2011:** \mathcal{S} is completely known ✓

$\mathcal{B} := \{G \in \mathcal{S} \mid b(G) = 2\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓

Soluble stabilisers

$\mathcal{S} := \{\text{almost simple primitive groups with soluble stabilisers}\}$

- **Li & Zhang, 2011:** \mathcal{S} is completely known ✓

$\mathcal{B} := \{G \in \mathcal{S} \mid b(G) = 2\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓
- **Burness & H, 2021:** The common neighbour property for \mathcal{B} ✓

Soluble stabilisers

$\mathcal{S} := \{\text{almost simple primitive groups with soluble stabilisers}\}$

- **Li & Zhang, 2011:** \mathcal{S} is completely known ✓

$\mathcal{B} := \{G \in \mathcal{S} \mid b(G) = 2\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓
- **Burness & H, 2021:** The common neighbour property for \mathcal{B} ✓

Clique number: maximal size $\omega(G)$ of a complete subgraph of $\Sigma(G)$.

Soluble stabilisers

$\mathcal{S} := \{\text{almost simple primitive groups with soluble stabilisers}\}$

- **Li & Zhang, 2011:** \mathcal{S} is completely known ✓

$\mathcal{B} := \{G \in \mathcal{S} \mid b(G) = 2\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓
- **Burness & H, 2021:** The common neighbour property for \mathcal{B} ✓

Clique number: maximal size $\omega(G)$ of a complete subgraph of $\Sigma(G)$.

Independence number: clique number $\alpha(G)$ of the complement of $\Sigma(G)$.

Soluble stabilisers

$\mathcal{S} := \{\text{almost simple primitive groups with soluble stabilisers}\}$

- **Li & Zhang, 2011:** \mathcal{S} is completely known ✓

$\mathcal{B} := \{G \in \mathcal{S} \mid b(G) = 2\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓
- **Burness & H, 2021:** The common neighbour property for \mathcal{B} ✓

Clique number: maximal size $\omega(G)$ of a complete subgraph of $\Sigma(G)$.

Independence number: clique number $\alpha(G)$ of the complement of $\Sigma(G)$.

Example: $(G, G_\alpha) = (A_5, S_3) \implies \Sigma(G) = J(5, 2), \omega(G) = 4, \alpha(G) = 2.$

Soluble stabilisers

$\mathcal{S} := \{\text{almost simple primitive groups with soluble stabilisers}\}$

- **Li & Zhang, 2011:** \mathcal{S} is completely known ✓

$\mathcal{B} := \{G \in \mathcal{S} \mid b(G) = 2\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓
- **Burness & H, 2021:** The common neighbour property for \mathcal{B} ✓

Clique number: maximal size $\omega(G)$ of a complete subgraph of $\Sigma(G)$.

Independence number: clique number $\alpha(G)$ of the complement of $\Sigma(G)$.

Example: $(G, G_\alpha) = (A_5, S_3) \implies \Sigma(G) = J(5, 2), \omega(G) = 4, \alpha(G) = 2.$

Theorem (Burness & H, 2021)

- $G \in \mathcal{B}$ is simple $\implies \omega(G) \geq 5$ or $(G, G_\alpha) = (A_5, S_3);$

Soluble stabilisers

$\mathcal{S} := \{\text{almost simple primitive groups with soluble stabilisers}\}$

- **Li & Zhang, 2011:** \mathcal{S} is completely known ✓

$\mathcal{B} := \{G \in \mathcal{S} \mid b(G) = 2\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓
- **Burness & H, 2021:** The common neighbour property for \mathcal{B} ✓

Clique number: maximal size $\omega(G)$ of a complete subgraph of $\Sigma(G)$.

Independence number: clique number $\alpha(G)$ of the complement of $\Sigma(G)$.

Example: $(G, G_\alpha) = (A_5, S_3) \implies \Sigma(G) = J(5, 2), \omega(G) = 4, \alpha(G) = 2$.

Theorem (Burness & H, 2021)

- $G \in \mathcal{B}$ is simple $\implies \omega(G) \geq 5$ or $(G, G_\alpha) = (A_5, S_3)$;
- $G \in \mathcal{B} \implies \alpha(G) \geq 4$ or $(G, G_\alpha) = (A_5, S_3)$.

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n and let $v(G)$ be the valency of $\Sigma(G)$.

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n and let $v(G)$ be the valency of $\Sigma(G)$. Then

$$Q(G) := \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2} = 1 - \frac{v(G)}{n}$$

is the probability that a random pair in Ω is not a base for G .

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n and let $v(G)$ be the valency of $\Sigma(G)$. Then

$$Q(G) := \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2} = 1 - \frac{v(G)}{n}$$

is the probability that a random pair in Ω is not a base for G .

Notes:

- $Q(G) < 1 \iff b(G) \leq 2$.

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n and let $v(G)$ be the valency of $\Sigma(G)$. Then

$$Q(G) := \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2} = 1 - \frac{v(G)}{n}$$

is the probability that a random pair in Ω is not a base for G .

Notes:

- $Q(G) < 1 \iff b(G) \leq 2$.
- $Q(G) < 1/2 \implies \Sigma(G)$ has the common neighbour property.

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n and let $v(G)$ be the valency of $\Sigma(G)$. Then

$$Q(G) := \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2} = 1 - \frac{v(G)}{n}$$

is the probability that a random pair in Ω is not a base for G .

Notes:

- $Q(G) < 1 \iff b(G) \leq 2$.
- $Q(G) < 1/2 \implies \Sigma(G)$ has the common neighbour property.
- $Q(G) < 1/t \implies \omega(G) \geq t + 1$.

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n and let $v(G)$ be the valency of $\Sigma(G)$. Then

$$Q(G) := \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2} = 1 - \frac{v(G)}{n}$$

is the probability that a random pair in Ω is not a base for G .

Notes:

- $Q(G) < 1 \iff b(G) \leq 2$.
- $Q(G) < 1/2 \implies \Sigma(G)$ has the common neighbour property.
- $Q(G) < 1/t \implies \omega(G) \geq t + 1$.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $Q(G) \rightarrow 1$ as $q \rightarrow \infty$. But $\Sigma(G) = J(q+1, 2)$ still has the common neighbour property.

Future work

Valency: Study the number $r(G)$ of regular G_α -orbits.

Future work

Valency: Study the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021:** some general methods

Future work

Valency: Study the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021:** some general methods
- Probabilistic methods: lower bounds for $r(G)$

Future work

Valency: Study the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021:** some general methods
- Probabilistic methods: lower bounds for $r(G)$

Problem. Classify the primitive groups G with $r(G) = 1$.

Future work

Valency: Study the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021:** some general methods
- Probabilistic methods: lower bounds for $r(G)$

Problem. Classify the primitive groups G with $r(G) = 1$.

Burness & H, 2021: $G \in \mathcal{B}$ and $r(G) = 1$ ✓

Future work

Valency: Study the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021:** some general methods
- Probabilistic methods: lower bounds for $r(G)$

Problem. Classify the primitive groups G with $r(G) = 1$.

Burness & H, 2021: $G \in \mathcal{B}$ and $r(G) = 1$ ✓

e.g. $G = \mathrm{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$.

Future work

Valency: Study the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021:** some general methods
- Probabilistic methods: lower bounds for $r(G)$

Problem. Classify the primitive groups G with $r(G) = 1$.

Burness & H, 2021: $G \in \mathcal{B}$ and $r(G) = 1$ ✓

e.g. $G = \mathrm{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$.

Product-type primitive groups: $G \leq L \wr P$ with its product action on $\Omega = \Gamma^k$.

Future work

Valency: Study the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021:** some general methods
- Probabilistic methods: lower bounds for $r(G)$

Problem. Classify the primitive groups G with $r(G) = 1$.

Burness & H, 2021: $G \in \mathcal{B}$ and $r(G) = 1$ ✓

e.g. $G = \mathrm{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$.

Product-type primitive groups: $G \leq L \wr P$ with its product action on $\Omega = \Gamma^k$.

Theorem (Bailey & Cameron, 2013)

$b(L \wr P) = 2 \iff r(L) \geq \text{the distinguishing number of } P.$

Future work

Valency: Study the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021:** some general methods
- Probabilistic methods: lower bounds for $r(G)$

Problem. Classify the primitive groups G with $r(G) = 1$.

Burness & H, 2021: $G \in \mathcal{B}$ and $r(G) = 1$ ✓

e.g. $G = \mathrm{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$.

Product-type primitive groups: $G \leq L \wr P$ with its product action on $\Omega = \Gamma^k$.

Theorem (Bailey & Cameron, 2013)

$b(L \wr P) = 2 \iff r(L) \geq \text{the distinguishing number of } P.$

Questions:

- Properties of the Saxl graph $\Sigma(G)$ when $G = L \wr P$?

Future work

Valency: Study the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021:** some general methods
- Probabilistic methods: lower bounds for $r(G)$

Problem. Classify the primitive groups G with $r(G) = 1$.

Burness & H, 2021: $G \in \mathcal{B}$ and $r(G) = 1$ ✓

e.g. $G = \mathrm{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$.

Product-type primitive groups: $G \leq L \wr P$ with its product action on $\Omega = \Gamma^k$.

Theorem (Bailey & Cameron, 2013)

$b(L \wr P) = 2 \iff r(L) \geq \text{the distinguishing number of } P.$

Questions:

- Properties of the Saxl graph $\Sigma(G)$ when $G = L \wr P$?
- If $G < L \wr P$ is primitive, then when do we have $b(G) = 2$?

Thank you!