

Base-two primitive permutation groups and their Saxl graphs

Hong Yi Huang

University of Bristol

LMS Graduate Student Meeting

8 November 2021



Let $G \leq \text{Sym}(\Omega)$ be a permutation group.

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G .

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Question: For a subset $\Delta \subseteq \Omega$, when do we have

$$G_{(\Delta)} := \bigcap_{\alpha \in \Delta} G_\alpha = 1?$$

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Question: For a subset $\Delta \subseteq \Omega$, when do we have

$$G_{(\Delta)} := \bigcap_{\alpha \in \Delta} G_\alpha = 1?$$

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$ and $\Delta = \{1, \dots, n-1\}$.

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Question: For a subset $\Delta \subseteq \Omega$, when do we have

$$G_{(\Delta)} := \bigcap_{\alpha \in \Delta} G_\alpha = 1?$$

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$ and $\Delta = \{1, \dots, n-1\}$.
- $G = \text{GL}(V)$, $\Omega = V$ and Δ contains a basis of V .

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

From now, we assume Ω is finite.

Base size $b(G)$: the minimal size of a base for G .

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

From now, we assume Ω is finite.

Base size $b(G)$: the minimal size of a base for G .

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 1$.

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

From now, we assume Ω is finite.

Base size $b(G)$: the minimal size of a base for G .

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 1$.
- $G = \text{GL}(V)$, $\Omega = V$: $b(G) = \dim(V)$.

Base-two groups

Notes: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;

Base-two groups

Notes: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Base-two groups

Notes: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Problem

Classify the finite transitive groups G with $b(G) = 2$.

Base-two groups

Notes: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Problem

Classify the finite transitive groups G with $b(G) = 2$.

G is called **primitive** if G_α is a maximal subgroup of G .

Base-two groups

Notes: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Problem

Classify the finite transitive groups G with $b(G) = 2$.

G is called **primitive** if G_α is a maximal subgroup of G .

Problem

Classify the finite primitive groups G with $b(G) = 2$.

Base-two groups

Notes: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Problem

Classify the finite transitive groups G with $b(G) = 2$.

G is called **primitive** if G_α is a maximal subgroup of G .

Problem

Classify the finite primitive groups G with $b(G) = 2$.

Example

Consider the action of $G = D_{2n}$ on $\{1, \dots, n\}$. Then

- $\{1, 2\}$ is a base, so $b(G) = 2$;

Base-two groups

Notes: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Problem

Classify the finite transitive groups G with $b(G) = 2$.

G is called **primitive** if G_α is a maximal subgroup of G .

Problem

Classify the finite primitive groups G with $b(G) = 2$.

Example

Consider the action of $G = D_{2n}$ on $\{1, \dots, n\}$. Then

- $\{1, 2\}$ is a base, so $b(G) = 2$;
- G is primitive iff n is a prime.

Some results

The **O’Nan-Scott Theorem** divides finite primitive groups into 5 types.

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem

Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem

Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

Almost simple: $\text{Inn}(T) \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem

Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

Almost simple: $\text{Inn}(T) \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

Some results

The **O’Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem

Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

Almost simple: $\text{Inn}(T) \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- CFSG is used. Partial results.

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Example

$G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$:

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Example

$$G = D_8 = \langle (1234), (24) \rangle, \Omega = \{1, 2, 3, 4\}: \Sigma(G) = C_4 = \square$$

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Example

$$G = D_8 = \langle (1234), (24) \rangle, \Omega = \{1, 2, 3, 4\}: \Sigma(G) = C_4 = \square$$

From now, we assume $b(G) = 2$ and G is transitive.

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Example

$$G = D_8 = \langle (1234), (24) \rangle, \Omega = \{1, 2, 3, 4\}: \Sigma(G) = C_4 = \square$$

From now, we assume $b(G) = 2$ and G is transitive. We have

- $\Sigma(G)$ is vertex-transitive;

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Example

$$G = D_8 = \langle (1234), (24) \rangle, \Omega = \{1, 2, 3, 4\}: \Sigma(G) = C_4 = \square$$

From now, we assume $b(G) = 2$ and G is transitive. We have

- $\Sigma(G)$ is vertex-transitive;
- $\Sigma(G)$ is connected if G is primitive;

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Example

$$G = D_8 = \langle (1234), (24) \rangle, \Omega = \{1, 2, 3, 4\}: \Sigma(G) = C_4 = \square$$

From now, we assume $b(G) = 2$ and G is transitive. We have

- $\Sigma(G)$ is vertex-transitive;
- $\Sigma(G)$ is connected if G is primitive;
- $\Sigma(G)$ has valency $r|G_\alpha|$, where r is the number of regular G_α -orbits.

A further example

Let $G = \mathrm{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 .

- $G_\alpha = D_{2(q-1)}$;
- α and β form a base iff they share a common 1-space.

A further example

Let $G = \mathrm{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 .

- $G_\alpha = D_{2(q-1)}$;
- α and β form a base iff they share a common 1-space.

Hence, $\Sigma(G) \cong J(q+1, 2)$ is a **Johnson graph**: vertices 2-subsets of $\{1, \dots, q+1\}$ and two vertices are adjacent if they are not disjoint.


A further example

Let $G = \text{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 .

- $G_\alpha = D_{2(q-1)}$;
- α and β form a base iff they share a common 1-space.

Hence, $\Sigma(G) \cong J(q+1, 2)$ is a **Johnson graph**: vertices 2-subsets of $\{1, \dots, q+1\}$ and two vertices are adjacent if they are not disjoint.

For example, when $q = 4$ we have the complement of the Petersen graph.



Diameter

Recall: $\Sigma(G)$ is connected if G is primitive.

Diameter

Recall: $\Sigma(G)$ is connected if G is primitive.

Question: What is the diameter of $\Sigma(G)$ if G is primitive?

Diameter

Recall: $\Sigma(G)$ is connected if G is primitive.

Question: What is the diameter of $\Sigma(G)$ if G is primitive?

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Diameter

Recall: $\Sigma(G)$ is connected if G is primitive.

Question: What is the diameter of $\Sigma(G)$ if G is primitive?

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Diameter

Recall: $\Sigma(G)$ is connected if G is primitive.

Question: What is the diameter of $\Sigma(G)$ if G is primitive?

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Theorem (Chen & Du, 2021+; Burness & H, 2021+)

$\text{soc}(G) = \text{L}_2(q) \implies \Sigma(G)$ has the common neighbour property.

Soluble stabiliser

Li & Zhang 2011: Almost simple primitive groups with G_α soluble ✓

Soluble stabiliser

Li & Zhang 2011: Almost simple primitive groups with G_α soluble ✓

Burness 2021: The set \mathcal{G} of those base-two groups ✓

Soluble stabiliser

Li & Zhang 2011: Almost simple primitive groups with G_α soluble ✓

Burness 2021: The set \mathcal{G} of those base-two groups ✓

Theorem (Burness & H, 2021+)

$G \in \mathcal{G} \implies \Sigma(G)$ has the common neighbour property.

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n and let $v(G)$ be the valency of $\Sigma(G)$.

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n and let $v(G)$ be the valency of $\Sigma(G)$. Then

$$Q(G) := \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2} = 1 - \frac{v(G)}{n}$$

is the probability that a random pair in Ω is not a base for G .

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n and let $v(G)$ be the valency of $\Sigma(G)$. Then

$$Q(G) := \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2} = 1 - \frac{v(G)}{n}$$

is the probability that a random pair in Ω is not a base for G .

Notes:

- $Q(G) < 1 \iff b(G) \leq 2$.

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n and let $v(G)$ be the valency of $\Sigma(G)$. Then

$$Q(G) := \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2} = 1 - \frac{v(G)}{n}$$

is the probability that a random pair in Ω is not a base for G .

Notes:

- $Q(G) < 1 \iff b(G) \leq 2$.
- $Q(G) < 1/2 \implies \Sigma(G)$ has the common neighbour property.

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n and let $v(G)$ be the valency of $\Sigma(G)$. Then

$$Q(G) := \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2} = 1 - \frac{v(G)}{n}$$

is the probability that a random pair in Ω is not a base for G .

Notes:

- $Q(G) < 1 \iff b(G) \leq 2$.
- $Q(G) < 1/2 \implies \Sigma(G)$ has the common neighbour property.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $Q(G) \rightarrow 1$ as $q \rightarrow \infty$. But $\Sigma(G) = J(q+1, 2)$ still has the common neighbour property.

Future work

Valency: Determine the number $r(G)$ of regular G_α -orbits.

Future work

Valency: Determine the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021+:** some general methods

Future work

Valency: Determine the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021+:** some general methods
- **Burness & H, 2021+:** $G \in \mathcal{G}$ and $r(G) = 1$ ✓

Future work

Valency: Determine the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021+:** some general methods
- **Burness & H, 2021+:** $G \in \mathcal{G}$ and $r(G) = 1$ ✓

Product type primitive groups: $G \leq L \wr P$.

Future work

Valency: Determine the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021+:** some general methods
- **Burness & H, 2021+:** $G \in \mathcal{G}$ and $r(G) = 1$ ✓

Product type primitive groups: $G \leq L \wr P$.

Theorem (Bailey & Cameron 2013)

$b(L \wr P) = 2 \iff r(L) \geq \text{the distinguishing number of } P$.

Future work

Valency: Determine the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021+:** some general methods
- **Burness & H, 2021+:** $G \in \mathcal{G}$ and $r(G) = 1$ ✓

Product type primitive groups: $G \leq L \wr P$.

Theorem (Bailey & Cameron 2013)

$b(L \wr P) = 2 \iff r(L) \geq \text{the distinguishing number of } P$.

Questions:

- The Saxl graph $\Sigma(G)$ when $G = L \wr P$?

Future work

Valency: Determine the number $r(G)$ of regular G_α -orbits.

- **Chen & H, 2021+:** some general methods
- **Burness & H, 2021+:** $G \in \mathcal{G}$ and $r(G) = 1$ ✓

Product type primitive groups: $G \leq L \wr P$.

Theorem (Bailey & Cameron 2013)

$b(L \wr P) = 2 \iff r(L) \geq \text{the distinguishing number of } P$.

Questions:

- The Saxl graph $\Sigma(G)$ when $G = L \wr P$?
- If $G < L \wr P$, then when do we have $b(G) = 2$?

Thank you!