

Base-two primitive groups and their Saxl graphs

Hong Yi Huang

Groups and Graphs

24 July 2022



Let $G \leq \text{Sym}(\Omega)$ be a permutation group.

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G .

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Question: For a subset $\Delta \subseteq \Omega$, when do we have

$$G_{(\Delta)} := \bigcap_{\alpha \in \Delta} G_\alpha = 1?$$

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Question: For a subset $\Delta \subseteq \Omega$, when do we have

$$G_{(\Delta)} := \bigcap_{\alpha \in \Delta} G_\alpha = 1?$$

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$ and $\Delta = \{1, \dots, n-1\}$.

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Question: For a subset $\Delta \subseteq \Omega$, when do we have

$$G_{(\Delta)} := \bigcap_{\alpha \in \Delta} G_\alpha = 1?$$

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$ and $\Delta = \{1, \dots, n-1\}$.
- $G = \text{GL}(V)$, $\Omega = V \setminus \{0\}$ and Δ contains a basis of V .

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

From now on, we will assume Ω is finite.

Base size $b(G)$: the minimal size of a base for G .

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

From now on, we will assume Ω is finite.

Base size $b(G)$: the minimal size of a base for G .

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 1$.

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

From now on, we will assume Ω is finite.

Base size $b(G)$: the minimal size of a base for G .

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 1$.
- $G = \text{GL}(V)$, $\Omega = V \setminus \{0\}$: $b(G) = \dim(V)$.

Base sizes

Observation: If Δ is a base and $x, y \in G$, then

$$\alpha^x = \alpha^y \text{ for all } \alpha \in \Delta \iff xy^{-1} \in \bigcap_{\alpha \in \Delta} G_\alpha = G_{(\Delta)} \iff x = y.$$

Base sizes

Observation: If Δ is a base and $x, y \in G$, then

$$\alpha^x = \alpha^y \text{ for all } \alpha \in \Delta \iff xy^{-1} \in \bigcap_{\alpha \in \Delta} G_\alpha = G_{(\Delta)} \iff x = y.$$

That is, each group element is uniquely determined by its action on Δ .

Base sizes

Observation: If Δ is a base and $x, y \in G$, then

$$\alpha^x = \alpha^y \text{ for all } \alpha \in \Delta \iff xy^{-1} \in \bigcap_{\alpha \in \Delta} G_\alpha = G_{(\Delta)} \iff x = y.$$

That is, each group element is uniquely determined by its action on Δ .

- A small base Δ provides an efficient way to store the elements of G , using $|\Delta|$ -tuples rather than $|\Omega|$ -tuples.

Base sizes

Observation: If Δ is a base and $x, y \in G$, then

$$\alpha^x = \alpha^y \text{ for all } \alpha \in \Delta \iff xy^{-1} \in \bigcap_{\alpha \in \Delta} G_\alpha = G_{(\Delta)} \iff x = y.$$

That is, each group element is uniquely determined by its action on Δ .

- A small base Δ provides an efficient way to store the elements of G , using $|\Delta|$ -tuples rather than $|\Omega|$ -tuples.

Question: How small can a base be?

Base sizes

Observation: If Δ is a base and $x, y \in G$, then

$$\alpha^x = \alpha^y \text{ for all } \alpha \in \Delta \iff xy^{-1} \in \bigcap_{\alpha \in \Delta} G_\alpha = G_{(\Delta)} \iff x = y.$$

That is, each group element is uniquely determined by its action on Δ .

- A small base Δ provides an efficient way to store the elements of G , using $|\Delta|$ -tuples rather than $|\Omega|$ -tuples.

Question: How small can a base be?

Some applications:

- Minimal dimensions
- Generation of finite groups
- Graphs defined on groups

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

G is called **primitive** if G_α is maximal in G .

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

G is called **primitive** if G_α is maximal in G .

Problem. Classify the finite primitive groups G with $b(G) = 2$.

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

G is called **primitive** if G_α is maximal in G .

Problem. Classify the finite primitive groups G with $b(G) = 2$.

Example

Consider the action of $G = D_{2n}$ on $\{1, \dots, n\}$. Then

- $\{1, 2\}$ is a base, so $b(G) = 2$;

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

G is called **primitive** if G_α is maximal in G .

Problem. Classify the finite primitive groups G with $b(G) = 2$.

Example

Consider the action of $G = D_{2n}$ on $\{1, \dots, n\}$. Then

- $\{1, 2\}$ is a base, so $b(G) = 2$;
- G is primitive iff n is a prime.

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: Partial classification in the setting $G_\alpha/Z(G_\alpha)$ is simple.

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: Partial classification in the setting $G_\alpha/Z(G_\alpha)$ is simple.

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: Partial classification in the setting $G_\alpha/Z(G_\alpha)$ is simple.

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: Partial classification in the setting $G_\alpha/Z(G_\alpha)$ is simple.

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- **Burness et al., 2010/11:** T alternating or sporadic ✓

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: Partial classification in the setting $G_\alpha/Z(G_\alpha)$ is simple.

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- **Burness et al., 2010/11:** T alternating or sporadic ✓
- G Lie type: Partial answers

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: Partial classification in the setting $G_\alpha/Z(G_\alpha)$ is simple.

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- **Burness et al., 2010/11:** T alternating or sporadic ✓
- G Lie type: Partial answers

Diagonal and twisted wreath types: Partial results (**Fawcett 2013/22**)

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: Partial classification in the setting $G_\alpha/Z(G_\alpha)$ is simple.

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- **Burness et al., 2010/11:** T alternating or sporadic ✓
- G Lie type: Partial answers

Diagonal and twisted wreath types: Partial results (**Fawcett 2013/22**)

e.g. $G \leq T^k.(\text{Out}(T) \times P)$ with $A_k \not\leq P \implies b(G) = 2$.

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: Partial classification in the setting $G_\alpha/Z(G_\alpha)$ is simple.

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- **Burness et al., 2010/11:** T alternating or sporadic ✓
- G Lie type: Partial answers

Diagonal and twisted wreath types: Partial results (**Fawcett 2013/22**)

e.g. $G \leq T^k.(\text{Out}(T) \times P)$ with $A_k \not\leq P \implies b(G) = 2$.

Product type: $G \leq L \wr P$ with its product action on Γ^k , where $L \leq \text{Sym}(\Gamma)$

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: Partial classification in the setting $G_\alpha/Z(G_\alpha)$ is simple.

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- **Burness et al., 2010/11:** T alternating or sporadic ✓
- G Lie type: Partial answers

Diagonal and twisted wreath types: Partial results (**Fawcett 2013/22**)

e.g. $G \leq T^k.(\text{Out}(T) \times P)$ with $A_k \not\leq P \implies b(G) = 2$.

Product type: $G \leq L \wr P$ with its product action on Γ^k , where $L \leq \text{Sym}(\Gamma)$

- **Bailey & Cameron, 2013:** Explicit $b(L \wr P)$ ✓

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: Partial classification in the setting $G_\alpha/Z(G_\alpha)$ is simple.

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question: Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- **Burness et al., 2010/11:** T alternating or sporadic ✓
- G Lie type: Partial answers

Diagonal and twisted wreath types: Partial results (**Fawcett 2013/22**)

e.g. $G \leq T^k.(\text{Out}(T) \times P)$ with $A_k \not\leq P \implies b(G) = 2$.

Product type: $G \leq L \wr P$ with its product action on Γ^k , where $L \leq \text{Sym}(\Gamma)$

- **Bailey & Cameron, 2013:** Explicit $b(L \wr P)$ ✓
- Progress where $G < L \wr P$ (**Burness & H, 2022+**)

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n . Then

$$Q(G) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_\alpha \cap G_\beta \neq 1\}|}{n^2}$$

is the probability that a random pair in Ω is not a base for G .

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n . Then

$$Q(G) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_\alpha \cap G_\beta \neq 1\}|}{n^2}$$

is the probability that a random pair in Ω is not a base for G .

Note. $Q(G) < 1 \iff b(G) \leq 2$.

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n . Then

$$Q(G) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_\alpha \cap G_\beta \neq 1\}|}{n^2}$$

is the probability that a random pair in Ω is not a base for G .

Note. $Q(G) < 1 \iff b(G) \leq 2$.

Chen & H, 2022: General method to calculate $Q(G)$.

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n . Then

$$Q(G) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_\alpha \cap G_\beta \neq 1\}|}{n^2}$$

is the probability that a random pair in Ω is not a base for G .

Note. $Q(G) < 1 \iff b(G) \leq 2$.

Chen & H, 2022: General method to calculate $Q(G)$.

To calculate exact $Q(G)$ is difficult, but we have

$$Q(G) < \sum_{x \in \mathcal{P}} \frac{|x^G \cap G_\alpha|}{|x^G|} =: \hat{Q}(G),$$

where \mathcal{P} is the set of elements of prime order in G .

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a transitive permutation group of degree n . Then

$$Q(G) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_\alpha \cap G_\beta \neq 1\}|}{n^2}$$

is the probability that a random pair in Ω is not a base for G .

Note. $Q(G) < 1 \iff b(G) \leq 2$.

Chen & H, 2022: General method to calculate $Q(G)$.

To calculate exact $Q(G)$ is difficult, but we have

$$Q(G) < \sum_{x \in \mathcal{P}} \frac{|x^G \cap G_\alpha|}{|x^G|} =: \hat{Q}(G),$$

where \mathcal{P} is the set of elements of prime order in G .

Probabilistic method: $\hat{Q}(G) < 1 \implies b(G) \leq 2$.

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

From now on, we assume G is transitive and $b(G) \leq 2$.

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

From now on, we assume G is transitive and $b(G) \leq 2$.

Examples

- $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$:

Saxl graphs


Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

From now on, we assume G is transitive and $b(G) \leq 2$.

Examples

- $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 

Saxl graphs


Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

From now on, we assume G is transitive and $b(G) \leq 2$.

Examples

- $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 
- $G = D_{10} = \langle (12345), (25)(34) \rangle$, $\Omega = \{1, 2, 3, 4, 5\}$:

Saxl graphs


Definition (Burness & Giudici, 2020)


Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

From now on, we assume G is transitive and $b(G) \leq 2$.

Examples

• $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 

• $G = D_{10} = \langle (12345), (25)(34) \rangle$, $\Omega = \{1, 2, 3, 4, 5\}$: $\Sigma(G) =$ 

Further examples

- $G = \mathrm{GL}_2(q)$ and $\Omega = \mathbb{F}_q^2 \setminus \{0\} \implies b(G) = \dim(\mathbb{F}_q^2) = 2$.

Further examples

- $G = \mathrm{GL}_2(q)$ and $\Omega = \mathbb{F}_q^2 \setminus \{0\} \implies b(G) = \dim(\mathbb{F}_q^2) = 2$.

Note. $\{v, w\}$ is a base $\iff \{v, w\}$ is linearly independent.

Further examples

- $G = \mathrm{GL}_2(q)$ and $\Omega = \mathbb{F}_q^2 \setminus \{0\} \implies b(G) = \dim(\mathbb{F}_q^2) = 2$.

Note. $\{v, w\}$ is a base $\iff \{v, w\}$ is linearly independent.

So $\Sigma(G) = \mathbf{K}_{q^2-1} - (q+1)\mathbf{K}_{q-1}$ is **complete multipartite**.

Further examples

- $G = \mathrm{GL}_2(q)$ and $\Omega = \mathbb{F}_q^2 \setminus \{0\} \implies b(G) = \dim(\mathbb{F}_q^2) = 2$.

Note. $\{v, w\}$ is a base $\iff \{v, w\}$ is linearly independent.

So $\Sigma(G) = \mathbf{K}_{q^2-1} - (q+1)\mathbf{K}_{q-1}$ is **complete multipartite**.

- $G = \mathrm{PGL}_2(q)$ and $\Omega = \{2\text{-subsets of } \{1\text{-spaces in } \mathbb{F}_q^2\}\}$.

Further examples

- $G = \mathrm{GL}_2(q)$ and $\Omega = \mathbb{F}_q^2 \setminus \{0\} \implies b(G) = \dim(\mathbb{F}_q^2) = 2$.

Note. $\{v, w\}$ is a base $\iff \{v, w\}$ is linearly independent.

So $\Sigma(G) = \mathbf{K}_{q^2-1} - (q+1)\mathbf{K}_{q-1}$ is **complete multipartite**.

- $G = \mathrm{PGL}_2(q)$ and $\Omega = \{2\text{-subsets of } \{1\text{-spaces in } \mathbb{F}_q^2\}\}$.

Note. $G_\alpha \cong D_{2(q-1)}$ and $\{\alpha, \beta\}$ is a base $\iff |\alpha \cap \beta| = 1$.

Further examples

- $G = \mathrm{GL}_2(q)$ and $\Omega = \mathbb{F}_q^2 \setminus \{0\} \implies b(G) = \dim(\mathbb{F}_q^2) = 2$.

Note. $\{v, w\}$ is a base $\iff \{v, w\}$ is linearly independent.

So $\Sigma(G) = \mathbf{K}_{q^2-1} - (q+1)\mathbf{K}_{q-1}$ is **complete multipartite**.

- $G = \mathrm{PGL}_2(q)$ and $\Omega = \{2\text{-subsets of } \{1\text{-spaces in } \mathbb{F}_q^2\}\}$.

Note. $G_\alpha \cong D_{2(q-1)}$ and $\{\alpha, \beta\}$ is a base $\iff |\alpha \cap \beta| = 1$.

Hence, $\Sigma(G) \cong J(q+1, 2)$ is a **Johnson graph**.

Further examples

- $G = \mathrm{GL}_2(q)$ and $\Omega = \mathbb{F}_q^2 \setminus \{0\} \implies b(G) = \dim(\mathbb{F}_q^2) = 2$.

Note. $\{v, w\}$ is a base $\iff \{v, w\}$ is linearly independent.


So $\Sigma(G) = \mathbf{K}_{q^2-1} - (q+1)\mathbf{K}_{q-1}$ is **complete multipartite**.

- $G = \mathrm{PGL}_2(q)$ and $\Omega = \{2\text{-subsets of } \{1\text{-spaces in } \mathbb{F}_q^2\}\}$.

Note. $G_\alpha \cong D_{2(q-1)}$ and $\{\alpha, \beta\}$ is a base $\iff |\alpha \cap \beta| = 1$.

Hence, $\Sigma(G) \cong J(q+1, 2)$ is a **Johnson graph**.

For example, when $q = 4$ we have the complement of the Petersen.



Diameter

Notes.

- $\Sigma(G)$ is the union of the **regular orbital graphs** of G .

Diameter

Notes.

- $\Sigma(G)$ is the union of the **regular orbital graphs** of G .
- G is primitive $\implies \Sigma(G)$ is connected.

Diameter

Notes.

- $\Sigma(G)$ is the union of the **regular orbital graphs** of G .
- G is primitive $\implies \Sigma(G)$ is connected.

Question: What is the diameter of $\Sigma(G)$ if G is primitive?

Diameter

Notes.

- $\Sigma(G)$ is the union of the **regular orbital graphs** of G .
- G is primitive $\implies \Sigma(G)$ is connected.

Question: What is the diameter of $\Sigma(G)$ if G is primitive?

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Diameter

Notes.

- $\Sigma(G)$ is the union of the **regular orbital graphs** of G .
- G is primitive $\implies \Sigma(G)$ is connected.

Question: What is the diameter of $\Sigma(G)$ if G is primitive?

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

In particular, it asserts that $\Sigma(G)$ has diameter at most 2.

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓
- “Most” almost simple groups with alternating or sporadic socles

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓
- “Most” almost simple groups with alternating or sporadic socles
- **Chen & Du, 2020+; Burness & H, 2021+:** $\text{soc}(G) = L_2(q)$ ✓

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓
- “Most” almost simple groups with alternating or sporadic socles
- **Chen & Du, 2020+; Burness & H, 2021+:** $\text{soc}(G) = L_2(q)$ ✓
- **Burness & H, 2021+:** almost simple groups with G_α soluble ✓

Diameter

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓
- “Most” almost simple groups with alternating or sporadic socles
- **Chen & Du, 2020+; Burness & H, 2021+:** $\text{soc}(G) = L_2(q)$ ✓
- **Burness & H, 2021+:** almost simple groups with G_α soluble ✓
- **Lee & Popiel, 2021+:** some affine groups

Probabilistic methods

Let $v(G)$ be the valency of $\Sigma(G)$ and recall that

$$Q(G) = 1 - \frac{v(G)}{|\Omega|}$$

is the probability that a random pair in Ω is not a base for G .

Probabilistic methods

Let $v(G)$ be the valency of $\Sigma(G)$ and recall that

$$Q(G) = 1 - \frac{v(G)}{|\Omega|}$$

is the probability that a random pair in Ω is not a base for G .

Notes.

- $Q(G) < 1 \iff b(G) \leq 2$.

Probabilistic methods

Let $v(G)$ be the valency of $\Sigma(G)$ and recall that

$$Q(G) = 1 - \frac{v(G)}{|\Omega|}$$

is the probability that a random pair in Ω is not a base for G .

Notes.

- $Q(G) < 1 \iff b(G) \leq 2$.
- $Q(G) < 1/2 \implies \Sigma(G)$ has the common neighbour property.

Probabilistic methods

Let $v(G)$ be the valency of $\Sigma(G)$ and recall that

$$Q(G) = 1 - \frac{v(G)}{|\Omega|}$$

is the probability that a random pair in Ω is not a base for G .

Notes.

- $Q(G) < 1 \iff b(G) \leq 2$.
- $Q(G) < 1/2 \implies \Sigma(G)$ has the common neighbour property.
- $Q(G) < 1/t \implies \Sigma(G)$ has clique number at least $t + 1$.

Probabilistic methods

Let $v(G)$ be the valency of $\Sigma(G)$ and recall that

$$Q(G) = 1 - \frac{v(G)}{|\Omega|}$$

is the probability that a random pair in Ω is not a base for G .

Notes.

- $Q(G) < 1 \iff b(G) \leq 2$.
- $Q(G) < 1/2 \implies \Sigma(G)$ has the common neighbour property.
- $Q(G) < 1/t \implies \Sigma(G)$ has clique number at least $t + 1$.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $Q(G) \rightarrow 1$ as $q \rightarrow \infty$. But $\Sigma(G) = J(q+1, 2)$ still has the common neighbour property.

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Note. $\Sigma(\alpha)$ is the union of regular G_α -orbits.

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Note. $\Sigma(\alpha)$ is the union of regular G_α -orbits.

Recall: BG conjecture: $\Sigma(\alpha)$ meets the union of regular G_β -orbits.

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Note. $\Sigma(\alpha)$ is the union of regular G_α -orbits.

Recall: BG conjecture: $\Sigma(\alpha)$ meets the union of regular G_β -orbits.

Conjecture (Burness & H, 2022+)

G primitive and $\alpha, \beta \in \Omega \implies \Sigma(\alpha)$ meets **every** regular G_β -orbit.

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Note. $\Sigma(\alpha)$ is the union of regular G_α -orbits.

Recall: BG conjecture: $\Sigma(\alpha)$ meets the union of regular G_β -orbits.

Conjecture (Burness & H, 2022+)

G primitive and $\alpha, \beta \in \Omega \implies \Sigma(\alpha)$ meets **every** regular G_β -orbit.

Theorem (Burness & H, 2022+)

BG conjecture and BH conjecture are equivalent.

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Note. $\Sigma(\alpha)$ is the union of regular G_α -orbits.

Recall: BG conjecture: $\Sigma(\alpha)$ meets the union of regular G_β -orbits.

Conjecture (Burness & H, 2022+)

G primitive and $\alpha, \beta \in \Omega \implies \Sigma(\alpha)$ meets **every** regular G_β -orbit.

Theorem (Burness & H, 2022+)

BG conjecture and BH conjecture are equivalent.

Evidence:

- All primitive groups of degree up to 4095 ✓

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Note. $\Sigma(\alpha)$ is the union of regular G_α -orbits.

Recall: BG conjecture: $\Sigma(\alpha)$ meets the union of regular G_β -orbits.

Conjecture (Burness & H, 2022+)

G primitive and $\alpha, \beta \in \Omega \implies \Sigma(\alpha)$ meets **every** regular G_β -orbit.

Theorem (Burness & H, 2022+)

BG conjecture and BH conjecture are equivalent.

Evidence:

- All primitive groups of degree up to 4095 ✓
- $G = \text{PSL}_2(q)$ and G_α of type $\text{GL}_1(q) \wr S_2$

Future work

Saxl graphs:

- Generalised Saxl graphs

Future work

Saxl graphs:

- Generalised Saxl graphs
- Valencies (**Chen & H, 2022**: some results)

Future work

Saxl graphs:

- Generalised Saxl graphs
- Valencies (**Chen & H, 2022**: some results)
- Other invariants of Saxl graphs (e.g. the independence number)

Future work

Saxl graphs:

- Generalised Saxl graphs
- Valencies (**Chen & H, 2022**: some results)
- Other invariants of Saxl graphs (e.g. the independence number)

Other problems on bases:

- Irredundant bases

Future work

Saxl graphs:

- Generalised Saxl graphs
- Valencies (**Chen & H, 2022**: some results)
- Other invariants of Saxl graphs (e.g. the independence number)

Other problems on bases:

- Irredundant bases
- Bounds for $b(G)$ in a general setting

Thank you!