

Groups, graphs and transitivity

Hong Yi Huang

University of Bristol

JAC, 21 May 2021

Outline

1 Automorphisms and transitivity

2 Cayley graphs

3 Vertex-transitive graphs

4 Asymptotic problems

5 Saxl graphs

Automorphisms

Let $\Gamma = (V, E)$ be a finite simple digraph: no loops or multiple edges.

Automorphisms

Let $\Gamma = (V, E)$ be a finite simple digraph: no loops or multiple edges.

Automorphism: $\sigma \in \text{Sym}(V)$ such that $\alpha^\sigma \rightarrow \beta^\sigma \iff \alpha \rightarrow \beta$.

Automorphisms

Let $\Gamma = (V, E)$ be a finite simple digraph: no loops or multiple edges.

Automorphism: $\sigma \in \text{Sym}(V)$ such that $\alpha^\sigma \rightarrow \beta^\sigma \iff \alpha \rightarrow \beta$.

Automorphism group $\text{Aut}(\Gamma)$: the set of automorphisms.

Automorphisms

Let $\Gamma = (V, E)$ be a finite simple digraph: no loops or multiple edges.

Automorphism: $\sigma \in \text{Sym}(V)$ such that $\alpha^\sigma \rightarrow \beta^\sigma \iff \alpha \rightarrow \beta$.

Automorphism group $\text{Aut}(\Gamma)$: the set of automorphisms.

Example

- $\text{Aut}(\mathbf{P}_n) \cong C_2$, where \mathbf{P}_n is a path of length n .

Automorphisms

Let $\Gamma = (V, E)$ be a finite simple digraph: no loops or multiple edges.

Automorphism: $\sigma \in \text{Sym}(V)$ such that $\alpha^\sigma \rightarrow \beta^\sigma \iff \alpha \rightarrow \beta$.

Automorphism group $\text{Aut}(\Gamma)$: the set of automorphisms.

Example

- $\text{Aut}(\mathbf{P}_n) \cong C_2$, where \mathbf{P}_n is a path of length n .
- $\text{Aut}(\mathbf{C}_n) \cong D_{2n}$, where \mathbf{C}_n is a cycle of length n .

Automorphisms

Let $\Gamma = (V, E)$ be a finite simple digraph: no loops or multiple edges.

Automorphism: $\sigma \in \text{Sym}(V)$ such that $\alpha^\sigma \rightarrow \beta^\sigma \iff \alpha \rightarrow \beta$.

Automorphism group $\text{Aut}(\Gamma)$: the set of automorphisms.

Example

- $\text{Aut}(\mathbf{P}_n) \cong C_2$, where \mathbf{P}_n is a path of length n .
- $\text{Aut}(\mathbf{C}_n) \cong D_{2n}$, where \mathbf{C}_n is a cycle of length n .
- $\text{Aut}(\mathbf{K}_n) \cong S_n$, where \mathbf{K}_n is a complete graph of order n .

Automorphisms

Let $\Gamma = (V, E)$ be a finite simple digraph: no loops or multiple edges.

Automorphism: $\sigma \in \text{Sym}(V)$ such that $\alpha^\sigma \rightarrow \beta^\sigma \iff \alpha \rightarrow \beta$.

Automorphism group $\text{Aut}(\Gamma)$: the set of automorphisms.

Example

- $\text{Aut}(\mathbf{P}_n) \cong C_2$, where \mathbf{P}_n is a path of length n .
- $\text{Aut}(\mathbf{C}_n) \cong D_{2n}$, where \mathbf{C}_n is a cycle of length n .
- $\text{Aut}(\mathbf{K}_n) \cong S_n$, where \mathbf{K}_n is a complete graph of order n .

Γ is called **vertex-transitive** if $\text{Aut}(\Gamma)$ is transitive on the set of vertices.

Automorphisms

Let $\Gamma = (V, E)$ be a finite simple digraph: no loops or multiple edges.

Automorphism: $\sigma \in \text{Sym}(V)$ such that $\alpha^\sigma \rightarrow \beta^\sigma \iff \alpha \rightarrow \beta$.

Automorphism group $\text{Aut}(\Gamma)$: the set of automorphisms.

Example

- $\text{Aut}(\mathbf{P}_n) \cong C_2$, where \mathbf{P}_n is a path of length n .
- $\text{Aut}(\mathbf{C}_n) \cong D_{2n}$, where \mathbf{C}_n is a cycle of length n .
- $\text{Aut}(\mathbf{K}_n) \cong S_n$, where \mathbf{K}_n is a complete graph of order n .

Γ is called **vertex-transitive** if $\text{Aut}(\Gamma)$ is transitive on the set of vertices.

Remark: one can replace “vertex” and “transitive” by almost anything. For example: edge-primitive or arc-semiregular.

The Petersen graph

The Petersen graph Γ can be constructed as follows:

- Vertices: 2-subsets of $\{1, 2, 3, 4, 5\}$;
- Edges: $\alpha \sim \beta$ if $\alpha \cap \beta = \emptyset$.

The Petersen graph

The Petersen graph Γ can be constructed as follows:

- Vertices: 2-subsets of $\{1, 2, 3, 4, 5\}$;
- Edges: $\alpha \sim \beta$ if $\alpha \cap \beta = \emptyset$.


One can show that $\text{Aut}(\Gamma) \cong S_5$.

The Petersen graph

The Petersen graph Γ can be constructed as follows:

- Vertices: 2-subsets of $\{1, 2, 3, 4, 5\}$;
- Edges: $\alpha \sim \beta$ if $\alpha \cap \beta = \emptyset$.

One can show that $\text{Aut}(\Gamma) \cong S_5$.



Outline

- 1 Automorphisms and transitivity
- 2 Cayley graphs
- 3 Vertex-transitive graphs
- 4 Asymptotic problems
- 5 Saxl graphs

Basic properties

Let G be a finite group and $S \subseteq G \setminus \{1\}$.

Cayley digraph $\text{Cay}(G, S)$: vertices G and $x \rightarrow y \iff yx^{-1} \in S$.

Basic properties

Let G be a finite group and $S \subseteq G \setminus \{1\}$.

Cayley digraph $\text{Cay}(G, S)$: vertices G and $x \rightarrow y \iff yx^{-1} \in S$.

- A Cayley digraph $\text{Cay}(G, S)$ is undirected $\iff S = S^{-1}$.

Basic properties

Let G be a finite group and $S \subseteq G \setminus \{1\}$.

Cayley digraph $\text{Cay}(G, S)$: vertices G and $x \rightarrow y \iff yx^{-1} \in S$.

- A Cayley digraph $\text{Cay}(G, S)$ is undirected $\iff S = S^{-1}$.

Cayley graph $\text{Cay}(G, S)$: undirected Cayley digraph (or $S = S^{-1}$).

Basic properties

Let G be a finite group and $S \subseteq G \setminus \{1\}$.

Cayley digraph $\text{Cay}(G, S)$: vertices G and $x \rightarrow y \iff yx^{-1} \in S$.

- A Cayley digraph $\text{Cay}(G, S)$ is undirected $\iff S = S^{-1}$.

Cayley graph $\text{Cay}(G, S)$: undirected Cayley digraph (or $S = S^{-1}$).

- A Cayley (di)graph is connected $\iff G = \langle S \rangle$.

Basic properties

Let G be a finite group and $S \subseteq G \setminus \{1\}$.

Cayley digraph $\text{Cay}(G, S)$: vertices G and $x \rightarrow y \iff yx^{-1} \in S$.

- A Cayley digraph $\text{Cay}(G, S)$ is undirected $\iff S = S^{-1}$.

Cayley graph $\text{Cay}(G, S)$: undirected Cayley digraph (or $S = S^{-1}$).

- A Cayley (di)graph is connected $\iff G = \langle S \rangle$.

Indeed, $1 \rightarrow s_1 \rightarrow s_1 s_2 \rightarrow \cdots \rightarrow s_1 s_2 \cdots s_n$ for $s_i \in S$ is a path.

Some examples

Example

- $\text{Cay}(G, G \setminus \{1\}) \cong \mathbf{K}_n$ if $|G| = n$.

Some examples

Example

- $\text{Cay}(G, G \setminus \{1\}) \cong \mathbf{K}_n$ if $|G| = n$.
- $\text{Cay}(G, G \setminus H) \cong \mathbf{K}_{n,n}$ if $|G| = 2n$ with $|G : H| = 2$, where $\mathbf{K}_{n,n}$ denotes the complete bipartite graph.

Some examples

Example

- $\text{Cay}(G, G \setminus \{1\}) \cong \mathbf{K}_n$ if $|G| = n$.
- $\text{Cay}(G, G \setminus H) \cong \mathbf{K}_{n,n}$ if $|G| = 2n$ with $|G : H| = 2$, where $\mathbf{K}_{n,n}$ denotes the complete bipartite graph.
- $\text{Cay}(G, \{x, x^{-1}\}) \cong \mathbf{C}_n$ and $\text{Cay}(G, \{x\}) \cong \overrightarrow{\mathbf{C}}_n$ if $G = \langle x \rangle \cong C_n$.

Some examples

Example

- $\text{Cay}(G, G \setminus \{1\}) \cong \mathbf{K}_n$ if $|G| = n$.
- $\text{Cay}(G, G \setminus H) \cong \mathbf{K}_{n,n}$ if $|G| = 2n$ with $|G : H| = 2$, where $\mathbf{K}_{n,n}$ denotes the complete bipartite graph.
- $\text{Cay}(G, \{x, x^{-1}\}) \cong \mathbf{C}_n$ and $\text{Cay}(G, \{x\}) \cong \overrightarrow{\mathbf{C}}_n$ if $G = \langle x \rangle \cong C_n$.
- $\text{Cay}(G, \{x, x^{-1}\}) \cong m\mathbf{C}_n$ if $|G| = mn$ and $|x| = n$.

Transitivity

Lemma

Γ is a Cayley digraph $\iff \text{Aut}(\Gamma)$ has a regular subgroup.


Transitivity

Lemma

Γ is a Cayley digraph $\iff \text{Aut}(\Gamma)$ has a regular subgroup.

Note that a Cayley graph is vertex-transitive. The converse is false!

The smallest example: the Petersen graph.




Transitivity

Lemma

Γ is a Cayley digraph $\iff \text{Aut}(\Gamma)$ has a regular subgroup.

Note that a Cayley graph is vertex-transitive. The converse is false!

The smallest example: the Petersen graph.



How can we use groups to construct non-Cayley vertex-transitive graphs?

Outline

- 1 Automorphisms and transitivity
- 2 Cayley graphs
- 3 Vertex-transitive graphs**
- 4 Asymptotic problems
- 5 Saxl graphs

Coset graphs

Let G be a finite group, $H < G$ and $S \subseteq G \setminus H$.

Coset digraph $\text{Cos}(G, H, HSH)$: vertices $[G : H]$ and

$$xH \rightarrow yH \iff yx^{-1} \in HSH.$$

Coset graphs

Let G be a finite group, $H < G$ and $S \subseteq G \setminus H$.

Coset digraph $\text{Cos}(G, H, HSH)$: vertices $[G : H]$ and

$$xH \rightarrow yH \iff yx^{-1} \in HSH.$$

A Cayley digraph is a coset digraph by taking $H = 1$.

Coset graphs

Let G be a finite group, $H < G$ and $S \subseteq G \setminus H$.

Coset digraph $\text{Cos}(G, H, HSH)$: vertices $[G : H]$ and

$$xH \rightarrow yH \iff yx^{-1} \in HSH.$$

A Cayley digraph is a coset digraph by taking $H = 1$.

- A coset digraph $\text{Cos}(G, H, HSH)$ is undirected $\iff HSH = HS^{-1}H$.

Coset graphs

Let G be a finite group, $H < G$ and $S \subseteq G \setminus H$.

Coset digraph $\text{Cos}(G, H, HSH)$: vertices $[G : H]$ and

$$xH \rightarrow yH \iff yx^{-1} \in HSH.$$

A Cayley digraph is a coset digraph by taking $H = 1$.

- A coset digraph $\text{Cos}(G, H, HSH)$ is undirected $\iff HSH = HS^{-1}H$.

Coset graph $\text{Cos}(G, H, HSH)$: undirected coset digraph (or $HSH = HS^{-1}H$).

Coset graphs

Let G be a finite group, $H < G$ and $S \subseteq G \setminus H$.

Coset digraph $\text{Cos}(G, H, HSH)$: vertices $[G : H]$ and

$$xH \rightarrow yH \iff yx^{-1} \in HSH.$$

A Cayley digraph is a coset digraph by taking $H = 1$.

- A coset digraph $\text{Cos}(G, H, HSH)$ is undirected $\iff HSH = HS^{-1}H$.

Coset graph $\text{Cos}(G, H, HSH)$: undirected coset digraph (or $HSH = HS^{-1}H$).

- A coset graph $\text{Cos}(G, H, HSH)$ is connected $\iff \langle H, S \rangle = G$.

Coset graphs

Let G be a finite group, $H < G$ and $S \subseteq G \setminus H$.

Coset digraph $\text{Cos}(G, H, HSH)$: vertices $[G : H]$ and

$$xH \rightarrow yH \iff yx^{-1} \in HSH.$$

A Cayley digraph is a coset digraph by taking $H = 1$.

- A coset digraph $\text{Cos}(G, H, HSH)$ is undirected $\iff HSH = HS^{-1}H$.


Coset graph $\text{Cos}(G, H, HSH)$: undirected coset digraph (or $HSH = HS^{-1}H$).

- A coset graph $\text{Cos}(G, H, HSH)$ is connected $\iff \langle H, S \rangle = G$.

Lemma

Γ is vertex-transitive $\iff \Gamma$ is a coset digraph.

Petersen graph as a coset graph



$\Gamma = \text{Cos}(S_5, H, HSH)$ with $H = \text{Sym}(\{1, 2\}) \times \text{Sym}(\{3, 4, 5\})$ and
 $S = \{(13)(24), (13)(25), (14)(25)\}.$

Arc-transitivity

Γ is called **arc-transitive** if $\text{Aut}(\Gamma)$ acts transitively on arcs of Γ .

Lemma

Γ is arc-transitive $\iff \Gamma$ is isomorphic to $\text{Cos}(G, H, HgH)$ for some G , H and g .

Arc-transitivity

Γ is called **arc-transitive** if $\text{Aut}(\Gamma)$ acts transitively on arcs of Γ .

Lemma

Γ is arc-transitive $\iff \Gamma$ is isomorphic to $\text{Cos}(G, H, HgH)$ for some G , H and g .

The Petersen graph Γ :

$\Gamma = \text{Cos}(S_5, H, HSH)$ with $H = \text{Sym}(\{1, 2\}) \times \text{Sym}(\{3, 4, 5\})$ and

$$S = \{(13)(24), (13)(25), (14)(25)\}.$$

Arc-transitivity

Γ is called **arc-transitive** if $\text{Aut}(\Gamma)$ acts transitively on arcs of Γ .

Lemma

Γ is arc-transitive $\iff \Gamma$ is isomorphic to $\text{Cos}(G, H, HgH)$ for some G , H and g .

The Petersen graph Γ :

$\Gamma = \text{Cos}(S_5, H, HSH)$ with $H = \text{Sym}(\{1, 2\}) \times \text{Sym}(\{3, 4, 5\})$ and

$$S = \{(13)(24), (13)(25), (14)(25)\}.$$

$\Gamma = \text{Cos}(S_5, H, HgH)$ with $H = \text{Sym}(\{1, 2\}) \times \text{Sym}(\{3, 4, 5\})$ and

$$S = \{g\} = \{(13)(24)\}.$$

Orbital graphs

Let $G \leq \text{Sym}(\Omega)$ be a permutation group.

Orbital graphs

Let $G \leq \text{Sym}(\Omega)$ be a permutation group.

An **orbital** of G on Ω is an orbit of G on $\Omega \times \Omega$.

Orbital graphs

Let $G \leq \text{Sym}(\Omega)$ be a permutation group.

An **orbital** of G on Ω is an orbit of G on $\Omega \times \Omega$.

For example, if $x, y \in \Omega$, then $\{(x^g, y^g) : g \in G\}$ is an orbital.

Orbital graphs

Let $G \leq \text{Sym}(\Omega)$ be a permutation group.

An **orbital** of G on Ω is an orbit of G on $\Omega \times \Omega$.

For example, if $x, y \in \Omega$, then $\{(x^g, y^g) : g \in G\}$ is an orbital.

If G is transitive, then $\{(x, x) : x \in \Omega\}$ is an orbital (diagonal).

Orbital graphs

Let $G \leq \text{Sym}(\Omega)$ be a permutation group.

An **orbital** of G on Ω is an orbit of G on $\Omega \times \Omega$.

For example, if $x, y \in \Omega$, then $\{(x^g, y^g) : g \in G\}$ is an orbital.

If G is transitive, then $\{(x, x) : x \in \Omega\}$ is an orbital (diagonal).

G is 2-transitive iff G has exactly 2 orbitals.

Orbital graphs

Let $G \leq \text{Sym}(\Omega)$ be a permutation group.

An **orbital** of G on Ω is an orbit of G on $\Omega \times \Omega$.

For example, if $x, y \in \Omega$, then $\{(x^g, y^g) : g \in G\}$ is an orbital.

If G is transitive, then $\{(x, x) : x \in \Omega\}$ is an orbital (diagonal).

G is 2-transitive iff G has exactly 2 orbitals.

The **orbital graph** $(x, y)^G$ of G is a graph with

- Vertices Ω ;
- Edges in the orbital $(x, y)^G = \{(x^g, y^g) : g \in G\}$ (non-diagonal).

Orbital graphs

Example

- \mathbf{K}_n is an orbital graph for S_n ;
- \mathbf{K}_n is an orbital graph for a 2-transitive group of degree n ;

Orbital graphs

Example

- \mathbf{K}_n is an orbital graph for S_n ;
- \mathbf{K}_n is an orbital graph for a 2-transitive group of degree n ;
- Let $G = \langle (012345) \rangle \cong C_6$ acting on $\Omega = \{0, 1, 2, 3, 4, 5\}$.
 - ▶ $(0, 1)^G$ and $(1, 0)^G$ are isomorphic to $\overrightarrow{\mathbf{C}_6}$;
 - ▶ $(0, 2)^G$ and $(2, 0)^G$ are isomorphic to $2\overrightarrow{\mathbf{C}_3}$;
 - ▶ $(0, 3)^G = (3, 0)^G$ is isomorphic to $3\mathbf{K}_2$;

Orbital graphs

Example

- \mathbf{K}_n is an orbital graph for S_n ;
- \mathbf{K}_n is an orbital graph for a 2-transitive group of degree n ;
- Let $G = \langle (012345) \rangle \cong C_6$ acting on $\Omega = \{0, 1, 2, 3, 4, 5\}$.
 - ▶ $(0, 1)^G$ and $(1, 0)^G$ are isomorphic to $\overrightarrow{\mathbf{C}_6}$;
 - ▶ $(0, 2)^G$ and $(2, 0)^G$ are isomorphic to $2\overrightarrow{\mathbf{C}_3}$;
 - ▶ $(0, 3)^G = (3, 0)^G$ is isomorphic to $3\mathbf{K}_2$;
- Let $G = \langle (012345), (15)(32) \rangle \cong D_{12}$ acting on $\Omega = \{0, 1, 2, 3, 4, 5\}$.
 - ▶ $(0, 1)^G = (1, 0)^G$ is isomorphic to \mathbf{C}_6 ;
 - ▶ $(0, 2)^G = (2, 0)^G$ is isomorphic to $2\mathbf{C}_3$;
 - ▶ $(0, 3)^G = (3, 0)^G$ is isomorphic to $3\mathbf{K}_2$;

Orbital graphs

Example

- \mathbf{K}_n is an orbital graph for S_n ;
- \mathbf{K}_n is an orbital graph for a 2-transitive group of degree n ;
- Let $G = \langle (012345) \rangle \cong C_6$ acting on $\Omega = \{0, 1, 2, 3, 4, 5\}$.
 - ▶ $(0, 1)^G$ and $(1, 0)^G$ are isomorphic to $\overrightarrow{\mathbf{C}_6}$;
 - ▶ $(0, 2)^G$ and $(2, 0)^G$ are isomorphic to $2\overrightarrow{\mathbf{C}_3}$;
 - ▶ $(0, 3)^G = (3, 0)^G$ is isomorphic to $3\mathbf{K}_2$;
- Let $G = \langle (012345), (15)(32) \rangle \cong D_{12}$ acting on $\Omega = \{0, 1, 2, 3, 4, 5\}$.
 - ▶ $(0, 1)^G = (1, 0)^G$ is isomorphic to \mathbf{C}_6 ;
 - ▶ $(0, 2)^G = (2, 0)^G$ is isomorphic to $2\mathbf{C}_3$;
 - ▶ $(0, 3)^G = (3, 0)^G$ is isomorphic to $3\mathbf{K}_2$;

An orbital Δ is called **self-paired** if $(x, y) \in \Delta \iff (y, x) \in \Delta$.

An orbital graph is undirected iff the orbital is self-paired.

Arc-transitivity

Lemma


Γ is arc-transitive $\iff \Gamma$ is an orbital graph for a transitive group.

Arc-transitivity

Lemma

Γ is arc-transitive $\iff \Gamma$ is an orbital graph for a transitive group.

The Petersen graph $\Gamma = (\{1, 2\}, \{3, 4\})^G$ for $G = \text{Sym}(5)$.



Generalised orbital graph

A **generalised orbital graph** for G is a union of orbital graphs for G .

Generalised orbital graph

A **generalised orbital graph** for G is a union of orbital graphs for G .

Lemma

Γ is vertex-transitive $\iff \Gamma$ is a generalised orbital graph for a transitive group.

Summary

Lemma

Let Γ be a finite simple digraph. Then Γ is Cayley $\iff \text{Aut}(\Gamma)$ has a regular subgroup.

Summary

Lemma

Let Γ be a finite simple digraph. Then Γ is Cayley \iff $\text{Aut}(\Gamma)$ has a regular subgroup.

Lemma

Let Γ be a finite simple digraph. Then the following are equivalent:

- Γ is vertex-transitive;
- Γ is a generalised orbital graph for a transitive group;
- $\Gamma = \text{Cos}(G, H, HSH)$ for some G , H and S .

Summary

Lemma

Let Γ be a finite simple digraph. Then Γ is Cayley \iff $\text{Aut}(\Gamma)$ has a regular subgroup.

Lemma

Let Γ be a finite simple digraph. Then the following are equivalent:

- Γ is vertex-transitive;
- Γ is a generalised orbital graph for a transitive group;
- $\Gamma = \text{Cos}(G, H, HSH)$ for some G, H and S .

Lemma

Let Γ be a finite simple digraph. Then the following are equivalent:

- Γ is arc-transitive;
- Γ is an orbital graph for a transitive group;
- $\Gamma = \text{Cos}(G, H, HgH)$ for some G, H and g .

Outline

- 1 Automorphisms and transitivity
- 2 Cayley graphs
- 3 Vertex-transitive graphs
- 4 Asymptotic problems
- 5 Saxl graphs

Asymptotic problems

Conjecture

Almost all vertex-transitive (di)graph are Cayley (di)graphs.

Asymptotic problems

Conjecture

Almost all vertex-transitive (di)graph are Cayley (di)graphs.

A graph is called **Hamiltonian** if it has a cycle visiting every vertex exactly once (**Hamiltonian cycle**).

Conjecture (Lovász, 1969)

- All but 5 connected vertex-transitive graphs are Hamiltonian;
- All connected Cayley graphs are Hamiltonian (except K_2).

Asymptotic problems

Conjecture

Almost all vertex-transitive (di)graph are Cayley (di)graphs.

A graph is called **Hamiltonian** if it has a cycle visiting every vertex exactly once (**Hamiltonian cycle**).

Conjecture (Lovász, 1969)

- All but 5 connected vertex-transitive graphs are Hamiltonian;
- All connected Cayley graphs are Hamiltonian (except K_2).

Theorem (Meng & Huang, 1996)

- Almost all Cayley (di)graphs are connected.
- Almost all Cayley graphs are Hamiltonian.

Outline

- 1 Automorphisms and transitivity
- 2 Cayley graphs
- 3 Vertex-transitive graphs
- 4 Asymptotic problems
- 5 Saxl graphs

Bases

Let $G \leq \text{Sym}(\Omega)$ be a permutation group with $|\Omega| < \infty$.

Definition

A subset Δ of Ω is a **base** for G if $\bigcap_{\alpha \in \Delta} G_\alpha = 1$.

Bases

Let $G \leq \text{Sym}(\Omega)$ be a permutation group with $|\Omega| < \infty$.

Definition

A subset Δ of Ω is a **base** for G if $\cap_{\alpha \in \Delta} G_\alpha = 1$.

The **base size** of G , denoted by $b(G)$, is the minimal cardinality of a base.

Bases

Let $G \leq \text{Sym}(\Omega)$ be a permutation group with $|\Omega| < \infty$.

Definition

A subset Δ of Ω is a **base** for G if $\cap_{\alpha \in \Delta} G_\alpha = 1$.

The **base size** of G , denoted by $b(G)$, is the minimal cardinality of a base.

- Images of a base determine the whole group G .

Bases

Let $G \leq \text{Sym}(\Omega)$ be a permutation group with $|\Omega| < \infty$.

Definition

A subset Δ of Ω is a **base** for G if $\bigcap_{\alpha \in \Delta} G_\alpha = 1$.

The **base size** of G , denoted by $b(G)$, is the minimal cardinality of a base.

- Images of a base determine the whole group G .
- If G is transitive and $H = G_\alpha$, then $b(G)$ is the minimal cardinality of a subset $S \subseteq G$ such that

$$\bigcap_{x \in S} H^x = 1.$$

Bases

Let $G \leq \text{Sym}(\Omega)$ be a permutation group with $|\Omega| < \infty$.

Definition

A subset Δ of Ω is a **base** for G if $\bigcap_{\alpha \in \Delta} G_\alpha = 1$.

The **base size** of G , denoted by $b(G)$, is the minimal cardinality of a base.

- Images of a base determine the whole group G .
- If G is transitive and $H = G_\alpha$, then $b(G)$ is the minimal cardinality of a subset $S \subseteq G$ such that

$$\bigcap_{x \in S} H^x = 1.$$

- There always exists a base by noting that Ω is a base.

Bases

Let $G \leq \text{Sym}(\Omega)$ be a permutation group with $|\Omega| < \infty$.

Definition

A subset Δ of Ω is a **base** for G if $\bigcap_{\alpha \in \Delta} G_\alpha = 1$.

The **base size** of G , denoted by $b(G)$, is the minimal cardinality of a base.

- Images of a base determine the whole group G .
- If G is transitive and $H = G_\alpha$, then $b(G)$ is the minimal cardinality of a subset $S \subseteq \Omega$ such that

$$\bigcap_{x \in S} H^x = 1.$$

- There always exists a base by noting that Ω is a base.
- $b(G) = 1 \iff G$ has a regular orbit on Ω .

Examples

Example

- $G = S_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 1$.

Examples

Example

- $G = S_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 1$.
- $G = A_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 2$.

Examples

Example

- $G = S_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 1$.
- $G = A_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 2$.
- $G = D_{2n}$, $\Omega = \{1, \dots, n\}$: $b(G) = 2$.

Examples

Example

- $G = S_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 1$.
- $G = A_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 2$.
- $G = D_{2n}$, $\Omega = \{1, \dots, n\}$: $b(G) = 2$.
- $G = \text{GL}(V)$, $\Omega = V$:
A subset of Ω is a base iff it contains a basis of V , so $b(G) = \dim V$.

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$ be a permutation group with $b(G) = 2$.

The **Saxl graph** $\Sigma(G)$: vertices Ω , $\alpha \sim \beta \iff \{\alpha, \beta\}$ is a base.

Saxl graphs


Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$ be a permutation group with $b(G) = 2$.

The **Saxl graph** $\Sigma(G)$: vertices Ω , $\alpha \sim \beta \iff \{\alpha, \beta\}$ is a base.

Example

- $G = D_8$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) \cong C_4$.



Saxl graphs


Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$ be a permutation group with $b(G) = 2$.

The **Saxl graph** $\Sigma(G)$: vertices Ω , $\alpha \sim \beta \iff \{\alpha, \beta\}$ is a base.

Example

- $G = D_8$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) \cong C_4$.



- $G = D_{10}$, $\Omega = \{1, 2, 3, 4, 5\}$: $\Sigma(G) \cong K_5$.


A further example

Let $G = \text{GL}_2(q)$ and $\Omega = \mathbb{F}_q^2 \setminus \{0\}$. Then $\alpha \sim \beta$ iff $\{\alpha, \beta\}$ is linearly independent. Thus, $\Sigma(G)$ is **complete multipartite** with $q + 1$ parts of size $q - 1$.

A further example

Let $G = \text{GL}_2(q)$ and $\Omega = \mathbb{F}_q^2 \setminus \{0\}$. Then $\alpha \sim \beta$ iff $\{\alpha, \beta\}$ is linearly independent. Thus, $\Sigma(G)$ is **complete multipartite** with $q + 1$ parts of size $q - 1$.

For example, when $q = 3$ we have $\Sigma(G) \cong K_8 - 4K_2$.



Not Petersen this time

Let $G = \text{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 . Then α and β form a base iff they share a common 1-space.

Not Petersen this time

Let $G = \text{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 . Then α and β form a base iff they share a common 1-space.


Hence, $\Sigma(G) \cong J(q+1, 2)$ is a **Johnson graph**: vertices 2-subsets of $\{1, \dots, q+1\}$ and being adjacent if not disjoint.

Not Petersen this time

Let $G = \text{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 . Then α and β form a base iff they share a common 1-space.

Hence, $\Sigma(G) \cong J(q+1, 2)$ is a **Johnson graph**: vertices 2-subsets of $\{1, \dots, q+1\}$ and being adjacent if not disjoint.

For example, when $q = 4$ we have the complement of the Petersen graph.



First observations

Proposition

Suppose G is transitive with $b(G) = 2$ and $\Sigma(G)$ is the Saxl graph of G .

1. $\Sigma(G)$ is G -vertex-transitive.

First observations

Proposition

Suppose G is transitive with $b(G) = 2$ and $\Sigma(G)$ is the Saxl graph of G .

- 1 $\Sigma(G)$ is G -vertex-transitive.
- 2 $\Sigma(G)$ is connected if G is primitive.

First observations

Proposition

Suppose G is transitive with $b(G) = 2$ and $\Sigma(G)$ is the Saxl graph of G .

- 1 $\Sigma(G)$ is G -vertex-transitive.
- 2 $\Sigma(G)$ is connected if G is primitive.
- 3 $\Sigma(G)$ is G -arc-semiregular.

First observations

Proposition

Suppose G is transitive with $b(G) = 2$ and $\Sigma(G)$ is the Saxl graph of G .

- 1 $\Sigma(G)$ is G -vertex-transitive.
- 2 $\Sigma(G)$ is connected if G is primitive.
- 3 $\Sigma(G)$ is G -arc-semiregular.
- 4 $\Sigma(G)$ is the union of all regular orbital graphs of G .

First observations

Proposition

Suppose G is transitive with $b(G) = 2$ and $\Sigma(G)$ is the Saxl graph of G .

- 1 $\Sigma(G)$ is G -vertex-transitive.
- 2 $\Sigma(G)$ is connected if G is primitive.
- 3 $\Sigma(G)$ is G -arc-semiregular.
- 4 $\Sigma(G)$ is the union of all regular orbital graphs of G .
- 5 $\Sigma(G)$ has valency $r|G_\alpha|$, where r is the number of regular suborbits.

Theorem (Burness & Giudici, 2020)

Let G be a finite transitive base-two permutation group with degree n . Then $\Sigma(G)$ has prime valency p iff one of the following holds:

- $G = C_p \wr C_2$, $n = 2p$, $\Sigma(G) \cong K_{p,p}$.
- $G = S_3$, $n = 3$, $\Sigma(G) \cong K_3$.
- $G = \text{AGL}_1(2^f)$, $n = p + 1 = 2^f$, $\Sigma(G) \cong K_{p+1}$.

Theorem (Burness & Giudici, 2020)

Let G be a finite transitive base-two permutation group with degree n . Then $\Sigma(G)$ has prime valency p iff one of the following holds:

- $G = C_p \wr C_2$, $n = 2p$, $\Sigma(G) \cong K_{p,p}$.
- $G = S_3$, $n = 3$, $\Sigma(G) \cong K_3$.
- $G = \text{AGL}_1(2^f)$, $n = p + 1 = 2^f$, $\Sigma(G) \cong K_{p+1}$.

A group G is called **almost simple** if $\text{soc}(G)$ is non-abelian simple.

Theorem (Chen & H, 2020)

Let G be an almost simple primitive group. Then $\Sigma(G)$ has prime-power valency p^f iff $p = 2$ and one of the following holds:

- $G = \text{PGL}_2(q)$, $q = 9$ or a Fermat prime, $\Sigma(G) \cong J(q + 1, 2)$.
- $G = M_{10}$, $G_\alpha = 8:2$.

Eulerian cycles

By a famous theorem of Euler, a connected graph has an Eulerian cycle iff the degree of every vertex is even.

Theorem (Burness & Giudici, 2020; Chen & H, 2020)

Let G be an almost simple primitive group with stabiliser H . Then one of the following holds:

- $\Sigma(G)$ is Eulerian.
- $G = M_{23}$ and $H = 23:11$.
- $\text{soc}(G) = L_r^\epsilon(q)$, r odd prime, $G \not\leq \text{PGL}_r^\epsilon(q)$ and H is of type $\text{GL}_1^\epsilon(q^r)$.

Eulerian cycles

By a famous theorem of Euler, a connected graph has an Eulerian cycle iff the degree of every vertex is even.

Theorem (Burness & Giudici, 2020; Chen & H, 2020)

Let G be an almost simple primitive group with stabiliser H . Then one of the following holds:

- $\Sigma(G)$ is Eulerian.
- $G = M_{23}$ and $H = 23:11$.
- $\text{soc}(G) = L_r^\epsilon(q)$, r odd prime, $G \not\leq \text{PGL}_r^\epsilon(q)$ and H is of type $\text{GL}_1^\epsilon(q^r)$.

Question. Is $(G, H) = (M_{23}, 23:11)$ the only non-Eulerian example?

Unique regular suborbit

Recall that the valency of $\Sigma(G)$ is $r|G_\alpha|$, where r is the number of regular suborbits.

Unique regular suborbit

Recall that the valency of $\Sigma(G)$ is $r|G_\alpha|$, where r is the number of regular suborbits.

When $r = 1$, $\Sigma(G)$ is an orbital graph and G -arc-transitive.

Unique regular suborbit

Recall that the valency of $\Sigma(G)$ is $r|G_\alpha|$, where r is the number of regular suborbits.

When $r = 1$, $\Sigma(G)$ is an orbital graph and G -arc-transitive.

Question. Can we classify primitive groups with $r = 1$?

Unique regular suborbit

Recall that the valency of $\Sigma(G)$ is $r|G_\alpha|$, where r is the number of regular suborbits.

When $r = 1$, $\Sigma(G)$ is an orbital graph and G -arc-transitive.

Question. Can we classify primitive groups with $r = 1$?

Example

Let $G = \text{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 . Then $\Sigma(G) \cong J(q+1, 2)$ has valency $2(q-1)$.

Indeed, $G_\alpha \cong D_{2(q-1)}$ and so $r = 1$.

Unique regular suborbit

Recall that the valency of $\Sigma(G)$ is $r|G_\alpha|$, where r is the number of regular suborbits.

When $r = 1$, $\Sigma(G)$ is an orbital graph and G -arc-transitive.

Question. Can we classify primitive groups with $r = 1$?

Example

Let $G = \text{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 . Then $\Sigma(G) \cong J(q+1, 2)$ has valency $2(q-1)$.

Indeed, $G_\alpha \cong D_{2(q-1)}$ and so $r = 1$.

- **Burness & Giudici, 2020:** most cases when $\text{soc}(G)$ alternating or sporadic.

Unique regular suborbit

Recall that the valency of $\Sigma(G)$ is $r|G_\alpha|$, where r is the number of regular suborbits.

When $r = 1$, $\Sigma(G)$ is an orbital graph and G -arc-transitive.

Question. Can we classify primitive groups with $r = 1$?

Example

Let $G = \text{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 . Then $\Sigma(G) \cong J(q+1, 2)$ has valency $2(q-1)$.

Indeed, $G_\alpha \cong D_{2(q-1)}$ and so $r = 1$.

- **Burness & Giudici, 2020:** most cases when $\text{soc}(G)$ alternating or sporadic.
- **Burness & H, in progress:** almost simple with soluble stabiliser.

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a base-two transitive permutation group with degree n . Let $\text{val}(G)$ be the valency of $\Sigma(G)$. Set

$$Q(G, 2) := 1 - \mathbb{P}(G, 2) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2} = 1 - \frac{\text{val}(G)}{n}.$$

Probabilistic methods

Let $G \leq \text{Sym}(\Omega)$ be a base-two transitive permutation group with degree n . Let $\text{val}(G)$ be the valency of $\Sigma(G)$. Set

$$Q(G, 2) := 1 - \mathbb{P}(G, 2) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2} = 1 - \frac{\text{val}(G)}{n}.$$

Lemma

If $Q(G, 2) < \frac{1}{t} \leq \frac{1}{2}$, then $\Sigma(G)$ has all of the following properties:

- Any t vertices in $\Sigma(G)$ have a common neighbour;
- $\Sigma(G)$ has diameter at most 2;
- $\Sigma(G)$ has clique number at least $t + 1$;
- $\Sigma(G)$ is Hamiltonian.

Burness-Giudici Conjecture

Recall that

$$Q(G, 2) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2}$$

is the probability that random chosen pairs in Ω do not form a base.

Burness-Giudici Conjecture

Recall that

$$Q(G, 2) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2}$$

is the probability that random chosen pairs in Ω do not form a base.

- $Q(G, 2) < 1 \implies b(G) \leq 2$.
- $Q(G, 2) < \frac{1}{t} \implies \Sigma(G)$ satisfies all statements in the above lemma.

Burness-Giudici Conjecture

Recall that

$$Q(G, 2) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2}$$

is the probability that random chosen pairs in Ω do not form a base.

- $Q(G, 2) < 1 \implies b(G) \leq 2$.
- $Q(G, 2) < \frac{1}{t} \implies \Sigma(G)$ satisfies all statements in the above lemma.

Conjecture (Burness & Giudici, 2020)

Let G be a primitive permutation group with $b(G) = 2$. Then any two vertices in $\Sigma(G)$ have a common neighbour.

Burness-Giudici Conjecture

Recall that

$$Q(G, 2) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_{\alpha\beta} \neq 1\}|}{n^2}$$

is the probability that random chosen pairs in Ω do not form a base.

- $Q(G, 2) < 1 \implies b(G) \leq 2$.
- $Q(G, 2) < \frac{1}{t} \implies \Sigma(G)$ satisfies all statements in the above lemma.

Conjecture (Burness & Giudici, 2020)

Let G be a primitive permutation group with $b(G) = 2$. Then any two vertices in $\Sigma(G)$ have a common neighbour.

Note that if $Q(G, 2) < \frac{1}{2}$, then the conjecture holds.

Some evidence

Example

Let $G = \text{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 . Then $\Sigma(G) \cong J(q+1, 2)$ has valency $2(q-1)$ and thus

$$Q(G, 2) = 1 - \frac{\text{val}(G)}{n} = 1 - \frac{4(q-1)}{q(q+1)} \rightarrow 1 \text{ as } q \rightarrow \infty$$

Some evidence

Example

Let $G = \text{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 . Then $\Sigma(G) \cong J(q+1, 2)$ has valency $2(q-1)$ and thus

$$Q(G, 2) = 1 - \frac{\text{val}(G)}{n} = 1 - \frac{4(q-1)}{q(q+1)} \rightarrow 1 \text{ as } q \rightarrow \infty$$

but $\Sigma(G) \cong J(q+1, 2)$ still satisfies the Burness-Giudici Conjecture.

Some evidence

Example

Let $G = \mathrm{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 . Then $\Sigma(G) \cong J(q+1, 2)$ has valency $2(q-1)$ and thus

$$Q(G, 2) = 1 - \frac{\mathrm{val}(G)}{n} = 1 - \frac{4(q-1)}{q(q+1)} \rightarrow 1 \text{ as } q \rightarrow \infty$$

but $\Sigma(G) \cong J(q+1, 2)$ still satisfies the Burness-Giudici Conjecture.

Example

Suppose $\mathrm{soc}(G) = L_3^\epsilon(q)$ with $q = p \equiv \epsilon \pmod{3}$ and $H = G_\alpha$ is of type $3^{1+2} \cdot \mathrm{Sp}_2(3)$. Then $Q(G, 2) < 8q^{-1} < \frac{1}{2}$ for all $q > 23$.

Some evidence

Example

Let $G = \text{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 . Then $\Sigma(G) \cong J(q+1, 2)$ has valency $2(q-1)$ and thus

$$Q(G, 2) = 1 - \frac{\text{val}(G)}{n} = 1 - \frac{4(q-1)}{q(q+1)} \rightarrow 1 \text{ as } q \rightarrow \infty$$

but $\Sigma(G) \cong J(q+1, 2)$ still satisfies the Burness-Giudici Conjecture.

Example

Suppose $\text{soc}(G) = L_3^\epsilon(q)$ with $q = p \equiv \epsilon \pmod{3}$ and $H = G_\alpha$ is of type $3^{1+2} \cdot \text{Sp}_2(3)$. Then $Q(G, 2) < 8q^{-1} < \frac{1}{2}$ for all $q > 23$.

When $q \leq 23$ we can also check using MAGMA that the conjecture holds.

Further evidence

- All primitive groups with degree $n \leq 4095$

Further evidence

- All primitive groups with degree $n \leq 4095$
- “Most” alternating groups, symmetric groups and sporadic groups

Further evidence

- All primitive groups with degree $n \leq 4095$
- “Most” alternating groups, symmetric groups and sporadic groups
- $\text{soc}(G) = L_2(q)$ (**Chen & Du, 2020; Burness & H, in progress**)

Further evidence

- All primitive groups with degree $n \leq 4095$
- “Most” alternating groups, symmetric groups and sporadic groups
- $\text{soc}(G) = L_2(q)$ (**Chen & Du, 2020; Burness & H, in progress**)
- All almost simple primitive groups with soluble stabilisers (**Burness & H, in progress**)

Further evidence

- All primitive groups with degree $n \leq 4095$
- “Most” alternating groups, symmetric groups and sporadic groups
- $\text{soc}(G) = L_2(q)$ (**Chen & Du, 2020; Burness & H, in progress**)
- All almost simple primitive groups with soluble stabilisers (**Burness & H, in progress**)
- Asymptotic results for many diagonal and twisted wreath type groups (**Fawcett, 2013/21**)

Other invariants

Clique number: maximal size of complete subgraph.

Other invariants

Clique number: maximal size of complete subgraph.

Theorem (Burness & H, in progress)

Let G be an almost simple primitive group with soluble stabiliser H . Suppose $\text{soc}(G) \neq L_2(q)$. Then one of the following holds:

- $\Sigma(G)$ has clique number at least 5.
- $G = A_5$ and $H = S_3$, the clique number of $\Sigma(G)$ is 4.

Other invariants

Clique number: maximal size of complete subgraph.

Theorem (Burness & H, in progress)

Let G be an almost simple primitive group with soluble stabiliser H . Suppose $\text{soc}(G) \neq L_2(q)$. Then one of the following holds:

- $\Sigma(G)$ has clique number at least 5.
- $G = A_5$ and $H = S_3$, the clique number of $\Sigma(G)$ is 4.

Independence number: maximal size of empty subgraph.

Other invariants

Clique number: maximal size of complete subgraph.

Theorem (Burness & H, in progress)

Let G be an almost simple primitive group with soluble stabiliser H . Suppose $\text{soc}(G) \neq L_2(q)$. Then one of the following holds:

- $\Sigma(G)$ has clique number at least 5.
- $G = A_5$ and $H = S_3$, the clique number of $\Sigma(G)$ is 4.

Independence number: maximal size of empty subgraph.

Theorem (Burness & H, in progress)

Let G be an almost simple primitive group with soluble stabiliser H . Then one of the following holds:

- $\Sigma(G)$ has independence number at least 4.
- $G = A_5$ and $H = S_3$, the independence number of $\Sigma(G)$ is 2.

Problems

- **Connectedness.** Characterise transitive groups with connected Saxl graph. If G is quasiprimitive?
- **Automorphisms.**
 - ▶ When do we have $G = \text{Aut}(\Sigma(G))$?
 - ▶ When is $\Sigma(G)$ Cayley?
- **Cycles.** Eulerian cycle? Hamiltonian cycle?
- **Unique regular suborbit.** Can we classify groups with $r = 1$?
- **Other invariants.** Chromatic numbers? Spectrum?

Thanks for your attention!