

Transitive subgroups of primitive groups

Hongyi Huang

Finite Groups, Fusion Systems and Applications

Mathematisches Forschungsinstitut Oberwolfach

20 March 2025



Group factorisations

Let G be a finite group and $H, K \leq G$.

Facts.

- $G = HK \iff K$ is **transitive** on G/H .

Group factorisations

Let G be a finite group and $H, K \leq G$.

Facts.

- $G = HK \iff K$ is **transitive** on G/H .
- $G = HK$ and $H \cap K = 1 \iff K$ is **regular** on G/H .

Group factorisations

Let G be a finite group and $H, K \leq G$.

Facts.

- $G = HK \iff K$ is **transitive** on G/H .
- $G = HK$ and $H \cap K = 1 \iff K$ is **regular** on G/H .

The expression $G = HK$ is called a **factorisation** of G .

Factorisations of almost simple groups (thanks Cheryl!)

Theorem. The factorisations of almost simple groups are classified.

Contributed earlier by Giudici, Liebeck, Praeger, Saxl...

Recent advances:

- H or K has at least two insoluble composition factors:

Li & Xia, 2019, JAlg

- Both H and K are insoluble (for classical groups):

Li, Wang & Xia, 2024, arXiv (102 pages)

- At least one of H and K is soluble:

Li & Xia, 2022, AMS Memoirs

Burness & Li, 2021, Adv

Feng, Li, Li, Wang, Xia & Zou, 2024, arXiv (final step)

Primitive groups

A permutation group G is called **primitive** if G_α is a maximal subgroup.

Main theme. Classify the transitive subgroups of primitive groups.

Primitive groups

A permutation group G is called **primitive** if G_α is a maximal subgroup.

Main theme. Classify the transitive subgroups of primitive groups.

The **O'Nan-Scott theorem** divides finite primitive groups into 5 types.

- Affine (HA)
- Almost simple (AS) ✓
- Diagonal type (HS & SD)
- Product type (HC, CD & PA)
- Twisted wreath products (TW)

Primitive groups

A permutation group G is called **primitive** if G_α is a maximal subgroup.

Main theme. Classify the transitive subgroups of primitive groups.

The **O'Nan-Scott theorem** divides finite primitive groups into 5 types.

- Affine (HA)
- Almost simple (AS) ✓
- Diagonal type (HS & SD)
- Product type (HC, CD & PA) (ask Barbara, Cheryl and Michael)
- Twisted wreath products (TW)

Primitive groups

A permutation group G is called **primitive** if G_α is a maximal subgroup.

Main theme. Classify the transitive subgroups of primitive groups.

The **O'Nan-Scott theorem** divides finite primitive groups into 5 types.

- Affine (HA)
- Almost simple (AS) ✓
- Diagonal type (HS & SD)
- Product type (HC, CD & PA) (ask Barbara, Cheryl and Michael)
- Twisted wreath products (TW)

Problem. Classify the **regular** subgroups and the **soluble** transitive subgroups of primitive groups of **diagonal type**, up to conjugacy.

Holomorph simple groups (HS)

Let T be a non-abelian finite simple group and let

$$G = \text{Hol}(T) = T : \text{Aut}(T) = T^2 : \text{Out}(T)$$

be the **holomorph** of T . So $G \leq \text{Sym}(T)$ is primitive.

Holomorph simple groups (HS)

Let T be a non-abelian finite simple group and let

$$G = \text{Hol}(T) = T : \text{Aut}(T) = T^2 : \text{Out}(T)$$

be the **holomorph** of T . So $G \leq \text{Sym}(T)$ is primitive.

Key observation (Liebeck, Praeger & Saxl, 2000).

If B is a transitive subgroup of G , then there exist $H, K \leq \text{Aut}(T)$ isomorphic to some quotient groups of B such that

$$T \trianglelefteq HK = HT = KT \leq \text{Aut}(T).$$

Holomorph simple groups (HS)

Let T be a non-abelian finite simple group and let

$$G = \text{Hol}(T) = T : \text{Aut}(T) = T^2 : \text{Out}(T)$$

be the **holomorph** of T . So $G \leq \text{Sym}(T)$ is primitive.

Key observation (Liebeck, Praeger & Saxl, 2000).

If B is a transitive subgroup of G , then there exist $H, K \leq \text{Aut}(T)$ isomorphic to some quotient groups of B such that

$$T \trianglelefteq HK = HT = KT \leq \text{Aut}(T).$$

Example

$T = A_{q+1}$, $B \cong (A_{q-2} \times \text{PSL}_2(q)).2$ associated to $S_{q+1} = S_{q-2} \text{PGL}_2(q)$.

Holomorph simple groups (HS)

Let T be a non-abelian finite simple group and let

$$G = \text{Hol}(T) = T : \text{Aut}(T) = T^2 : \text{Out}(T)$$

be the **holomorph** of T . So $G \leq \text{Sym}(T)$ is primitive.

Key observation (Liebeck, Praeger & Saxl, 2000).

If B is a transitive subgroup of G , then there exist $H, K \leq \text{Aut}(T)$ isomorphic to some quotient groups of B such that

$$T \trianglelefteq HK = HT = KT \leq \text{Aut}(T).$$

Example

$$T = A_{q+1}, B \cong (A_{q-2} \times \text{PSL}_2(q)).2 \text{ associated to } S_{q+1} = S_{q-2} \text{PGL}_2(q).$$

Conversely, if $T = HK$, then there exists a transitive subgroup of G isomorphic to $H \times K$.

The key step

Key observation (Liebeck, Praeger & Saxl, 2000).

If B is a transitive subgroup of G , then there exist $H, K \leq \text{Aut}(T)$ isomorphic to some quotient groups of B such that

$$T \trianglelefteq HK = HT = KT \leq \text{Aut}(T). \quad (\star)$$

The key step

Key observation (Liebeck, Praeger & Saxl, 2000).

If B is a transitive subgroup of G , then there exist $H, K \leq \text{Aut}(T)$ isomorphic to some quotient groups of B such that

$$T \trianglelefteq HK = HT = KT \leq \text{Aut}(T). \quad (\star)$$

If B is soluble, then both H and K are soluble, so (basically) $T = \text{PSL}_2(q)$.

The key step

Key observation (Liebeck, Praeger & Saxl, 2000).

If B is a transitive subgroup of G , then there exist $H, K \leq \text{Aut}(T)$ isomorphic to some quotient groups of B such that

$$T \trianglelefteq HK = HT = KT \leq \text{Aut}(T). \quad (\star)$$

If B is soluble, then both H and K are soluble, so (basically) $T = \text{PSL}_2(q)$.

If B is regular, then there exists $N \trianglelefteq H$ and $M \trianglelefteq K$ such that

$$H/N \cong K/M \text{ and } |H : N| = |HK : T||H \cap K|. \quad (\star\star)$$

The key step

Key observation (Liebeck, Praeger & Saxl, 2000).

If B is a transitive subgroup of G , then there exist $H, K \leq \text{Aut}(T)$ isomorphic to some quotient groups of B such that

$$T \trianglelefteq HK = HT = KT \leq \text{Aut}(T). \quad (\star)$$

If B is soluble, then both H and K are soluble, so (basically) $T = \text{PSL}_2(q)$.

If B is regular, then there exists $N \trianglelefteq H$ and $M \trianglelefteq K$ such that

$$H/N \cong K/M \text{ and } |H : N| = |HK : T||H \cap K|. \quad (\star\star)$$

It is NOT easy to determine the factorisations satisfying (\star) and $(\star\star)$.

Suppose $T = A_n$ and $L \in \{A_n, S_n\}$. If $L = HK$ then H is transitive on m -subsets and $A_{n-m} \trianglelefteq K \leq S_{n-m} \times S_m$, with $1 \leq m \leq 5$.

Suppose $T = A_n$ and $L \in \{A_n, S_n\}$. If $L = HK$ then H is transitive on m -subsets and $A_{n-m} \trianglelefteq K \leq S_{n-m} \times S_m$, with $1 \leq m \leq 5$.

Example

Assume (\star) and $(\star\star)$, with $m = 1$. Then H is regular on $[n]$, and

- the Sylow 2-subgroups of H are not cyclic $\implies (L, K) = (A_n, A_{n-1})$.
- otherwise $(L, K) = (S_n, S_{n-1})$.

Suppose $T = A_n$ and $L \in \{A_n, S_n\}$. If $L = HK$ then H is transitive on m -subsets and $A_{n-m} \trianglelefteq K \leq S_{n-m} \times S_m$, with $1 \leq m \leq 5$.

Example

Assume (\star) and $(\star\star)$, with $m = 1$. Then H is regular on $[n]$, and

- the Sylow 2-subgroups of H are not cyclic $\implies (L, K) = (A_n, A_{n-1})$.
- otherwise $(L, K) = (S_n, S_{n-1})$.

Example

Assume (\star) and $(\star\star)$, with $m = 3$. Then (basically) $n = q + 1$ and

- q even, $L = A_n$, $H = \text{PSL}_2(q)$, $K = A_{n-3}$.
- $q \equiv 3 \pmod{4}$, $L = A_n$, $H = \text{PSL}_2(q)$, $K = S_{n-3}$.
- $\sqrt{q} \equiv 3 \pmod{4}$, $L = A_n$, $H = \text{PSL}_2(q).2_3$, $K \in \{A_{n-3}, S_{n-3}\}$.
- $L = S_n$, $H = \text{PGL}_2(q)$, $K = S_{n-3}$.
- $q \equiv 1 \pmod{4}$, $L = S_n$, $H = \text{PGL}_2(q)$, $K = A_{n-3} \times S_2$.
- $\sqrt{q} \equiv 1 \pmod{4}$, $L = S_n$, $H = \text{PSL}_2(q).2_3$, $K \in \{S_{n-3}, A_{n-3} \times S_2\}$.

Main results on holomorph simple groups

Theorem (H & Wang, 2025+)

For every finite simple group T , the **soluble transitive** subgroups of $\text{Hol}(T)$ are determined, up to conjugacy.

Theorem (H & Wang, 2025+)

For every finite simple group T , the **regular** subgroups of $\text{Hol}(T)$ are determined, up to conjugacy.

Application: Hopf-Galois structure and skew braces

For a finite group G , TFAE:

- B is isomorphic to a regular subgroup of $\text{Hol}(G)$;
- There exists a **Hopf-Galois structure** of type B on any Galois G -extension.
- $G \cong (X, +)$ and $B \cong (X, \circ)$ for some **skew brace** $(X, +, \circ)$.

Application: Hopf-Galois structure and skew braces

For a finite group G , TFAE:

- B is isomorphic to a regular subgroup of $\text{Hol}(G)$;
- There exists a **Hopf-Galois structure** of type B on any Galois G -extension.
- $G \cong (X, +)$ and $B \cong (X, \circ)$ for some **skew brace** $(X, +, \circ)$.

Theorem (H & Wang, 2025+)

The types of Hopf-Galois structures are determined on any Galois extension whose Galois group is finite simple.

Theorem (H & Wang, 2025+)

The skew braces with finite simple additive groups are classified.

General diagonal type groups

Let $k \geq 2$, T be a non-abelian simple group, and let

$$X := \{(t, \dots, t) : t \in T\} \leq T^k.$$

Then $T^k \leq \text{Sym}(\Omega)$ with $\Omega = T^k/X$ of size $|T|^{k-1}$.

General diagonal type groups

Let $k \geq 2$, T be a non-abelian simple group, and let

$$X := \{(t, \dots, t) : t \in T\} \leq T^k.$$

Then $T^k \leq \text{Sym}(\Omega)$ with $\Omega = T^k/X$ of size $|T|^{k-1}$.

A group $G \leq \text{Sym}(\Omega)$ is of **diagonal type** if

$$T^k \trianglelefteq G \leq N_{\text{Sym}(\Omega)}(T^k) \cong T^k.(\text{Out}(T) \times S_k).$$

General diagonal type groups

Let $k \geq 2$, T be a non-abelian simple group, and let

$$X := \{(t, \dots, t) : t \in T\} \leq T^k.$$

Then $T^k \leq \text{Sym}(\Omega)$ with $\Omega = T^k/X$ of size $|T|^{k-1}$.

A group $G \leq \text{Sym}(\Omega)$ is of **diagonal type** if

$$T^k \trianglelefteq G \leq N_{\text{Sym}(\Omega)}(T^k) \cong T^k.(\text{Out}(T) \times S_k).$$

Note. G induces $P_G \leq S_k$. And G is **primitive** if and only if

P_G is primitive (SD), or $k = 2$ and $P_G = A_2 = 1$ (HS).

General diagonal type groups

Let $k \geq 2$, T be a non-abelian simple group, and let

$$X := \{(t, \dots, t) : t \in T\} \leq T^k.$$

Then $T^k \leq \text{Sym}(\Omega)$ with $\Omega = T^k/X$ of size $|T|^{k-1}$.

A group $G \leq \text{Sym}(\Omega)$ is of **diagonal type** if

$$T^k \trianglelefteq G \leq N_{\text{Sym}(\Omega)}(T^k) \cong T^k.(\text{Out}(T) \times S_k).$$

Note. G induces $P_G \leq S_k$. And G is **primitive** if and only if

P_G is primitive (SD), or $k = 2$ and $P_G = A_2 = 1$ (HS).

Example. $\text{Hol}(T) = T^2. \text{Out}(T)$.

General diagonal type groups

Suppose $B \leq G \leq T^k$. $(\text{Out}(T) \times S_k)$ is transitive.

- ① Assume B **does not contain** any simple direct factor of T^k .

General diagonal type groups

Suppose $B \leq G \leq T^k$. $(\text{Out}(T) \times S_k)$ is transitive.

① Assume B **does not contain** any simple direct factor of T^k .

Liebeck, Praeger & Saxl, 2000: $k \leq 3$ and $B \leq T^k \cdot \text{Out}(T)$.

General diagonal type groups

Suppose $B \leq G \leq T^k$. $(\text{Out}(T) \times S_k)$ is transitive.

① Assume B **does not contain** any simple direct factor of T^k .

Liebeck, Praeger & Saxl, 2000: $k \leq 3$ and $B \leq T^k \cdot \text{Out}(T)$.

Theorem (H & Wang, 2025+)

If B is regular or soluble, then $k = 2$ and $B \leq \text{Hol}(T)$ is determined.

General diagonal type groups

Suppose $B \leq G \leq T^k$. $(\text{Out}(T) \times S_k)$ is transitive.

① Assume B **does not contain** any simple direct factor of T^k .

Liebeck, Praeger & Saxl, 2000: $k \leq 3$ and $B \leq T^k \cdot \text{Out}(T)$.

Theorem (H & Wang, 2025+)

If B is regular or soluble, then $k = 2$ and $B \leq \text{Hol}(T)$ is determined.

② Assume $B \leq G$ is **regular** and assume B **contains** exactly $\ell \geq 1$ simple direct factors of T^k .

General diagonal type groups

Suppose $B \leq G \leq T^k$. ($\text{Out}(T) \times S_k$) is transitive.

① Assume B **does not contain** any simple direct factor of T^k .

Liebeck, Praeger & Saxl, 2000: $k \leq 3$ and $B \leq T^k \cdot \text{Out}(T)$.

Theorem (H & Wang, 2025+)

If B is regular or soluble, then $k = 2$ and $B \leq \text{Hol}(T)$ is determined.

② Assume $B \leq G$ is **regular** and assume B **contains** exactly $\ell \geq 1$ simple direct factors of T^k .

Note. If $\ell = k - 1$ then $B \cong T^{k-1}$. So we assume $\ell \leq k - 2$.

General diagonal type groups

Suppose $B \leq G \leq T^k$. ($\text{Out}(T) \times S_k$) is transitive.

① Assume B **does not contain** any simple direct factor of T^k .

Liebeck, Praeger & Saxl, 2000: $k \leq 3$ and $B \leq T^k \cdot \text{Out}(T)$.

Theorem (H & Wang, 2025+)

If B is regular or soluble, then $k = 2$ and $B \leq \text{Hol}(T)$ is determined.

② Assume $B \leq G$ is **regular** and assume B **contains** exactly $\ell \geq 1$ simple direct factors of T^k .

Note. If $\ell = k - 1$ then $B \cong T^{k-1}$. So we assume $\ell \leq k - 2$.

Morris & Spiga, 2021: $\ell = k - 3$ or $k - 2$, and $B \leq T^k \cdot \text{Out}(T)$.

General diagonal type groups

Suppose $B \leq G \leq T^k$. ($\text{Out}(T) \times S_k$) is transitive.

① Assume B **does not contain** any simple direct factor of T^k .

Liebeck, Praeger & Saxl, 2000: $k \leq 3$ and $B \leq T^k$. $\text{Out}(T)$.

Theorem (H & Wang, 2025+)

If B is regular or soluble, then $k = 2$ and $B \leq \text{Hol}(T)$ is determined.

② Assume $B \leq G$ is **regular** and assume B **contains** exactly $\ell \geq 1$ simple direct factors of T^k .

Note. If $\ell = k - 1$ then $B \cong T^{k-1}$. So we assume $\ell \leq k - 2$.

Morris & Spiga, 2021: $\ell = k - 3$ or $k - 2$, and $B \leq T^k$. $\text{Out}(T)$.

Theorem (H & Wang, 2025+)

$\ell = k - 2$, and B is determined.

General diagonal type groups

Suppose $B \leq G \leq T^k \cdot (\text{Out}(T) \times S_k)$ is transitive.

① Assume B **does not contain** any simple direct factor of T^k .

Liebeck, Praeger & Saxl, 2000: $k \leq 3$ and $B \leq T^k \cdot \text{Out}(T)$.

Theorem (H & Wang, 2025+)

If B is regular or soluble, then $k = 2$ and $B \leq \text{Hol}(T)$ is determined.

② Assume $B \leq G$ is **regular** and assume B **contains** exactly $\ell \geq 1$ simple direct factors of T^k .

Note. If $\ell = k - 1$ then $B \cong T^{k-1}$. So we assume $\ell \leq k - 2$.

Morris & Spiga, 2021: $\ell = k - 3$ or $k - 2$, and $B \leq T^k \cdot \text{Out}(T)$.

Theorem (H & Wang, 2025+)

$\ell = k - 2$, and B is determined.

Example. $B = T^{k-2} \times H \times K$, with $H \times K$ a regular subgroup of $\text{Hol}(T)$.

The main result

Theorem (H & Wang, 2025+). The regular and the soluble transitive subgroups of diagonal type groups are determined, up to conjugacy.

Danke schön!