

Permutation groups of rank three

Hongyi Huang

Group Theory in Florence IV

2 July 2024

joint with C.H. Li and Y.Z. Zhu



Ranks

Let $G \leq \text{Sym}(\Omega)$ be a finite transitive permutation group.

rank(G): The number of G_α -orbits on Ω (G -orbits on Ω^2).

Ranks

Let $G \leq \text{Sym}(\Omega)$ be a finite transitive permutation group.

rank(G): The number of G_α -orbits on Ω (G -orbits on Ω^2).

Low rank groups:

$$\text{rank}(G) = 1 \iff |\Omega| = 1.$$

$$\text{rank}(G) = 2 \iff G \text{ is 2-transitive.}$$

- G is determined via CFSG;

Ranks

Let $G \leq \text{Sym}(\Omega)$ be a finite transitive permutation group.

rank(G): The number of G_α -orbits on Ω (G -orbits on Ω^2).

Low rank groups:

$$\text{rank}(G) = 1 \iff |\Omega| = 1.$$

$$\text{rank}(G) = 2 \iff G \text{ is 2-transitive.}$$

- G is determined via CFSG;
- G is affine or almost simple (Burnside 1900).

Ranks

Let $G \leq \text{Sym}(\Omega)$ be a finite transitive permutation group.

rank(G): The number of G_α -orbits on Ω (G -orbits on Ω^2).

Low rank groups:

$\text{rank}(G) = 1 \iff |\Omega| = 1.$

$\text{rank}(G) = 2 \iff G$ is 2-transitive.

- G is determined via CFSG;
- G is affine or almost simple (Burnside 1900).

Problem. *Classify the transitive groups G with $\text{rank}(G) = 3$.*

Ranks

Let $G \leq \text{Sym}(\Omega)$ be a finite transitive permutation group.

rank(G): The number of G_α -orbits on Ω (G -orbits on Ω^2).

Low rank groups:

$$\text{rank}(G) = 1 \iff |\Omega| = 1.$$

$$\text{rank}(G) = 2 \iff G \text{ is 2-transitive.}$$

- G is determined via CFSG;
- G is affine or almost simple (Burnside 1900).

Problem. *Classify the transitive groups G with $\text{rank}(G) = 3$.*

Example. $G = \text{GL}_n(3)$ and $\Omega = \mathbb{F}_3^n \setminus \{0\}$.

Primitivity

Primitive: G_α is a maximal subgroup of G .

- Bannai, Foulser, Kantor, Liebeck, Liebler and Saxl (1969–1986)

Primitivity

Primitive: G_α is a maximal subgroup of G .

- Bannai, Foulser, Kantor, Liebeck, Liebler and Saxl (1969–1986)

Quasiprimitive: Every non-trivial normal subgroup of G is transitive.

- Devillers, Giudici, Li, Pearce & Praeger (2011)

primitive \implies quasiprimitive

Primitivity

Primitive: G_α is a maximal subgroup of G .

- Bannai, Foulser, Kantor, Liebeck, Liebler and Saxl (1969–1986)

Quasiprimitive: Every non-trivial normal subgroup of G is transitive.

- Devillers, Giudici, Li, Pearce & Praeger (2011)

Innately transitive: G has a transitive minimal normal subgroup.

- Baykalov, Devillers & Praeger (2023)

primitive \implies quasiprimitive \implies innately transitive

Primitivity

Primitive: G_α is a maximal subgroup of G .

- Bannai, Foulser, Kantor, Liebeck, Liebler and Saxl (1969–1986)

Quasiprimitive: Every non-trivial normal subgroup of G is transitive.

- Devillers, Giudici, Li, Pearce & Praeger (2011)

Innately transitive: G has a transitive minimal normal subgroup.

- Baykalov, Devillers & Praeger (2023)

e.g. $GL_n(3) \cong C_2 \times PSL_n(3)$ when n is odd.

primitive \implies quasiprimitive \implies innately transitive

Primitivity

Primitive: G_α is a maximal subgroup of G .

- Bannai, Foulser, Kantor, Liebeck, Liebler and Saxl (1969–1986)

Quasiprimitive: Every non-trivial normal subgroup of G is transitive.

- Devillers, Giudici, Li, Pearce & Praeger (2011)

Innately transitive: G has a transitive minimal normal subgroup.

- Baykalov, Devillers & Praeger (2023)

e.g. $\mathrm{GL}_n(3) \cong C_2 \times \mathrm{PSL}_n(3)$ when n is odd.

Remark. When n is even, $\mathrm{GL}_n(3)$ is not innately transitive.

primitive \implies quasiprimitive \implies innately transitive

Primitivity

Primitive: G_α is a maximal subgroup of G .

- Bannai, Foulser, Kantor, Liebeck, Liebler and Saxl (1969–1986)

Quasiprimitive: Every non-trivial normal subgroup of G is transitive.

- Devillers, Giudici, Li, Pearce & Praeger (2011)

Innately transitive: G has a transitive minimal normal subgroup.

- Baykalov, Devillers & Praeger (2023)

e.g. $GL_n(3) \cong C_2 \times PSL_n(3)$ when n is odd.

Remark. When n is even, $GL_n(3)$ is not innately transitive.

Note. Every normal subgroup of $GL_n(3)$ is transitive or semiregular (such a group is said to be **semiprimitive**).

primitive \implies quasiprimitive \implies innately transitive \implies semiprimitive

Imprimitive groups

Imprimitive: G_α is not a maximal subgroup of G .

Imprimitive groups

Imprimitive: G_α is not a maximal subgroup of G .

G is imprimitive \iff there exists $B \subseteq \Omega$ with $|B| \notin \{1, |\Omega|\}$ s.t.

$$B^g \cap B = B \text{ or } \emptyset \text{ for all } g \in G.$$

Imprimitive groups

Imprimitive: G_α is not a maximal subgroup of G .

G is imprimitive \iff there exists $B \subseteq \Omega$ with $|B| \notin \{1, |\Omega|\}$ s.t.

$$B^g \cap B = B \text{ or } \emptyset \text{ for all } g \in G.$$

The set B is called a (non-trivial) **block**, and B^G is called a **block system**.

Imprimitive groups

Imprimitive: G_α is not a maximal subgroup of G .

G is imprimitive \iff there exists $B \subseteq \Omega$ with $|B| \notin \{1, |\Omega|\}$ s.t.

$$B^g \cap B = B \text{ or } \emptyset \text{ for all } g \in G.$$

The set B is called a (non-trivial) **block**, and B^G is called a **block system**.

Imprimitive rank 3 groups:

- G has a unique non-trivial block system $\mathcal{B} = B^G$.

Imprimitive groups

Imprimitive: G_α is not a maximal subgroup of G .


G is imprimitive \iff there exists $B \subseteq \Omega$ with $|B| \notin \{1, |\Omega|\}$ s.t.

$$B^g \cap B = B \text{ or } \emptyset \text{ for all } g \in G.$$

The set B is called a (non-trivial) **block**, and B^G is called a **block system**.

Imprimitive rank 3 groups:

- G has a unique non-trivial block system $\mathcal{B} = B^G$.



Imprimitive groups

Imprimitive: G_α is not a maximal subgroup of G .


G is imprimitive \iff there exists $B \subseteq \Omega$ with $|B| \notin \{1, |\Omega|\}$ s.t.

$$B^g \cap B = B \text{ or } \emptyset \text{ for all } g \in G.$$

The set B is called a (non-trivial) **block**, and B^G is called a **block system**.


Imprimitive rank 3 groups:

- G has a unique non-trivial block system $\mathcal{B} = B^G$.




- The induced groups $G^{\mathcal{B}}$ (on \mathcal{B}) and G_B^B (on B) are 2-transitive.

Imprimitive rank three groups



Imprimitive rank three groups



G_B^B affine

Theorem (H, Li & Zhu, 2024+)

Suppose G_B^B is affine and $\text{rank}(G) = 3$.

G_B^B affine

Theorem (H, Li & Zhu, 2024+)

Suppose G_B^B is affine and $\text{rank}(G) = 3$. Then one of the following holds.

- G^B is almost simple and G is **semiprimitive**;
- G^B is affine and $N \trianglelefteq G \leq N: \text{Aut}(N)$;

G_B^B affine

Theorem (H, Li & Zhu, 2024+)

Suppose G_B^B is affine and $\text{rank}(G) = 3$. Then one of the following holds.

- G^B is almost simple and G is **semiprimitive**;
- G^B is affine and $N \trianglelefteq G \leq N: \text{Aut}(N)$;
 $(N, \text{Aut}(N))$ is determined (Li & Zhu, 2024+)

Theorem (H, Li & Zhu, 2024+)

Suppose G_B^B is affine and $\text{rank}(G) = 3$. Then one of the following holds.

- G^B is almost simple and G is **semiprimitive**;
- G^B is affine and $N \trianglelefteq G \leq N: \text{Aut}(N)$;
 $(N, \text{Aut}(N))$ is determined (Li & Zhu, 2024+)
- $K_{(B)}$ is transitive on B' ;
- $K_{(B)} \neq 1$ is intransitive on B' , and G has an elementary abelian self-centralising normal subgroup,

where $K = G_{(B)}$ and $B, B' \in \mathcal{B}$.

Theorem (H, Li & Zhu, 2024+)

Suppose G_B^B is affine and $\text{rank}(G) = 3$. Then one of the following holds.

- G^B is almost simple and G is **semiprimitive**;
- G^B is affine and $N \trianglelefteq G \leq N: \text{Aut}(N)$;
 $(N, \text{Aut}(N))$ is determined (Li & Zhu, 2024+)
- $K_{(B)}$ is transitive on B' ;
always rank 3
- $K_{(B)} \neq 1$ is intransitive on B' , and G has an elementary abelian self-centralising normal subgroup,

where $K = G_{(B)}$ and $B, B' \in \mathcal{B}$.

G_B^B affine

Theorem (H, Li & Zhu, 2024+)

Suppose G_B^B is affine and $\text{rank}(G) = 3$. Then one of the following holds.

- G^B is almost simple and G is **semiprimitive**;
- G^B is affine and $N \trianglelefteq G \leq N: \text{Aut}(N)$;
 $(N, \text{Aut}(N))$ is determined (Li & Zhu, 2024+)
- $K_{(B)}$ is transitive on B' ;
always rank 3
- $K_{(B)} \neq 1$ is intransitive on B' , and G has an elementary abelian self-centralising normal subgroup,
no known examples

where $K = G_{(B)}$ and $B, B' \in \mathcal{B}$.

Imprimitive rank three groups (G_B^B affine)

G^B affine	$N \trianglelefteq G$ regular N is known	$K_{(B)}$ trans on B'	$K_{(B)} \neq 1$ intrans on B'
G^B almost simple	semiprimitive		

Semiprimitive linear groups

Let $X = \mathrm{GL}_n(q) \leq \mathrm{Sym}(\Gamma)$ with $\Gamma = \mathbb{F}_q^n \setminus \{0\}$.

Let $C \leq Z(X)$ and $\Omega = \{C\text{-orbits on } \Gamma\}$. So $G := X/C \leq \mathrm{Sym}(\Omega)$.

Semiprimitive linear groups

Let $X = \mathrm{GL}_n(q) \leq \mathrm{Sym}(\Gamma)$ with $\Gamma = \mathbb{F}_q^n \setminus \{0\}$.

Let $C \leq Z(X)$ and $\Omega = \{C\text{-orbits on } \Gamma\}$. So $G := X/C \leq \mathrm{Sym}(\Omega)$.

- G is semiprimitive.
- $\mathrm{rank}(G) = |Z(X) : C| + 1$.

Semiprimitive linear groups

Let $X = \mathrm{GL}_n(q) \leq \mathrm{Sym}(\Gamma)$ with $\Gamma = \mathbb{F}_q^n \setminus \{0\}$.

Let $C \leq Z(X)$ and $\Omega = \{C\text{-orbits on } \Gamma\}$. So $G := X/C \leq \mathrm{Sym}(\Omega)$.

- G is semiprimitive.
- $\mathrm{rank}(G) = |Z(X) : C| + 1$.

Examples

If $C = Z(X)$, then $G = \mathrm{PGL}_n(q)$ is 2-transitive.

Semiprimitive linear groups

Let $X = \mathrm{GL}_n(q) \leq \mathrm{Sym}(\Gamma)$ with $\Gamma = \mathbb{F}_q^n \setminus \{0\}$.

Let $C \leq Z(X)$ and $\Omega = \{C\text{-orbits on } \Gamma\}$. So $G := X/C \leq \mathrm{Sym}(\Omega)$.

- G is semiprimitive.
- $\mathrm{rank}(G) = |Z(X) : C| + 1$.

Examples

If $C = Z(X)$, then $G = \mathrm{PGL}_n(q)$ is 2-transitive.

If $C = 1$, then $G = \mathrm{GL}_n(q)$ and $\mathrm{rank}(G) = q$.

Semiprimitive linear groups

Let $X = \mathrm{GL}_n(q) \leq \mathrm{Sym}(\Gamma)$ with $\Gamma = \mathbb{F}_q^n \setminus \{0\}$.

Let $C \leq Z(X)$ and $\Omega = \{C\text{-orbits on } \Gamma\}$. So $G := X/C \leq \mathrm{Sym}(\Omega)$.

- G is semiprimitive.
- $\mathrm{rank}(G) = |Z(X) : C| + 1$.

Examples

If $C = Z(X)$, then $G = \mathrm{PGL}_n(q)$ is 2-transitive.

If $C = 1$, then $G = \mathrm{GL}_n(q)$ and $\mathrm{rank}(G) = q$.

If $|Z(X) : C| = 2$ (so $G = 2.\mathrm{PGL}_n(q)$), then $\mathrm{rank}(G) = 3$ with

- $|B| = 2$ and $G_B^B = S_2$;
- $\mathcal{B} = \{1\text{-spaces}\}$ and $G^{\mathcal{B}} = \mathrm{PGL}_n(q)$.

Semiprimitive rank three groups

Theorem (H, Li & Zhu, 2024+)

Suppose G^B is almost simple, G_B^B is affine and G is semiprimitive but not innately transitive.

Semiprimitive rank three groups

Theorem (H, Li & Zhu, 2024+)

Suppose $G^{\mathcal{B}}$ is almost simple, $G_B^{\mathcal{B}}$ is affine and G is semiprimitive but not innately transitive. Then $\text{rank}(G) = 3$ if one of the following holds.

- $(G, G_{\alpha}) = (3.S_6, S_5)$ or $(2.M_{12}, M_{11})$.
- Ω is the set of C -orbits on $\mathbb{F}_q^n \setminus \{0\}$ for $C \leq Z(\text{GL}_d(q))$, and

$$r. \text{PSL}_d(q) \cong \text{SL}_d(q)C/C \leq G \leq \Gamma\text{L}_d(q)/C \cong r. \text{P}\Gamma\text{L}_d(q).$$

Semiprimitive rank three groups

Theorem (H, Li & Zhu, 2024+)

Suppose $G^{\mathcal{B}}$ is almost simple, $G_B^{\mathcal{B}}$ is affine and G is semiprimitive but not innately transitive. Then $\text{rank}(G) = 3$ if one of the following holds.

- $(G, G_{\alpha}) = (3.S_6, S_5)$ or $(2.M_{12}, M_{11})$.
- Ω is the set of C -orbits on $\mathbb{F}_q^n \setminus \{0\}$ for $C \leq Z(\text{GL}_d(q))$, and

$$r. \text{PSL}_d(q) \cong \text{SL}_d(q)C/C \leq G \leq \Gamma\text{L}_d(q)/C \cong r. \text{P}\Gamma\text{L}_d(q).$$

e.g. $2. \text{PGL}_n(q)$, necessary and sufficient conditions are given

Semiprimitive rank three groups

Theorem (H, Li & Zhu, 2024+)

Suppose G^B is almost simple, G_B^B is affine and G is semiprimitive but not innately transitive. Then $\text{rank}(G) = 3$ if one of the following holds.

- $(G, G_\alpha) = (3.S_6, S_5)$ or $(2.M_{12}, M_{11})$.
- Ω is the set of C -orbits on $\mathbb{F}_q^n \setminus \{0\}$ for $C \leq Z(\text{GL}_d(q))$, and

$$r. \text{PSL}_d(q) \cong \text{SL}_d(q)C/C \leq G \leq \Gamma\text{L}_d(q)/C \cong r. \text{P}\Gamma\text{L}_d(q).$$

e.g. $2. \text{PGL}_n(q)$, necessary and sufficient conditions are given

This classifies the semiprimitive rank 3 groups G with G^B almost simple.

Semiprimitive rank three groups

Theorem (H, Li & Zhu, 2024+)

Suppose $G^{\mathcal{B}}$ is almost simple, $G_B^{\mathcal{B}}$ is affine and G is semiprimitive but not innately transitive. Then $\text{rank}(G) = 3$ if one of the following holds.

- $(G, G_{\alpha}) = (3.S_6, S_5)$ or $(2.M_{12}, M_{11})$.
- Ω is the set of C -orbits on $\mathbb{F}_q^n \setminus \{0\}$ for $C \leq Z(\text{GL}_d(q))$, and

$$r. \text{PSL}_d(q) \cong \text{SL}_d(q)C/C \leq G \leq \Gamma\text{L}_d(q)/C \cong r. \text{P}\Gamma\text{L}_d(q).$$

e.g. $2. \text{PGL}_n(q)$, necessary and sufficient conditions are given

This classifies the semiprimitive rank 3 groups G with $G^{\mathcal{B}}$ almost simple.

H, Li & Zhu, 2024+: If $G^{\mathcal{B}}$ is affine, G is semiprimitive and $\text{rank}(G) = 3$, then $G_B^{\mathcal{B}}$ is affine and $N \trianglelefteq G \leq N: \text{Aut}(N)$ with N a special group.

Semiprimitive rank three groups

Theorem (H, Li & Zhu, 2024+)

Suppose G^B is almost simple, G_B^B is affine and G is semiprimitive but not innately transitive. Then $\text{rank}(G) = 3$ if one of the following holds.

- $(G, G_\alpha) = (3.S_6, S_5)$ or $(2.M_{12}, M_{11})$.
- Ω is the set of C -orbits on $\mathbb{F}_q^n \setminus \{0\}$ for $C \leq Z(\text{GL}_d(q))$, and

$$r. \text{PSL}_d(q) \cong \text{SL}_d(q)C/C \leq G \leq \Gamma\text{L}_d(q)/C \cong r. \text{P}\Gamma\text{L}_d(q).$$

e.g. $2. \text{PGL}_n(q)$, necessary and sufficient conditions are given

This classifies the semiprimitive rank 3 groups G with G^B almost simple.

H, Li & Zhu, 2024+: If G^B is affine, G is semiprimitive and $\text{rank}(G) = 3$, then G_B^B is affine and $N \trianglelefteq G \leq N: \text{Aut}(N)$ with N a special group.

Li, Yi & Zhu (last Sunday): such G is determined.

Summary

