

# Base-two primitive permutation groups

Hong Yi Huang

Young Group Theorist Workshop, SwissMAP, Les Diablerets

5 September 2022



# Bases

Let  $G \leq \text{Sym}(\Omega)$  be a **transitive** permutation group, where  $|\Omega|$  is finite.

## Definition

A **base** for  $G$  is a subset  $\Delta$  of  $\Omega$  such that  $\bigcap_{\alpha \in \Delta} G_\alpha = 1$ .

Base size  $b(G)$ : the minimal size of a base for  $G$ .

## Examples

- $G = S_n$ ,  $\Omega = \{1, \dots, n\}$ :  $b(G) = n - 1$ .
- $G = \text{GL}(V)$ ,  $\Omega = V \setminus \{0\}$ :  $b(G) = \dim(V)$ .

# Bases

Let  $G \leq \text{Sym}(\Omega)$  be a **transitive** permutation group, where  $|\Omega|$  is finite.

## Definition

A **base** for  $G$  is a subset  $\Delta$  of  $\Omega$  such that  $\bigcap_{\alpha \in \Delta} G_\alpha = 1$ .

Base size  $b(G)$ : the minimal size of a base for  $G$ .

## Examples

- $G = S_n$ ,  $\Omega = \{1, \dots, n\}$ :  $b(G) = n - 1$ .
- $G = \text{GL}(V)$ ,  $\Omega = V \setminus \{0\}$ :  $b(G) = \dim(V)$ .
- $G = D_{2n}$ ,  $\Omega = \{1, \dots, n\}$ :  $b(G) = 2$ .

## Base-two groups

**Observation:**  $b(G) = 1 \iff G$  is regular.

## Base-two groups

**Observation:**  $b(G) = 1 \iff G$  is regular.

Let  $H = G_\alpha \neq 1$  be a point stabiliser.

$$\begin{aligned} b(G) = 2 &\iff G \text{ has a regular suborbit on } \Omega \\ &\iff H \cap H^g = 1 \text{ for some } g \in G \end{aligned}$$

## Base-two groups

**Observation:**  $b(G) = 1 \iff G$  is regular.

Let  $H = G_\alpha \neq 1$  be a point stabiliser.

$$\begin{aligned} b(G) = 2 &\iff G \text{ has a regular suborbit on } \Omega \\ &\iff H \cap H^g = 1 \text{ for some } g \in G \end{aligned}$$

**Recall.**  $G$  is called **primitive** if  $H$  is maximal in  $G$ .

## Base-two groups

**Observation:**  $b(G) = 1 \iff G$  is regular.

Let  $H = G_\alpha \neq 1$  be a point stabiliser.

$$\begin{aligned} b(G) = 2 &\iff G \text{ has a regular suborbit on } \Omega \\ &\iff H \cap H^g = 1 \text{ for some } g \in G \end{aligned}$$

**Recall.**  $G$  is called **primitive** if  $H$  is maximal in  $G$ .

**Problem.** Classify the finite primitive groups  $G$  with  $b(G) = 2$ .

## Base-two groups

**Observation:**  $b(G) = 1 \iff G$  is regular.

Let  $H = G_\alpha \neq 1$  be a point stabiliser.

$$\begin{aligned} b(G) = 2 &\iff G \text{ has a regular suborbit on } \Omega \\ &\iff H \cap H^g = 1 \text{ for some } g \in G \end{aligned}$$

**Recall.**  $G$  is called **primitive** if  $H$  is maximal in  $G$ .

**Problem.** Classify the finite primitive groups  $G$  with  $b(G) = 2$ .

### Example

$p$  prime,  $G = D_{2p}$  and  $\Omega = \{1, \dots, p\} \implies G$  primitive and  $b(G) = 2$ .



## Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

## Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

**Affine:** ask Melissa.

## Some results

The **O’Nan-Scott Theorem** divides finite primitive groups into 5 types.

**Affine:** ask Melissa.

**Almost simple:**  $T \leq G \leq \text{Aut}(T)$  for non-abelian simple  $T$ .

## Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

**Affine:** ask Melissa.

**Almost simple:**  $T \leq G \leq \text{Aut}(T)$  for non-abelian simple  $T$ .

- **Burness et al., 2010/11:**  $T$  alternating or sporadic ✓

## Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

**Affine:** ask Melissa.

**Almost simple:**  $T \leq G \leq \text{Aut}(T)$  for non-abelian simple  $T$ .

- **Burness et al., 2010/11:**  $T$  alternating or sporadic ✓
- $G$  Lie type: Partial answers

## Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

**Affine:** ask Melissa.

**Almost simple:**  $T \leq G \leq \text{Aut}(T)$  for non-abelian simple  $T$ .

- **Burness et al., 2010/11:**  $T$  alternating or sporadic ✓
- $G$  Lie type: Partial answers

**Diagonal and twisted wreath types:** Partial results (**Fawcett 2013/22**)

## Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

**Affine:** ask Melissa.

**Almost simple:**  $T \leq G \leq \text{Aut}(T)$  for non-abelian simple  $T$ .

- **Burness et al., 2010/11:**  $T$  alternating or sporadic ✓
- $G$  Lie type: Partial answers

**Diagonal and twisted wreath types:** Partial results (**Fawcett 2013/22**)

e.g.  $G \leq T^k.(\text{Out}(T) \times P)$  with  $P \neq A_k, S_k \implies b(G) = 2$ .

## Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

**Affine:** ask Melissa.

**Almost simple:**  $T \leq G \leq \text{Aut}(T)$  for non-abelian simple  $T$ .

- **Burness et al., 2010/11:**  $T$  alternating or sporadic ✓
- $G$  Lie type: Partial answers

**Diagonal and twisted wreath types:** Partial results (**Fawcett 2013/22**)

e.g.  $G \leq T^k.(\text{Out}(T) \times P)$  with  $P \neq A_k, S_k \implies b(G) = 2$ .

**H, 2022+:** If  $P = A_k$ , then  $b(G) = 2 \iff 2 < k < |T|$ .



## Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

**Affine:** ask Melissa.

**Almost simple:**  $T \leq G \leq \text{Aut}(T)$  for non-abelian simple  $T$ .

- **Burness et al., 2010/11:**  $T$  alternating or sporadic ✓
- $G$  Lie type: Partial answers

**Diagonal and twisted wreath types:** Partial results (**Fawcett 2013/22**)

e.g.  $G \leq T^k.(\text{Out}(T) \times P)$  with  $P \neq A_k, S_k \implies b(G) = 2$ .

**H, 2022+:** If  $P = A_k$ , then  $b(G) = 2 \iff 2 < k < |T|$ .

**Product type:**  $G \leq L \wr P$  with its product action on  $\Gamma^k$ , where  $L \leq \text{Sym}(\Gamma)$

## Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

**Affine:** ask Melissa.

**Almost simple:**  $T \leq G \leq \text{Aut}(T)$  for non-abelian simple  $T$ .

- **Burness et al., 2010/11:**  $T$  alternating or sporadic ✓
- $G$  Lie type: Partial answers

**Diagonal and twisted wreath types:** Partial results (**Fawcett 2013/22**)

e.g.  $G \leq T^k.(\text{Out}(T) \times P)$  with  $P \neq A_k, S_k \implies b(G) = 2$ .

**H, 2022+:** If  $P = A_k$ , then  $b(G) = 2 \iff 2 < k < |T|$ .

**Product type:**  $G \leq L \wr P$  with its product action on  $\Gamma^k$ , where  $L \leq \text{Sym}(\Gamma)$

- **Bailey & Cameron, 2011:**  $b(L \wr P) = 2 \iff L$  has at least  $D(P)$  regular suborbits, where  $D(P)$  is the **distinguishing number** of  $P$ .

## Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

**Affine:** ask Melissa.

**Almost simple:**  $T \leq G \leq \text{Aut}(T)$  for non-abelian simple  $T$ .

- **Burness et al., 2010/11:**  $T$  alternating or sporadic ✓
- $G$  Lie type: Partial answers

**Diagonal and twisted wreath types:** Partial results (**Fawcett 2013/22**)

e.g.  $G \leq T^k.(\text{Out}(T) \times P)$  with  $P \neq A_k, S_k \implies b(G) = 2$ .

**H, 2022+:** If  $P = A_k$ , then  $b(G) = 2 \iff 2 < k < |T|$ .

**Product type:**  $G \leq L \wr P$  with its product action on  $\Gamma^k$ , where  $L \leq \text{Sym}(\Gamma)$

- **Bailey & Cameron, 2011:**  $b(L \wr P) = 2 \iff L$  has at least  $D(P)$  regular suborbits, where  $D(P)$  is the **distinguishing number** of  $P$ .
- Progress where  $G < L \wr P$  (**Burness & H, 2022+**)

# Probabilistic methods

Consider

$$Q(G) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_\alpha \cap G_\beta \neq 1\}|}{|\Omega|^2},$$

the probability that a random pair in  $\Omega$  is **not** a base for  $G$ .

# Probabilistic methods

Consider

$$Q(G) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_\alpha \cap G_\beta \neq 1\}|}{|\Omega|^2},$$

the probability that a random pair in  $\Omega$  is **not** a base for  $G$ .

**Note.**  $Q(G) < 1 \iff b(G) \leq 2$ .

# Probabilistic methods

Consider

$$Q(G) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_\alpha \cap G_\beta \neq 1\}|}{|\Omega|^2},$$

the probability that a random pair in  $\Omega$  is **not** a base for  $G$ .

**Note.**  $Q(G) < 1 \iff b(G) \leq 2$ .

**Chen & H, 2022:** General method to calculate  $Q(G)$ .

# Probabilistic methods

Consider

$$Q(G) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_\alpha \cap G_\beta \neq 1\}|}{|\Omega|^2},$$

the probability that a random pair in  $\Omega$  is **not** a base for  $G$ .

**Note.**  $Q(G) < 1 \iff b(G) \leq 2$ .

**Chen & H, 2022:** General method to calculate  $Q(G)$ .

To calculate exact  $Q(G)$  is difficult, but we have

$$Q(G) < \sum_{x \in \mathcal{P}} \frac{|x^G \cap G_\alpha|}{|x^G|} =: \hat{Q}(G),$$

where  $\mathcal{P}$  is the set of elements of prime order in  $G$ .

# Probabilistic methods

Consider

$$Q(G) = \frac{|\{(\alpha, \beta) \in \Omega^2 : G_\alpha \cap G_\beta \neq 1\}|}{|\Omega|^2},$$

the probability that a random pair in  $\Omega$  is **not** a base for  $G$ .

**Note.**  $Q(G) < 1 \iff b(G) \leq 2$ .

**Chen & H, 2022:** General method to calculate  $Q(G)$ .

To calculate exact  $Q(G)$  is difficult, but we have

$$Q(G) < \sum_{x \in \mathcal{P}} \frac{|x^G \cap G_\alpha|}{|x^G|} =: \hat{Q}(G),$$

where  $\mathcal{P}$  is the set of elements of prime order in  $G$ .

**Probabilistic method:**  $\hat{Q}(G) < 1 \implies b(G) \leq 2$ .



# Saxl graphs

Assume  $G \leq \text{Sym}(\Omega)$  is transitive of degree  $n$  and  $b(G) = 2$ .

**Burness & Giudici, 2020: Saxl graph**  $\Sigma(G)$ :

vertices  $\Omega$ , with  $\alpha \sim \beta \iff \{\alpha, \beta\}$  is a base.

# Saxl graphs

Assume  $G \leq \text{Sym}(\Omega)$  is transitive of degree  $n$  and  $b(G) = 2$ .

**Burness & Giudici, 2020: Saxl graph**  $\Sigma(G)$ :

vertices  $\Omega$ , with  $\alpha \sim \beta \iff \{\alpha, \beta\}$  is a base.

## Example

- $G$  is **Frobenius**  $\iff \Sigma(G) \cong K_n$ .

# Saxl graphs

Assume  $G \leq \text{Sym}(\Omega)$  is transitive of degree  $n$  and  $b(G) = 2$ .

**Burness & Giudici, 2020: Saxl graph**  $\Sigma(G)$ :

vertices  $\Omega$ , with  $\alpha \sim \beta \iff \{\alpha, \beta\}$  is a base.

## Example

- $G$  is **Frobenius**  $\iff \Sigma(G) \cong K_n$ .

e.g.  $G = D_{2p}$  and  $n = p$  prime.

# Saxl graphs

Assume  $G \leq \text{Sym}(\Omega)$  is transitive of degree  $n$  and  $b(G) = 2$ .

**Burness & Giudici, 2020: Saxl graph**  $\Sigma(G)$ :

vertices  $\Omega$ , with  $\alpha \sim \beta \iff \{\alpha, \beta\}$  is a base.

## Example

- $G$  is **Frobenius**  $\iff \Sigma(G) \cong K_n$ .

e.g.  $G = D_{2p}$  and  $n = p$  prime.

- $G = \text{GL}_2(q)$  and  $\Omega = \mathbb{F}_q^2 \setminus \{0\}$ .

# Saxl graphs

Assume  $G \leq \text{Sym}(\Omega)$  is transitive of degree  $n$  and  $b(G) = 2$ .

**Burness & Giudici, 2020: Saxl graph**  $\Sigma(G)$ :

vertices  $\Omega$ , with  $\alpha \sim \beta \iff \{\alpha, \beta\}$  is a base.

## Example

- $G$  is **Frobenius**  $\iff \Sigma(G) \cong K_n$ .

e.g.  $G = D_{2p}$  and  $n = p$  prime.

- $G = \text{GL}_2(q)$  and  $\Omega = \mathbb{F}_q^2 \setminus \{0\}$ .

**Note.**  $\{\alpha, \beta\}$  is a base  $\iff \{\alpha, \beta\}$  is a basis for  $\mathbb{F}_q^2$ .

# Saxl graphs

Assume  $G \leq \text{Sym}(\Omega)$  is transitive of degree  $n$  and  $b(G) = 2$ .

**Burness & Giudici, 2020:** **Saxl graph**  $\Sigma(G)$ :

vertices  $\Omega$ , with  $\alpha \sim \beta \iff \{\alpha, \beta\}$  is a base.

## Example

- $G$  is **Frobenius**  $\iff \Sigma(G) \cong K_n$ .

e.g.  $G = D_{2p}$  and  $n = p$  prime.

- $G = \text{GL}_2(q)$  and  $\Omega = \mathbb{F}_q^2 \setminus \{0\}$ .

**Note.**  $\{\alpha, \beta\}$  is a base  $\iff \{\alpha, \beta\}$  is a basis for  $\mathbb{F}_q^2$ .

Hence,  $\Sigma(G) \cong K_{q^2-1} - (q+1)K_{q-1}$  is **complete multipartite**.

## Further example

- $G = \mathrm{PGL}_2(q)$  and  $\Omega = \{2\text{-subsets of } \{1\text{-spaces in } \mathbb{F}_q^2\}\}$ .

## Further example

- $G = \mathrm{PGL}_2(q)$  and  $\Omega = \{2\text{-subsets of } \{1\text{-spaces in } \mathbb{F}_q^2\}\}$ .

**Note.**  $G_\alpha \cong D_{2(q-1)}$  and  $\{\alpha, \beta\}$  is a base  $\iff |\alpha \cap \beta| = 1$ .



## Further example

- $G = \mathrm{PGL}_2(q)$  and  $\Omega = \{2\text{-subsets of } \{1\text{-spaces in } \mathbb{F}_q^2\}\}$ .

**Note.**  $G_\alpha \cong D_{2(q-1)}$  and  $\{\alpha, \beta\}$  is a base  $\iff |\alpha \cap \beta| = 1$ .

Hence,  $\Sigma(G) \cong J(q+1, 2)$  is a **Johnson graph**.


## Further example

- $G = \text{PGL}_2(q)$  and  $\Omega = \{2\text{-subsets of } \{1\text{-spaces in } \mathbb{F}_q^2\}\}$ .

**Note.**  $G_\alpha \cong D_{2(q-1)}$  and  $\{\alpha, \beta\}$  is a base  $\iff |\alpha \cap \beta| = 1$ .

Hence,  $\Sigma(G) \cong J(q+1, 2)$  is a **Johnson graph**.

For example, when  $q = 4$  we have the complement of the Petersen.



# Neighbours

Let  $\Sigma(\alpha)$  be the set of neighbours of  $\alpha$  in  $\Sigma(G)$ .

# Neighbours

Let  $\Sigma(\alpha)$  be the set of neighbours of  $\alpha$  in  $\Sigma(G)$ .

## Observations:

- $\Sigma(\alpha)$  is the union of regular  $G_\alpha$ -orbits.

# Neighbours

Let  $\Sigma(\alpha)$  be the set of neighbours of  $\alpha$  in  $\Sigma(G)$ .

## Observations:

- $\Sigma(\alpha)$  is the union of regular  $G_\alpha$ -orbits.
- $\Sigma(G)$  is the union of regular **orbital graphs** of  $G$ .

# Neighbours

Let  $\Sigma(\alpha)$  be the set of neighbours of  $\alpha$  in  $\Sigma(G)$ .

## Observations:

- $\Sigma(\alpha)$  is the union of regular  $G_\alpha$ -orbits.
- $\Sigma(G)$  is the union of regular **orbital graphs** of  $G$ .
- $G$  is primitive  $\implies \Sigma(G)$  is connected.

# Neighbours

Let  $\Sigma(\alpha)$  be the set of neighbours of  $\alpha$  in  $\Sigma(G)$ .

## Observations:

- $\Sigma(\alpha)$  is the union of regular  $G_\alpha$ -orbits.
- $\Sigma(G)$  is the union of regular **orbital graphs** of  $G$ .
- $G$  is primitive  $\implies \Sigma(G)$  is connected.

**Question:** What is the diameter of  $\Sigma(G)$  if  $G$  is primitive?

# Neighbours

Let  $\Sigma(\alpha)$  be the set of neighbours of  $\alpha$  in  $\Sigma(G)$ .

## Observations:

- $\Sigma(\alpha)$  is the union of regular  $G_\alpha$ -orbits.
- $\Sigma(G)$  is the union of regular **orbital graphs** of  $G$ .
- $G$  is primitive  $\implies \Sigma(G)$  is connected.

**Question:** What is the diameter of  $\Sigma(G)$  if  $G$  is primitive?

**Conjecture (Burness & Giudici, 2020)**

$G$  primitive and  $\alpha, \beta \in \Omega \implies \Sigma(\alpha) \cap \Sigma(\beta) \neq \emptyset$ .



# Probabilistic methods

Conjecture (Burness & Giudici, 2020)

$G$  primitive and  $\alpha, \beta \in \Omega \implies \Sigma(\alpha) \cap \Sigma(\beta) \neq \emptyset$ .

# Probabilistic methods

## Conjecture (Burness & Giudici, 2020)

$G$  primitive and  $\alpha, \beta \in \Omega \implies \Sigma(\alpha) \cap \Sigma(\beta) \neq \emptyset$ .

Recall that

$$Q(G) = 1 - \frac{|\Sigma(\alpha)|}{|\Omega|}$$

is the probability that a random pair in  $\Omega$  is not a base for  $G$ .

# Probabilistic methods

## Conjecture (Burness & Giudici, 2020)

$G$  primitive and  $\alpha, \beta \in \Omega \implies \Sigma(\alpha) \cap \Sigma(\beta) \neq \emptyset$ .

Recall that

$$Q(G) = 1 - \frac{|\Sigma(\alpha)|}{|\Omega|}$$

is the probability that a random pair in  $\Omega$  is not a base for  $G$ .

### Observations:

- $Q(G) < 1 \iff b(G) \leq 2$ .

# Probabilistic methods

## Conjecture (Burness & Giudici, 2020)

$G$  primitive and  $\alpha, \beta \in \Omega \implies \Sigma(\alpha) \cap \Sigma(\beta) \neq \emptyset$ .

Recall that

$$Q(G) = 1 - \frac{|\Sigma(\alpha)|}{|\Omega|}$$

is the probability that a random pair in  $\Omega$  is not a base for  $G$ .

### Observations:

- $Q(G) < 1 \iff b(G) \leq 2$ .
- $Q(G) < 1/2 \iff |\Sigma(\alpha)| > \frac{1}{2}|\Omega|$

# Probabilistic methods

## Conjecture (Burness & Giudici, 2020)

$G$  primitive and  $\alpha, \beta \in \Omega \implies \Sigma(\alpha) \cap \Sigma(\beta) \neq \emptyset$ .

Recall that

$$Q(G) = 1 - \frac{|\Sigma(\alpha)|}{|\Omega|}$$

is the probability that a random pair in  $\Omega$  is not a base for  $G$ .

### Observations:

- $Q(G) < 1 \iff b(G) \leq 2$ .
- $Q(G) < 1/2 \iff |\Sigma(\alpha)| > \frac{1}{2}|\Omega| \implies \Sigma(\alpha) \cap \Sigma(\beta) \neq \emptyset$ .

# Burness-Giudici conjecture

## Example

If  $G = \mathrm{PGL}_2(q)$  and  $G_\alpha = D_{2(q-1)}$ , then  $\Sigma(G) = J(q+1, 2)$  has the common neighbour property, though  $Q(G) \rightarrow 1$  as  $q \rightarrow \infty$ .

# Burness-Giudici conjecture

## Example

If  $G = \text{PGL}_2(q)$  and  $G_\alpha = D_{2(q-1)}$ , then  $\Sigma(G) = J(q+1, 2)$  has the common neighbour property, though  $Q(G) \rightarrow 1$  as  $q \rightarrow \infty$ .

Some other evidence:

- **Chen & Du 2020+; Burness & H, 2021+:**  $\text{soc}(G) = L_2(q)$  ✓

# Burness-Giudici conjecture

## Example

If  $G = \mathrm{PGL}_2(q)$  and  $G_\alpha = D_{2(q-1)}$ , then  $\Sigma(G) = J(q+1, 2)$  has the common neighbour property, though  $Q(G) \rightarrow 1$  as  $q \rightarrow \infty$ .

Some other evidence:

- **Chen & Du 2020+; Burness & H, 2021+:**  $\mathrm{soc}(G) = L_2(q)$  ✓
- **Burness & H, 2021+:**  $G$  almost simple with  $G_\alpha$  soluble ✓



# Burness-Giudici conjecture

## Example

If  $G = \mathrm{PGL}_2(q)$  and  $G_\alpha = D_{2(q-1)}$ , then  $\Sigma(G) = J(q+1, 2)$  has the common neighbour property, though  $Q(G) \rightarrow 1$  as  $q \rightarrow \infty$ .

Some other evidence:

- **Chen & Du 2020+; Burness & H, 2021+:**  $\mathrm{soc}(G) = L_2(q)$  ✓
- **Burness & H, 2021+:**  $G$  almost simple with  $G_\alpha$  soluble ✓
- **Lee & Popiel, 2021+:** some affine groups

# Burness-Giudici conjecture

## Example

If  $G = \mathrm{PGL}_2(q)$  and  $G_\alpha = D_{2(q-1)}$ , then  $\Sigma(G) = J(q+1, 2)$  has the common neighbour property, though  $Q(G) \rightarrow 1$  as  $q \rightarrow \infty$ .

Some other evidence:

- **Chen & Du 2020+; Burness & H, 2021+:**  $\mathrm{soc}(G) = L_2(q)$  ✓
- **Burness & H, 2021+:**  $G$  almost simple with  $G_\alpha$  soluble ✓
- **Lee & Popiel, 2021+:** some affine groups

**Recall.**  $\Sigma(\alpha)$  is the union of regular  $G_\alpha$ -orbits.

# Burness-Giudici conjecture

## Example

If  $G = \mathrm{PGL}_2(q)$  and  $G_\alpha = D_{2(q-1)}$ , then  $\Sigma(G) = J(q+1, 2)$  has the common neighbour property, though  $Q(G) \rightarrow 1$  as  $q \rightarrow \infty$ .

Some other evidence:

- **Chen & Du 2020+; Burness & H, 2021+:**  $\mathrm{soc}(G) = L_2(q)$  ✓
- **Burness & H, 2021+:**  $G$  almost simple with  $G_\alpha$  soluble ✓
- **Lee & Popiel, 2021+:** some affine groups

**Recall.**  $\Sigma(\alpha)$  is the union of regular  $G_\alpha$ -orbits.

**BG conjecture:**  $\Sigma(\alpha)$  meets **some** regular  $G_\beta$ -orbits.

# Burness-Giudici conjecture

## Example

If  $G = \mathrm{PGL}_2(q)$  and  $G_\alpha = D_{2(q-1)}$ , then  $\Sigma(G) = J(q+1, 2)$  has the common neighbour property, though  $Q(G) \rightarrow 1$  as  $q \rightarrow \infty$ .

Some other evidence:

- **Chen & Du 2020+; Burness & H, 2021+:**  $\mathrm{soc}(G) = L_2(q)$  ✓
- **Burness & H, 2021+:**  $G$  almost simple with  $G_\alpha$  soluble ✓
- **Lee & Popiel, 2021+:** some affine groups

**Recall.**  $\Sigma(\alpha)$  is the union of regular  $G_\alpha$ -orbits.

**BG conjecture:**  $\Sigma(\alpha)$  meets **some** regular  $G_\beta$ -orbits.

**Conjecture (Burness & H, 2022+)**

$\Sigma(\alpha)$  meets **every** regular  $G_\beta$ -orbit.

# Future work

## Saxl graphs:

- Other invariants

# Future work

## Saxl graphs:

- Other invariants

**Burness & H, 2021+:** Results on clique and independence numbers

**Chen & H, 2022:** Some results on valencies

# Future work

## Saxl graphs:

- Other invariants

**Burness & H, 2021+:** Results on clique and independence numbers

**Chen & H, 2022:** Some results on valencies

- Some generalisations

# Future work

## Saxl graphs:

- Other invariants

**Burness & H, 2021+:** Results on clique and independence numbers

**Chen & H, 2022:** Some results on valencies

- Some generalisations

## The base-two project:

- Affine groups  $G = VH$



# Future work

## Saxl graphs:

- Other invariants

**Burness & H, 2021+:** Results on clique and independence numbers

**Chen & H, 2022:** Some results on valencies

- Some generalisations

## The base-two project:

- Affine groups  $G = VH$

**Seress, 1996:**  $b(G) \leq 4$  if  $H$  is soluble

# Future work

## Saxl graphs:

- Other invariants

**Burness & H, 2021+:** Results on clique and independence numbers

**Chen & H, 2022:** Some results on valencies

- Some generalisations

## The base-two project:

- Affine groups  $G = VH$

**Seress, 1996:**  $b(G) \leq 4$  if  $H$  is soluble

**Halasi & Podoski, 2016:**  $b(G) \leq 3$  if  $(|V|, |H|) = 1$

# Future work

## Saxl graphs:

- Other invariants

**Burness & H, 2021+:** Results on clique and independence numbers

**Chen & H, 2022:** Some results on valencies

- Some generalisations

## The base-two project:

- Affine groups  $G = VH$

**Seress, 1996:**  $b(G) \leq 4$  if  $H$  is soluble

**Halasi & Podoski, 2016:**  $b(G) \leq 3$  if  $(|V|, |H|) = 1$

- Diagonal type groups

**Thank you!**