

Base-two primitive permutation groups

Hong Yi Huang

23rd Postgraduate Group Theory Conference

8 July 2022



Let $G \leq \text{Sym}(\Omega)$ be a permutation group.

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G .

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Question: For a subset $\Delta \subseteq \Omega$, when do we have

$$G_{(\Delta)} := \bigcap_{\alpha \in \Delta} G_\alpha = 1?$$

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Question: For a subset $\Delta \subseteq \Omega$, when do we have

$$G_{(\Delta)} := \bigcap_{\alpha \in \Delta} G_\alpha = 1?$$

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$ and $\Delta = \{1, \dots, n-1\}$.

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Denote by

$$G_\alpha = \{g \in G \mid \alpha^g = \alpha\}$$

the **stabiliser** of α in G . Then

$$\bigcap_{\alpha \in \Omega} G_\alpha = 1.$$

Question: For a subset $\Delta \subseteq \Omega$, when do we have

$$G_{(\Delta)} := \bigcap_{\alpha \in \Delta} G_\alpha = 1?$$

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$ and $\Delta = \{1, \dots, n-1\}$.
- $G = \text{GL}(V)$, $\Omega = V \setminus \{0\}$ and Δ contains a basis of V .

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

From now on, we will assume Ω is finite.

Base size $b(G)$: the minimal size of a base for G .

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

From now on, we will assume Ω is finite.

Base size $b(G)$: the minimal size of a base for G .

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 1$.

Bases

Definition

A **base** for $G \leq \text{Sym}(\Omega)$ is a subset Δ of Ω such that $G_{(\Delta)} = 1$.

From now on, we will assume Ω is finite.

Base size $b(G)$: the minimal size of a base for G .

Examples

- $G = S_n$, $\Omega = \{1, \dots, n\}$: $b(G) = n - 1$.
- $G = \text{GL}(V)$, $\Omega = V \setminus \{0\}$: $b(G) = \dim(V)$.

Base sizes

Observation. If Δ is a base and $x, y \in G$, then

$$\alpha^x = \alpha^y \text{ for all } \alpha \in \Delta \iff xy^{-1} \in \bigcap_{\alpha \in \Delta} G_\alpha \iff x = y.$$

Base sizes

Observation. If Δ is a base and $x, y \in G$, then

$$\alpha^x = \alpha^y \text{ for all } \alpha \in \Delta \iff xy^{-1} \in \bigcap_{\alpha \in \Delta} G_\alpha \iff x = y.$$

That is, each group element is uniquely determined by its action on Δ .

Base sizes

Observation. If Δ is a base and $x, y \in G$, then

$$\alpha^x = \alpha^y \text{ for all } \alpha \in \Delta \iff xy^{-1} \in \bigcap_{\alpha \in \Delta} G_\alpha \iff x = y.$$

That is, each group element is uniquely determined by its action on Δ .

- A small base Δ provides an efficient way to store the elements of G , using $|\Delta|$ -tuples rather than $|\Omega|$ -tuples.

Base sizes

Observation. If Δ is a base and $x, y \in G$, then

$$\alpha^x = \alpha^y \text{ for all } \alpha \in \Delta \iff xy^{-1} \in \bigcap_{\alpha \in \Delta} G_\alpha \iff x = y.$$

That is, each group element is uniquely determined by its action on Δ .

- A small base Δ provides an efficient way to store the elements of G , using $|\Delta|$ -tuples rather than $|\Omega|$ -tuples.

Question. How small can a base be?

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

G is called **primitive** if G_α is maximal in G .

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

G is called **primitive** if G_α is maximal in G .

Problem. Classify the finite primitive groups G with $b(G) = 2$.

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

G is called **primitive** if G_α is maximal in G .

Problem. Classify the finite primitive groups G with $b(G) = 2$.

Example

Consider the action of $G = D_{2n}$ on $\{1, \dots, n\}$. Then

- $\{1, 2\}$ is a base, so $b(G) = 2$;

Base-two groups

Observations: If G is transitive, then

- $b(G) = 1 \iff G$ is regular;
- $b(G) = 2 \iff G_\alpha \neq 1$ has a regular orbit.

G is called **primitive** if G_α is maximal in G .

Problem. Classify the finite primitive groups G with $b(G) = 2$.

Example

Consider the action of $G = D_{2n}$ on $\{1, \dots, n\}$. Then

- $\{1, 2\}$ is a base, so $b(G) = 2$;
- G is primitive iff n is a prime.

Some results

The **O’Nan-Scott Theorem** divides finite primitive groups into 5 types.

Some results

The **O’Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Some results

The **O’Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem. Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

Some results

The **O’Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem. Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem. Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem. Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question. Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

Some results

The **O’Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem. Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question. Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- CFSG is used. Partial results (**Burness et al.**)

Some results

The **O'Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem. Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question. Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- CFSG is used. Partial results (**Burness et al.**)

Diagonal and twisted wreath types: Partial results (**Fawcett 2013/22**)

Some results

The **O’Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem. Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question. Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- CFSG is used. Partial results (**Burness et al.**)

Diagonal and twisted wreath types: Partial results (**Fawcett 2013/22**)

Product type: $G \leq L \wr P$ with its product action on Γ^k , where $L \leq \text{Sym}(\Gamma)$

Some results

The **O’Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem. Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question. Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- CFSG is used. Partial results (**Burness et al.**)

Diagonal and twisted wreath types: Partial results (**Fawcett 2013/22**)

Product type: $G \leq L \wr P$ with its product action on Γ^k , where $L \leq \text{Sym}(\Gamma)$

- **Bailey & Cameron, 2013:** Explicit $b(L \wr P)$ ✓

Some results

The **O’Nan-Scott Theorem** divides finite primitive groups into 5 types.

Affine: $G = V:H \leq \text{AGL}(V)$, where $V = \mathbb{F}_p^d$ and $H \leq \text{GL}(V)$ irreducible.

Problem. Determine the pairs (H, V) , where H is a finite group, V is a faithful irreducible $\mathbb{F}_p H$ -module and H has a regular orbit on V .

- Partial results (e.g. $H/Z(H)$ is simple).

Almost simple: $T \leq G \leq \text{Aut}(T)$ for non-abelian simple T .

Question. Let H be a maximal subgroup of G . When is $H \cap H^g = 1$?

- CFSG is used. Partial results (**Burness et al.**)

Diagonal and twisted wreath types: Partial results (**Fawcett 2013/22**)

Product type: $G \leq L \wr P$ with its product action on Γ^k , where $L \leq \text{Sym}(\Gamma)$

- **Bailey & Cameron, 2013:** Explicit $b(L \wr P)$ ✓
- Progress where G_α is soluble and $G < L \wr P$ (**Burness & H, 2022+**)

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Examples

- $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$:


Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Examples

- $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 


Saxl graphs

Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Examples

- $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 
- $G = C_4 = \langle (1234) \rangle$, $\Omega = \{1, 2, 3, 4\}$:


Saxl graphs


Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Examples

• $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 

• $G = C_4 = \langle (1234) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 


Saxl graphs


Definition (Burness & Giudici, 2020)

Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Examples

- $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 

- $G = C_4 = \langle (1234) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 

- $G = D_{10} = \langle (12345), (25)(34) \rangle$, $\Omega = \{1, 2, 3, 4, 5\}$:


Saxl graphs


Definition (Burness & Giudici, 2020)


Let $G \leq \text{Sym}(\Omega)$. Then the **Saxl graph** $\Sigma(G)$ is a graph with

- vertex set Ω ;
- α and β are adjacent $\iff \{\alpha, \beta\}$ is a base for G .

Examples

• $G = D_8 = \langle (1234), (24) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 

• $G = C_4 = \langle (1234) \rangle$, $\Omega = \{1, 2, 3, 4\}$: $\Sigma(G) =$ 

• $G = D_{10} = \langle (12345), (25)(34) \rangle$, $\Omega = \{1, 2, 3, 4, 5\}$: $\Sigma(G) =$ 

Another example

Let $G = \mathrm{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 .

- $G_\alpha = D_{2(q-1)}$;
- α and β form a base iff they share a common 1-space.

Another example

Let $G = \mathrm{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 .

- $G_\alpha = D_{2(q-1)}$;
- α and β form a base iff they share a common 1-space.

Hence, $\Sigma(G) \cong J(q+1, 2)$ is a **Johnson graph**: vertices 2-subsets of $\{1, \dots, q+1\}$ and two vertices are adjacent if they are not disjoint.


Another example

Let $G = \text{PGL}_2(q)$ and Ω be the set of distinct pairs of 1-spaces in \mathbb{F}_q^2 .

- $G_\alpha = D_{2(q-1)}$;
- α and β form a base iff they share a common 1-space.

Hence, $\Sigma(G) \cong J(q+1, 2)$ is a **Johnson graph**: vertices 2-subsets of $\{1, \dots, q+1\}$ and two vertices are adjacent if they are not disjoint.

For example, when $q = 4$ we have the complement of the Petersen graph.



Valency

Assume G is transitive and $b(G) = 2$.

Valency

Assume G is transitive and $b(G) = 2$.

Notes.

- $\Sigma(G)$ is vertex-transitive.

Valency

Assume G is transitive and $b(G) = 2$.

Notes.

- $\Sigma(G)$ is vertex-transitive.
- $\Sigma(G)$ has valency $r(G)|G_\alpha|$, where $r(G)$ is the number of regular G_α -orbits.

Valency

Assume G is transitive and $b(G) = 2$.

Notes.

- $\Sigma(G)$ is vertex-transitive.
- $\Sigma(G)$ has valency $r(G)|G_\alpha|$, where $r(G)$ is the number of regular G_α -orbits.

Chen & H, 2022: A general method for computing $r(G)$.

Valency

Assume G is transitive and $b(G) = 2$.

Notes.

- $\Sigma(G)$ is vertex-transitive.
- $\Sigma(G)$ has valency $r(G)|G_\alpha|$, where $r(G)$ is the number of regular G_α -orbits.

Chen & H, 2022: A general method for computing $r(G)$.

Problem. Classify the finite primitive groups G with $r(G) = 1$.

Valency

Assume G is transitive and $b(G) = 2$.

Notes.

- $\Sigma(G)$ is vertex-transitive.
- $\Sigma(G)$ has valency $r(G)|G_\alpha|$, where $r(G)$ is the number of regular G_α -orbits.

Chen & H, 2022: A general method for computing $r(G)$.

Problem. Classify the finite primitive groups G with $r(G) = 1$.

Burness & H, 2021+: G almost simple, G_α soluble and $r(G) = 1$ ✓

Valency

Assume G is transitive and $b(G) = 2$.

Notes.

- $\Sigma(G)$ is vertex-transitive.
- $\Sigma(G)$ has valency $r(G)|G_\alpha|$, where $r(G)$ is the number of regular G_α -orbits.

Chen & H, 2022: A general method for computing $r(G)$.

Problem. Classify the finite primitive groups G with $r(G) = 1$.

Burness & H, 2021+: G almost simple, G_α soluble and $r(G) = 1$ ✓

e.g. $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$

Valency

Assume G is transitive and $b(G) = 2$.

Notes.

- $\Sigma(G)$ is vertex-transitive.
- $\Sigma(G)$ has valency $r(G)|G_\alpha|$, where $r(G)$ is the number of regular G_α -orbits.

Chen & H, 2022: A general method for computing $r(G)$.

Problem. Classify the finite primitive groups G with $r(G) = 1$.

Burness & H, 2021+: G almost simple, G_α soluble and $r(G) = 1$ ✓

e.g. $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$

Remark. $r(G) = 1 \iff \Sigma(G)$ is an **orbital graph**.

Diameter

Notes.

- $\Sigma(G)$ is the union of the $r(G)$ regular orbital graphs of G .

Diameter

Notes.

- $\Sigma(G)$ is the union of the $r(G)$ regular orbital graphs of G .
- G is primitive $\implies \Sigma(G)$ is connected.

Diameter

Notes.

- $\Sigma(G)$ is the union of the $r(G)$ regular orbital graphs of G .
- G is primitive $\implies \Sigma(G)$ is connected.

Question: What is the diameter of $\Sigma(G)$ if G is primitive?

Diameter

Notes.

- $\Sigma(G)$ is the union of the $r(G)$ regular orbital graphs of G .
- G is primitive $\implies \Sigma(G)$ is connected.

Question: What is the diameter of $\Sigma(G)$ if G is primitive?

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

Diameter

Notes.

- $\Sigma(G)$ is the union of the $r(G)$ regular orbital graphs of G .
- G is primitive $\implies \Sigma(G)$ is connected.

Question: What is the diameter of $\Sigma(G)$ if G is primitive?

Conjecture (Burness & Giudici, 2020)

G primitive \implies any two vertices in $\Sigma(G)$ have a common neighbour.

In particular, it asserts that $\Sigma(G)$ has diameter at most 2.

Diameter

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Diameter

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓

Diameter

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓
- “Most” almost simple groups with alternating or sporadic socles

Diameter

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓
- “Most” almost simple groups with alternating or sporadic socles
- **Chen & Du, 2020+; Burness & H, 2021+:** $\text{soc}(G) = L_2(q)$ ✓

Diameter

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓
- “Most” almost simple groups with alternating or sporadic socles
- **Chen & Du, 2020+; Burness & H, 2021+:** $\text{soc}(G) = L_2(q)$ ✓
- **Burness & H, 2021+:** almost simple groups with G_α soluble ✓

Diameter

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $\Sigma(G) = J(q+1, 2)$ has the common neighbour property.

Evidence:

- All primitive groups of degree up to 4095 ✓
- “Most” almost simple groups with alternating or sporadic socles
- **Chen & Du, 2020+; Burness & H, 2021+:** $\text{soc}(G) = L_2(q)$ ✓
- **Burness & H, 2021+:** almost simple groups with G_α soluble ✓
- **Lee & Popiel, 2021+:** some affine groups

Other invariants

$\mathcal{B} := \{\text{base-two almost simple primitive groups with soluble stabilisers}\}$

Other invariants

$\mathcal{B} := \{\text{base-two almost simple primitive groups with soluble stabilisers}\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓

Other invariants

$\mathcal{B} := \{\text{base-two almost simple primitive groups with soluble stabilisers}\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓
- **Burness & H, 2021+:** The common neighbour property for \mathcal{B} ✓

Other invariants

$\mathcal{B} := \{\text{base-two almost simple primitive groups with soluble stabilisers}\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓
- **Burness & H, 2021+:** The common neighbour property for \mathcal{B} ✓

Clique number: maximal size $\omega(G)$ of a complete subgraph of $\Sigma(G)$.

Other invariants

$\mathcal{B} := \{\text{base-two almost simple primitive groups with soluble stabilisers}\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓
- **Burness & H, 2021+:** The common neighbour property for \mathcal{B} ✓

Clique number: maximal size $\omega(G)$ of a complete subgraph of $\Sigma(G)$.

Independence number: clique number $\alpha(G)$ of the complement of $\Sigma(G)$.

Other invariants

$\mathcal{B} := \{\text{base-two almost simple primitive groups with soluble stabilisers}\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓
- **Burness & H, 2021+:** The common neighbour property for \mathcal{B} ✓

Clique number: maximal size $\omega(G)$ of a complete subgraph of $\Sigma(G)$.

Independence number: clique number $\alpha(G)$ of the complement of $\Sigma(G)$.

Example: $(G, G_\alpha) = (A_5, S_3) \implies \Sigma(G) = J(5, 2), \omega(G) = 4, \alpha(G) = 2.$

Other invariants

$\mathcal{B} := \{\text{base-two almost simple primitive groups with soluble stabilisers}\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓
- **Burness & H, 2021+:** The common neighbour property for \mathcal{B} ✓

Clique number: maximal size $\omega(G)$ of a complete subgraph of $\Sigma(G)$.

Independence number: clique number $\alpha(G)$ of the complement of $\Sigma(G)$.

Example: $(G, G_\alpha) = (A_5, S_3) \implies \Sigma(G) = J(5, 2), \omega(G) = 4, \alpha(G) = 2.$

Theorem (Burness & H, 2021+)

- $G \in \mathcal{B}$ is simple $\implies \omega(G) \geq 5$ or $(G, G_\alpha) = (A_5, S_3);$

Other invariants

$\mathcal{B} := \{\text{base-two almost simple primitive groups with soluble stabilisers}\}$

- **Burness, 2021:** \mathcal{B} is completely known ✓
- **Burness & H, 2021+:** The common neighbour property for \mathcal{B} ✓

Clique number: maximal size $\omega(G)$ of a complete subgraph of $\Sigma(G)$.

Independence number: clique number $\alpha(G)$ of the complement of $\Sigma(G)$.

Example: $(G, G_\alpha) = (A_5, S_3) \implies \Sigma(G) = J(5, 2), \omega(G) = 4, \alpha(G) = 2.$

Theorem (Burness & H, 2021+)

- $G \in \mathcal{B}$ is simple $\implies \omega(G) \geq 5$ or $(G, G_\alpha) = (A_5, S_3);$
- $G \in \mathcal{B} \implies \alpha(G) \geq 4$ or $(G, G_\alpha) = (A_5, S_3).$

Probabilistic methods

Recall that $\Sigma(G)$ has valency $v(G) = r(G)|G_\alpha|$

Probabilistic methods

Recall that $\Sigma(G)$ has valency $v(G) = r(G)|G_\alpha|$ and let

$$Q(G) = 1 - \frac{r(G)|G_\alpha|}{n} = 1 - \frac{v(G)}{n}$$

be the probability that a random pair in Ω is not a base for G .

Probabilistic methods

Recall that $\Sigma(G)$ has valency $v(G) = r(G)|G_\alpha|$ and let

$$Q(G) = 1 - \frac{r(G)|G_\alpha|}{n} = 1 - \frac{v(G)}{n}$$

be the probability that a random pair in Ω is not a base for G .

Notes:

- $Q(G) < 1 \iff b(G) \leq 2$.

Probabilistic methods

Recall that $\Sigma(G)$ has valency $v(G) = r(G)|G_\alpha|$ and let

$$Q(G) = 1 - \frac{r(G)|G_\alpha|}{n} = 1 - \frac{v(G)}{n}$$

be the probability that a random pair in Ω is not a base for G .

Notes:

- $Q(G) < 1 \iff b(G) \leq 2$.
- $Q(G) < 1/2 \implies \Sigma(G)$ has the common neighbour property.

Probabilistic methods

Recall that $\Sigma(G)$ has valency $v(G) = r(G)|G_\alpha|$ and let

$$Q(G) = 1 - \frac{r(G)|G_\alpha|}{n} = 1 - \frac{v(G)}{n}$$

be the probability that a random pair in Ω is not a base for G .

Notes:

- $Q(G) < 1 \iff b(G) \leq 2$.
- $Q(G) < 1/2 \implies \Sigma(G)$ has the common neighbour property.
- $Q(G) < 1/t \implies \omega(G) \geq t + 1$.

Probabilistic methods

Recall that $\Sigma(G)$ has valency $v(G) = r(G)|G_\alpha|$ and let

$$Q(G) = 1 - \frac{r(G)|G_\alpha|}{n} = 1 - \frac{v(G)}{n}$$

be the probability that a random pair in Ω is not a base for G .

Notes:

- $Q(G) < 1 \iff b(G) \leq 2$.
- $Q(G) < 1/2 \implies \Sigma(G)$ has the common neighbour property.
- $Q(G) < 1/t \implies \omega(G) \geq t + 1$.

Example

If $G = \text{PGL}_2(q)$ and $G_\alpha = D_{2(q-1)}$, then $Q(G) \rightarrow 1$ as $q \rightarrow \infty$. But $\Sigma(G) = J(q+1, 2)$ still has the common neighbour property.

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Conjecture (Burness & H, 2022+)

G primitive and $\alpha, \beta \in \Omega \implies \Sigma(\alpha)$ meets every regular G_β -orbit.

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Conjecture (Burness & H, 2022+)

G primitive and $\alpha, \beta \in \Omega \implies \Sigma(\alpha)$ meets every regular G_β -orbit.

Remark. BG conjecture: $\Sigma(\alpha)$ meets the union of regular G_β -orbits.

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Conjecture (Burness & H, 2022+)

G primitive and $\alpha, \beta \in \Omega \implies \Sigma(\alpha)$ meets every regular G_β -orbit.

Remark. BG conjecture: $\Sigma(\alpha)$ meets the union of regular G_β -orbits.

Theorem (Burness & H, 2022+)

BG conjecture and BH conjecture are equivalent.

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Conjecture (Burness & H, 2022+)

G primitive and $\alpha, \beta \in \Omega \implies \Sigma(\alpha)$ meets every regular G_β -orbit.

Remark. BG conjecture: $\Sigma(\alpha)$ meets the union of regular G_β -orbits.

Theorem (Burness & H, 2022+)

BG conjecture and BH conjecture are equivalent.

Indeed, G does not satisfy BH $\implies G \wr S_{r(G)}$ does not satisfy BG.

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Conjecture (Burness & H, 2022+)

G primitive and $\alpha, \beta \in \Omega \implies \Sigma(\alpha)$ meets every regular G_β -orbit.

Remark. BG conjecture: $\Sigma(\alpha)$ meets the union of regular G_β -orbits.

Theorem (Burness & H, 2022+)

BG conjecture and BH conjecture are equivalent.

Indeed, G does not satisfy BH $\implies G \wr S_{r(G)}$ does not satisfy BG.

Evidence:

- The cases $r(G) = 1$ and BG holds

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Conjecture (Burness & H, 2022+)

G primitive and $\alpha, \beta \in \Omega \implies \Sigma(\alpha)$ meets every regular G_β -orbit.

Remark. BG conjecture: $\Sigma(\alpha)$ meets the union of regular G_β -orbits.

Theorem (Burness & H, 2022+)

BG conjecture and BH conjecture are equivalent.

Indeed, G does not satisfy BH $\implies G \wr S_{r(G)}$ does not satisfy BG.

Evidence:

- The cases $r(G) = 1$ and BG holds
- All primitive groups of degree up to 4095 ✓

A strong conjecture

Let $\Sigma(\alpha)$ be the set of neighbours of α in $\Sigma(G)$.

Conjecture (Burness & H, 2022+)

G primitive and $\alpha, \beta \in \Omega \implies \Sigma(\alpha)$ meets every regular G_β -orbit.

Remark. BG conjecture: $\Sigma(\alpha)$ meets the union of regular G_β -orbits.

Theorem (Burness & H, 2022+)

BG conjecture and BH conjecture are equivalent.

Indeed, G does not satisfy BH $\implies G \wr S_{r(G)}$ does not satisfy BG.

Evidence:

- The cases $r(G) = 1$ and BG holds
- All primitive groups of degree up to 4095 ✓
- $G = \text{PSL}_2(q)$ and G_α of type $\text{GL}_1(q) \wr S_2$

Future work

Saxl graphs:

- Generalised Saxl graphs?

Future work

Saxl graphs:

- Generalised Saxl graphs?
- More evidence on BG/BH conjecture?

Future work

Saxl graphs:

- Generalised Saxl graphs?
- More evidence on BG/BH conjecture?
- Determine $r(G)$ in some special cases.

Future work

Saxl graphs:

- Generalised Saxl graphs?
- More evidence on BG/BH conjecture?
- Determine $r(G)$ in some special cases.
- Other invariants of Saxl graphs? (e.g. the spectrum)

Future work

Saxl graphs:

- Generalised Saxl graphs?
- More evidence on BG/BH conjecture?
- Determine $r(G)$ in some special cases.
- Other invariants of Saxl graphs? (e.g. the spectrum)

Diagonal type groups: $G \leq T^k.(\text{Out}(T) \times P)$

Future work

Saxl graphs:

- Generalised Saxl graphs?
- More evidence on BG/BH conjecture?
- Determine $r(G)$ in some special cases.
- Other invariants of Saxl graphs? (e.g. the spectrum)

Diagonal type groups: $G \leq T^k.(\text{Out}(T) \times P)$

Fawcett 2013: $b(G) = 2$ if $P \neq A_k, S_k$

Future work

Saxl graphs:

- Generalised Saxl graphs?
- More evidence on BG/BH conjecture?
- Determine $r(G)$ in some special cases.
- Other invariants of Saxl graphs? (e.g. the spectrum)

Diagonal type groups: $G \leq T^k.(\text{Out}(T) \times P)$

Fawcett 2013: $b(G) = 2$ if $P \neq A_k, S_k$

- Determine $b(G)$ when $P = A_k$ or S_k .

Future work

Saxl graphs:

- Generalised Saxl graphs?
- More evidence on BG/BH conjecture?
- Determine $r(G)$ in some special cases.
- Other invariants of Saxl graphs? (e.g. the spectrum)

Diagonal type groups: $G \leq T^k.(\text{Out}(T) \times P)$

Fawcett 2013: $b(G) = 2$ if $P \neq A_k, S_k$

- Determine $b(G)$ when $P = A_k$ or S_k .
- The Saxl graph $\Sigma(G)$?

Thank you!