

Intraplate earthquakes in Australia

DAN CLARK, ANDREW MCPHERSON, AND TREVOR ALLEN

Abstract

Relative to other intraplate areas of the world, Australia has a short recorded history of seismicity, spanning only a couple of centuries. As a consequence, there is significant uncertainty as to whether patterns evident in the contemporary seismic record are representative of the longer term, or constitute a bias resulting from the short sampling period. This problem can, in part, be overcome by validation against Australia's rich record of morphogenic earthquakes – Australia boasts arguably the richest late Neogene to Quaternary faulting record of any stable continental region. Long-term patterns in large earthquake occurrence, both temporal and spatial, can be deduced from the landscape record and used to inform contemporary earthquake hazard science. Seismicity source parameters such as large earthquake recurrence and magnitude vary across the Australian continent, and can be interpreted in a framework of large-scale neotectonic domains defined on the basis of geology and crustal setting. Temporal and spatial clustering of earthquakes is apparent at the scale of a single fault, and at the 1,000 km scale of a domain. The utility of the domains approach, which ties seismicity characteristics to crustal architecture and geology, is that behaviours can be extrapolated from well-characterised regions to poorly known analogous regions, both within Australia and worldwide.

2.1 Introduction

The Australian continent resides entirely within the Indo-Australian Plate, and is classified as a Stable Continental Region (SCR) in terms of its plate tectonic setting and seismicity (Johnston *et al.*, 1994; Schulte and Mooney, 2005). Such settings host approximately 0.2% of the world's seismic moment release (Johnston, 1994), and moderate to large earthquakes are rare. Analysis of focal mechanisms from SCR crust (Zoback, 1992;

Intraplate Earthquakes, ed. Pradeep Talwani. Published by Cambridge University Press. © Cambridge University Press 2014.

Reinecker *et al.*, 2003; Schulte and Mooney, 2005) indicates that compressive stress regimes dominate within continental interiors, with maximum compressive stresses oriented predominantly in accordance with absolute plate motion (see also Richardson, 1992). Australia is anomalous in this respect. While earthquake focal mechanisms suggest that the crustal stress field at seismogenic depths in Australia is everywhere compressive, with significant strike-slip components along the northwest margin (NWSSZ, Figure 2.1) and in the Flinders/Mount Lofty Ranges (FRSZ, Figure 2.1) (Leonard *et al.*, 2002; Clark *et al.*, 2003), stress orientations in southern Australia are typically not parallel to the north–northeast-directed plate motion vector (Coblentz *et al.*, 1995; Hillis and Reynolds, 2000, 2003; Sandiford *et al.*, 2004, 2005; Hillis *et al.*, 2008).

In the southern half of the continent, the maximum horizontal stress orientation (S_{Hmax}) is essentially east–west in western and central Australia and rotates to northwest–southeast in eastern Australia (Figure 2.1). In the northern half of the continent, the stress field transitions from the generally east–west trend in the south, to a broadly northeast–southwest trend. To a first order these regional stress orientations are not influenced by tectonic terrane, crustal thickness, heat flow, regional structural trends, geological age, or by the depth at which orientations are sampled (e.g., Hillis *et al.*, 2008; Sandiford and Quigley, 2009; Holford *et al.*, 2011). The trends have been satisfactorily modelled in terms of a balance between plate driving and resisting torques generated at the margins of the Indo-Australian Plate (Cloetingh and Wortel, 1986; Coblentz *et al.*, 1995, 1998; Reynolds *et al.*, 2002; Burbidge, 2004; Sandiford *et al.*, 2004; Dyksterhuis and Müller, 2008) (Figure 2.2).

Stratigraphic relationships establish that fault-related and presumably seismogenic deformation consistent with the present stress field commenced in the late Miocene, in the interval 10–6 Ma (Dickinson *et al.*, 2002; Sandiford *et al.*, 2004; Keep and Haig, 2010), as the result of complex evolving plate boundary conditions (Sandiford and Quigley, 2009). It has been proposed that the onset of deformation at specific locations may reflect rising stress levels related to the combination of all plate boundary forcings (Sandiford and Quigley, 2009), with variations in the thermal structure of the Australian lithosphere influencing the localisation of deformation at the regional or terrane scale (e.g., Celerier *et al.*, 2005; Sandiford and Egholm, 2008; Holford *et al.*, 2011). Analysis of the spatial and temporal patterns of contemporary seismicity might therefore be improved by studying the extraordinarily rich Neogene to Quaternary record of seismicity in the Australian landscape (e.g., Clark *et al.*, 2012). In the sections that follow, earthquake occurrence in Australia is therefore examined from both the historic and prehistoric (Neogene and Quaternary) records.

2.2 Two centuries of earthquake observations in Australia

Between 1788, when European colonists reported the first earthquake felt in Australia (Historical Records of Australia, 1914), and the early 1900s, newspaper articles were the main source of information about earthquakes (e.g., Malpas, 1991), and continued to be important until the 1960s (McCue, 2004). The first seismographs in Australia for




Figure 2.1 Map of the Australian continent plotting historical seismicity from the complete Australian catalogue for earthquakes $M \geq 4$. Zones of higher seismicity are shown: SWSZ, Southwest Seismic Zone; NWSSZ, Northwest Shelf Seismic Zone; FRSZ, Flinders Ranges Seismic Zone; SESZ, Southeast Seismic Zone. Significant earthquakes or earthquake sequences mentioned in text are indicated by numbers: 1, Meeberrie; 2, Tennant Creek; 3, offshore northeast Tasmania; 4, Dalton-Gunning; 5, Newcastle; 6, Beachport; 7, Warooka; 8, Adelaide; 9, Meckering; 10, Calingiri; 11, Cadoux; 12, Kalgoorlie-Boulder; 13, Ernabella; 14, Marryat Creek; 15, Katanning; 16, Norseman; 17, Ravensthorpe; 18, Mount Hotham. Numbers with asterisks indicate confirmed surface-rupturing earthquakes. The maximum horizontal stress ($S_{H_{max}}$) direction is indicated by fine dashed lines. The continent is broadly divided on the basis of geology and tectonic history, showing the dominance of non-extended cratonic crust in the central and western parts, non-extended non-cratonic crust in the east, and extended crust around much of the continental margin.




Figure 2.2 Indo-Australian Plate with plate boundary forces and orientation of modelled maximum and minimum horizontal stresses used in the finite element stress modelling of Reynolds *et al.* (2002). Much of the southern part of the continent has an east–west-oriented maximum horizontal compressive stress oriented at a high angle to the NNE-oriented plate velocity vector. Solid triangles indicate the direction of subduction and open triangles delineate the Banda Arc. TK, Tonga–Kermadec Trench; AAD, Australian–Antarctic discordance. (Modified after Reynolds *et al.*, 2002 and Hillis *et al.*, 2008.)

which records remain were four Milne seismographs established between 1901 and 1909 (Doyle and Underwood, 1965; McCue, 2004). The seismograph network remained sparse until 1962–3, after which time the Australian catalogue could be considered complete for magnitude (M) 5.0 and above earthquakes, and instrumental records largely replaced felt reports as the primary information source (McCue, 2004; Leonard, 2008). Doyle *et al.* (1968) summarised the history of earthquakes and seismological research in Australia up to 1966, and prepared a comprehensive reference list of historical earthquake sources. The authors tabulated over 160 Australian earthquakes, which formed the core of the preliminary digital Australian earthquake catalogue (Denham *et al.*, 1975). Despite enhancements resulting in the lowering of the completeness magnitude for the national catalogue to

M 3.5+ by the 1980s, the Australian National Seismic Network (ANSN – <http://www.ga.gov.au/earthquakes/seismicSearch.do>) remains relatively sparse, with no more than 70 seismic stations suitable for general earthquake monitoring in a continent of a similar size to the conterminous United States or Western Europe (Leonard, 2008). This results in generally poor constraints being placed on the location of the more than 36,000 onshore earthquakes recorded in the current Australian earthquake catalogue (Allen *et al.*, 2012a).

Approximately 20 earthquakes of $M > 6.0$ have been recorded in Australian continental crust, at the rate of one every ~ 6 –8 years (McCue, 1990; Leonard, 2008). On average, less than one $M > 5.0$ event has occurred per year (Leonard, 2008). The largest instrumentally recorded earthquake occurred in 1941 near Meeberrie, Western Australia (Everingham, 1982) (Figure 2.1 – 1). The event is associated with a surface wave magnitude of M_S 6.8 (Allen *et al.*, 2012a). In January 1988 three large “mainshock” earthquakes of magnitude M_S 6.3–6.7 (Choy and Bowman, 1990) occurred in a 12-hour period near Tennant Creek in the Northern Territory (Jones *et al.*, 1991) (Figure 2.1 – 2). The events were preceded by a year-long foreshock sequence, and followed by a continuing aftershock sequence (e.g., Bowman *et al.*, 1990). The most notable pre-instrumental earthquake sequence occurred in Bass Strait, off the northeast coast of Tasmania (Figure 2.1 – 3), in the 1880s–1890s. Over 2,400 events were felt during that period (Ripper, 1963), four of which caused damage in Tasmania, and two of which are estimated from felt effects to rival the Meeberrie earthquake in magnitude (Michael-Leiba, 1989). The Dalton–Gunning region of southeast Australia (Figure 2.1 – 4) continues to experience a less intense earthquake sequence (comprising more than 600 recorded events) that began with an M_L 5.6 event in the 1930s (Cleary, 1967; Denham *et al.*, 1981; Michael-Leiba *et al.*, 1994).

Earthquake clusters (e.g., Dalton–Gunning) are considered to be distinct from swarms, which are defined as occurring within a limited volume, lasting over a period from hours to months, with the largest event occurring well after the swarm commences, and not having a magnitude significantly greater than the second largest event (~ 0.5 magnitude unit; Gibson *et al.*, 1994; Dent, 2008). Geomechanical testing suggests that earthquake swarms occur preferentially in regions of extremely heterogeneous geological structure (Mogi, 1963). Increasing regional stress manifests as high stress concentration around numerous cracks and faults within the structured volume, resulting in failure on many local fractures at low stress. This has the effect of reducing the probability of failure on a single large fracture. Earthquake swarms are an important component of Australian seismicity, and can represent a large percentage of events in earthquake catalogues, particularly in regions of granitic geology (Dent, 2008). A non-exhaustive list of 42 earthquake swarms was compiled by Dent (2008, 2009), which includes perhaps the most significant of recent swarms, the 2001–2 Burakin Swarm (Leonard, 2002, 2003) in the Southwest Seismic Zone (SWSZ; Doyle, 1971) of Western Australia (Figure 2.1). This swarm involved six events in the magnitude range M_W 4.0–4.6 (Allen *et al.*, 2006) with over 18,000 smaller events, all purportedly occurring within a volume of ~ 5 km diameter (Leonard, 2003). The centre of activity for the swarm occurs approximately 20 km north of the surface rupture relating to the 1979 Cadoux M_S 6.0 earthquake (Figure 2.1 – 11). Several other swarms in the area (Kalannie, Beacon)

Table 2.1 Documented surface-rupturing earthquakes in Australia. Refer to Figure 2.1 for locations.

Year	Location	Magnitude (M_W)	Reference(s)
1968	Meckering, WA	6.6	Gordon and Lewis (1980), Vogtfjörd and Langston (1987), Johnston (1994)
1970	Calingiri, WA	5.5	Gordon and Lewis (1980), Johnston (1994)
1979	Cadoux, WA	6.1	Lewis <i>et al.</i> (1981), Fredrich <i>et al.</i> (1988)
1986	Marryat Creek, SA	5.8	Fredrich <i>et al.</i> (1988), Machette <i>et al.</i> (1993)
1988	Tennant Creek, NT*	6.3–6.6	Bowman <i>et al.</i> (1990), Chung <i>et al.</i> (1992), Crone <i>et al.</i> (1992)
2008	Katanning, WA	4.7	Dawson <i>et al.</i> (2008)
2012	Ernabella, SA	5.4	Clark <i>et al.</i> (2013)

* Tennant Creek involved a series of three consecutive events within one day.

suggest that the events might reflect continuing crustal adjustment relating to the Cadoux rupture (Dent, 2008, 2009). However, many other swarm sequences are apparently unrelated to larger events (Gibson *et al.*, 1994; Dent, 2008).

On 27 December 1989 an M_L 5.6 earthquake occurred near Newcastle, NSW (Figure 2.1 – 5), resulting in 13 fatalities and significant damage. The event was one of the most destructive and costly natural disasters to have occurred in Australia (McCue *et al.*, 1990; Sinadinovski *et al.*, 2002; Insurance Council of Australia, 2013), but not the first event to affect a populated centre. The 1897 Beachport, 1902 Warooka, and 1954 Adelaide events (McCue, 1975; Greenhalgh and Singh, 1988; Love, 1996) (Figure 2.1 – 6, 7, 8) caused damage in South Australia; the Warooka event also claimed two lives (McCue, 1975). The 1968 Meckering, 1970 Calingiri (Gordon and Lewis, 1980), 1979 Cadoux (Lewis *et al.*, 1981), and 2010 Kalgoorlie–Boulder (Bathgate *et al.*, 2010) (Figure 2.1 – 9, 10, 11, 12) events all caused damage in Western Australia, as did the 1988 Tennant Creek sequence in the Northern Territory (Bowman, 1992) (Figure 2.1 – 2).

Several of the damaging earthquakes noted previously also ruptured the ground surface. If the three 1988 Tennant Creek mainshocks (Bowman, 1988; Choy and Bowman, 1990; Bowman, 1992; Crone *et al.*, 1992) are treated as a single scarp-forming event, then the 23 March 2012 M_W 5.4 Ernabella earthquake (Clark *et al.*, 2013) (Figure 2.1 – 13) represents the seventh historic Australian event that can be unequivocally associated with surface rupture (Figure 2.1; Table 2.1). All seven of these events ruptured non-extended SCR cratonic crust (Figure 2.1) and involved a dominant reverse component to motion (Gordon and Lewis, 1980; Lewis *et al.*, 1981; Crone *et al.*, 1992; Machette *et al.*, 1993; Dawson *et al.*, 2008). As such, they are similar to the 1989 Ungava (Canada) (Adams *et al.*, 1991, 1992; Bent, 1994) and the 1993 Killari Latur (India) (Rajendran *et al.*, 1996, 2001; Seeber *et al.*, 1996; Rajendran, 2000) earthquakes.

In addition to the few documented surface-rupturing events, isoseismal maps have been compiled for almost 400 historic Australian earthquakes (Everingham, 1982; Rynn *et al.*, 1987; McCue, 1996). McCue (1980) derived a relationship between Richter magnitude (M) and the radius of a circle of equivalent area to the Modified Mercalli Intensity III isoseismal for Australian earthquakes with recorded magnitudes ranging from 3.6 to 7.0 (reproduced in McCue, 2004). This permitted magnitudes to be assigned for pre-instrumental earthquakes or recent earthquakes that were widely felt but either too close to or too far from the nearest seismograph to measure the magnitude (McCue, 2004).

2.2.1 Mechanism, geographic distribution, and strain rate

The Australian crustal stress regime is generally considered to be compressive (Denham *et al.*, 1981; Hillis and Reynolds, 2000, 2003). This assessment is supported by the majority of earthquake focal mechanisms, which range from thrust to oblique strike-slip (e.g., Leonard *et al.*, 2002; Keep *et al.*, 2012). Notable exceptions, with a dominant normal component, are the 1985 Norseman and 2001 Ravensthorpe earthquakes (Figure 2.1 – 16, 17), both located along the Albany–Fraser Orogen margin of the Yilgarn Craton, with the latter being anomalously deep at 18 km (McCue, 1989; Clark, 2004). The 1966 Mount Hotham earthquake in the SESZ (Denham *et al.*, 1982) (Figure 2.1 – 18), several of the Burakin Swarm events in the SWSZ (Leonard *et al.*, 2002), and two Tennant Creek aftershocks (Clark and Leonard, 2003) also have a dominant normal component to their focal mechanisms. Note that Leonard *et al.* (2002) incorrectly transposes the P and T axes, and dilatant and compressional quadrants, for the Norseman 1985 event (cf. McCue, 1989).

Assuming a maximum possible magnitude (M_{\max}) of ~ 7.0 (refer also to Section 2.5), the maximum seismogenic strain rate estimates for the continent (averaged over the last ~ 50 years of complete data) are $\sim 10^{-17}$ to 10^{-16} s^{-1} (Leonard, 2008; Braun *et al.*, 2009; Sandiford and Quigley, 2009). Comparable results (i.e., 10^{-17} s^{-1}) are obtained from thin-plate finite element modelling using plate boundary conditions, heat flow, stress, and geodetic data as inputs (Burbidge, 2004). This has been equated to an east–west shortening rate across southern Australia of approximately 0.3–0.4 mm/yr (Leonard, 2008), which compares to estimates based upon laser ranging and geodetic GPS of $0.65\text{--}3.0 \pm 2.0 \text{ mm/yr}$ (Smith and Kolenkiewicz, 1990; Tregonning, 2003; Leonard, 2008).

However, earthquake epicentres are not randomly distributed across the Australian continent in time or space (Denham, 1988; Leonard, 2008; Sinadinovski and McCue, 2010). At the continental scale, concentrations of epicentres occur in four major “seismogenic zones” (Figure 2.1) (Hillis *et al.*, 2008; Leonard, 2008; Sandiford and Egholm, 2008), with the continental margins also demonstrating a comparatively high number of epicentres relative to the interior (Sandiford and Egholm, 2008). Strain rates calculated for the higher seismicity zones are no more than 10^{-16} to 10^{-15} s^{-1} (Leonard, 2008; Braun *et al.*, 2009). Sandiford and Quigley (2009) provide an example from the Flinders Ranges Seismic Zone (FRSZ, Figure 2.1), where the bulk strain rate of 10^{-16} s^{-1} implies a total shortening of

~250 m/Ma across the ~100 km wide zone, which could be accommodated by 10 faults with slip rates consistent with the current topography and neotectonic faulting record (e.g., Sandiford, 2003b; Quigley *et al.*, 2006; Clark *et al.*, 2011a). Braun *et al.* (2009) estimate that half of the current relief of the Flinders Ranges, and a non-negligible proportion of the relief in the Eastern Highlands (SESZ, Figure 2.1), might plausibly have been built since the inception of the current stress regime. In contrast, similar seismogenic strain rates calculated for the SWSZ (Leonard, 2008; Braun *et al.*, 2009) (Figure 2.1) are not consistent with the present low-relief landscape (Clark, 2010), the distribution of paleo-earthquake fault scarps (Leonard, 2008; Leonard and Clark, 2011), or GPS strain rates (Leonard *et al.*, 2007). This implies migration of the locus of seismicity with time (Leonard, 2008; Leonard and Clark, 2011), an assertion supported by evidence suggesting that the rate of seismic activity in the SWSZ has increased dramatically in only the last 50 years (Michael-Leiba, 1987; Leonard and Clark, 2011).

2.2.2 Seismogenic depth

Depth constraint for Australian earthquakes is generally poor. Leonard (2008) selected a subset of the Australian earthquake catalogue for detailed hypocentral depth analysis using only depth values whose uncertainty was either less than the depth itself, or less than 5 km. More than 75% of epicentres from the current earthquake catalogue are rejected using these criteria. The selected data correlate with the regions of highest station density in private networks, and in the ANSN (i.e., southern Australia).

The concentration of epicentres in the SWSZ (Figure 2.1) occurs in non-extended cratonic crust. The Leonard (2008) dataset shows that earthquakes in this region are typically very shallow (see also Figure 2.3), with 95% of events found to have occurred within the upper 5 km of crust, consistent with previous analyses (Doyle, 1971; Everingham and Smith, 1979; Denham, 1988; McCue, 1990). This region has produced four surface-rupturing earthquakes in the last five decades (Figure 2.1; Table 2.1). The largest of these earthquakes (1968 Meckering; Figure 2.1 – 9) is calculated to have initiated at 1.5 km depth and to have ruptured both upwards to the surface and down to ~6 km depth (Langston, 1987; Fredrich *et al.*, 1988), while the smallest (2008 Katanning; Figure 2.1 – 15) ruptured from ~0.6 km depth to the surface (Dawson *et al.*, 2008). Earthquake swarms are also particularly prevalent in the SWSZ, with well-located depths being in the upper 2–3 km (Leonard, 2002, 2003). Analysis of the characteristics of modern and prehistoric fault scarps (Clark, 2010; Leonard and Clark, 2011; Clark *et al.*, 2012) suggests that large earthquake characteristics in the SWSZ (e.g., shallow depth) are typical of what might be expected throughout non-extended cratonic crust in Australia. However, Australia's deepest known earthquakes are also from this crustal setting: the 1989 Uluru earthquake (Figure 2.3 – 1), with a calculated depth of 31 km (Michael-Leiba *et al.*, 1994), and a 1992 earthquake beneath the Arafura Sea, north of the Northern Territory coastline (Figure 2.3 – 2), which had a depth of 39 km (McCue and Michael-Leiba, 1993).




Figure 2.3 Interpolated surface of variation in earthquake hypocentral depths across the Australian continent constructed using the ARCGIS Point Statistics tool (plotted value is the mean of values within a moving 3 degree search box, output cell size 0.25 degrees). Events with a catalogued depth of greater than 50 km are not considered. The earthquake catalogue used is from Allen *et al.* (2012a), and is not filtered according to hypocentral uncertainty (cf. Leonard, 2008). Higher seismicity zones are shown (cf. Figure 2.1). Locations referred to in text are marked. Significant earthquakes mentioned in text are indicated by numbers: 1, Uluru; 2, Banda Sea. Note the greater depth of events in the FRSZ and NWSSZ compared to the SWSZ and SESZ.

Despite being one of the most seismogenic regions of Australia (Leonard, 2008; Sandiford and Egholm, 2008), the Northwest Shelf Seismic Zone (NWSSZ – Figure 2.1) is characterized by extremely poorly constrained hypocentres, with depth uncertainties typically significantly larger than the depth values (e.g., Revets *et al.*, 2009; Allen *et al.*, 2012a). Consequently, this region was not analysed in detail by Leonard (2008). However, earthquake hypocentres tend to be deeper in this region of extended continental crust than in the adjoining non-extended cratonic crust (Figure 2.2). Approximately 40% of events recorded in the NWSSZ occur at depths greater than 10 km.

The FRSZ in South Australia consistently produces deeper earthquakes than other parts of the continent (Figure 2.3), with over 75% of well-constrained events occurring at depths greater than 10 km (Leonard, 2008). A temporary seismograph deployment (September to December 2003) suggested spatial variation in the depth distribution, with a predominance

of hypocentres in the 10–20 km depth range in the central Flinders Ranges, and less than 10 km depth in the southern Flinders Ranges (Cummins *et al.*, 2009), consistent with previous work (Greenhalgh *et al.*, 1994). The high proportion of hypocentres with depths greater than 10 km suggests that the depth to brittle–ductile transition in the Flinders Ranges is deeper than might be inferred from heat flow data (cf. Celerier *et al.*, 2005; Holford *et al.*, 2011).

Within the Southeast Seismic Zone (SESZ – Figure 2.3), and non-cratonic eastern Australia in general, hypocentral depth estimates are bimodal (Close and Seeber, 2007), ranging between very shallow (<5 km; 30% of data) and mid-crustal depths (10–20 km; 40% of data) (Allen *et al.*, 2012). Aftershocks tend to be very shallow and numerous, perhaps accounting for the shallower mode (Gibson *et al.*, 1981; Leonard, 2008). Within a sub-zone that encompasses the Pliocene to Holocene Newer Volcanic Province (e.g., Sutton *et al.*, 1977; Sheard, 1995), mid-crustal earthquakes are generally deeper, with 95% of recorded events occurring in the range 9–17 km (Leonard, 2008).

The Otway and Gippsland basins of southeast Australia (Figure 2.3) form part of an aulacogen developed in non-cratonic Paleozoic crust that was extended by rifting in the late Mesozoic–early Cenozoic. Earthquake hypocentres are deep in this region compared to those in the non-extended parts of the same Paleozoic basement province that is more typical of the SESZ. More than 70% of hypocentres in the extended basins are in the 10–25 km depth range (Allen *et al.*, 2012a), exemplified by the well-located 19 June 2012 M_L 5.4 Moe earthquake at a depth of 17 km (Sandiford and Gibson, 2012). It is probable that the contribution from these basins accounts for the greater depth element present in the southern southeast Australia (SEA-S) zone of Leonard (2008), as he does not distinguish between these two geological settings.

2.2.3 Attenuation and scaling relations

Several studies have shown that earthquakes in SCRs, such as Australia, are generally felt at larger distances than earthquakes in active tectonic regions (McCue, 1990; Frankel, 1994; Bakun and McGarr, 2002; Atkinson and Wald, 2007; Wald *et al.*, 2011). This is because the seismic energy propagates more efficiently through cold, relatively homogeneous continental crust, which is less susceptible to anelastic and scattering effects. It is recognised from analysis of isoseismal radii (Gaull *et al.*, 1990) that attenuation of seismic wave energy varies transversely across the Australian continent, with relatively low attenuation in the Archaean and Proterozoic terranes of western and central Australia and higher attenuation in the younger Phanerozoic terranes of eastern Australia (cf. Figure 2.1). Moreover, it is also observed that ground-motion amplitudes at large distances (>100 km) for an earthquake of given magnitude are lower in southeast Australia (SESZ) than in eastern North America (ENA) based on macroseismic intensities (Bakun and McGarr, 2002) and instrumental data (Allen and Atkinson, 2007). The higher attenuation observed in SESZ at distances greater than ~100 km is likely to be due to the broad crustal velocity

gradient (Collins *et al.*, 2003), which allows dispersion of Lg-wave energy into the upper mantle (e.g., Bowman and Kennett, 1991; Atkinson and Mereu, 1992). However, there appears to be no discernible difference between the attenuation of Australian and ENA ground motions in the distance range of engineering significance (i.e., <100 km; Allen and Atkinson, 2007; Allen, 2012). Furthermore, attenuation rates at these near-source distances are also comparable to those in active tectonic regions (e.g., Hanks and Johnston, 1992; Atkinson and Morrison, 2009; Campbell, 2011).

Rupture dimensions have been modelled for only two of Australia's large earthquake sequences – the 1968 Meckering earthquake and the 1988 Tennant Creek sequence (Somerville *et al.*, 2010) – both of which occurred in the cratonic western and central regions of Australia respectively (cf. Figure 2.1; Table 2.1). Significant simplifications were required to model the subsurface rupture geometries of both events, which involved complex surface rupture on intersecting structural elements of varying orientation (e.g., Crone *et al.*, 1992; Dentith *et al.*, 2009). While this might limit the confidence that can be placed in the rupture models, earthquake scaling relations developed from the models compare favourably with empirically derived relations for intraplate dip-slip earthquakes (Johnston, 1994), and self-similar scaling relations (Leonard, 2010). Following the 2012 Ernabella earthquake, updated relations between seismic moment and surface rupture length were developed, and suggest that earthquakes in nonextended cratonic Australia produce longer surface ruptures than is estimated from published scaling relations for earthquakes larger than $M_w \sim 6.5$ (Clark *et al.*, 2013). At present there are no estimates of the rupture dimensions of any earthquakes in the non-cratonic or extended crust regions of eastern Australia.

2.3 A long-term landscape record of large (morphogenic) earthquakes

Australia is one of the lowest, flattest, most arid and slowly eroding continents on Earth. Accordingly, large parts of Australia are favourable for the preservation of tectono-geomorphic features, such as fault scarps, for tens of thousands to millions of years (e.g., Quigley *et al.*, 2010). In regions with extremely low erosion rates, such as the Nullarbor Plain (Figure 2.4), it has been claimed that a morphogenic earthquake record spanning the last 15 Ma has been preserved essentially intact (Hillis *et al.*, 2008). On this basis, Australia boasts arguably the richest Late Neogene to Quaternary faulting record in all of the world's SCR crust (Sandiford, 2003b; Quigley *et al.*, 2010; Clark *et al.*, 2011a, 2012). Over 300 features (mainly fault scarps and folds) suspected or known to have been displaced under the current crustal stress regime have been identified and recorded (Clark *et al.*, 2011a, 2012) (Figure 2.4). The majority of these features, by virtue of their length and/or vertical displacement (see Section 3.1), are likely to reflect multiple surface-rupturing events (cf. Leonard, 2010). This remarkable archive, while undoubtedly incomplete, has the potential to extend the historic record of seismicity to a timescale commensurate with the recurrence time of large earthquakes in this intraplate setting.

Variation in fault scarp length, vertical displacement, proximity to other faults, and relationship to topography permits further division of this neotectonic record according




Figure 2.4 Neotectonic domains map of Australia (after Clark *et al.*, 2011a, 2012) showing features known or suspected of hosting displacement under the current stress field (grey lines). Neotectonic domains are indicated by numbers: 1, Precambrian Craton and Non-reactivated Proterozoic Crust; 2, Sprigg Orogen; 3, Reactivated Proterozoic Crust; 4, Eastern Australian Phanerozoic Accretionary Terranes; 5 Eastern Extended Continental Crust; 6, Western Extended Continental Crust; 7, Passive Margin Extended Continental and Transitional Crust. Shading and hatching as per Figure 2.1 shows the domains in their broader context of non-extended cratonic crust in the central and western parts, non-extended non-cratonic crust in the east, and extended margins around much of the continent.

to fault character. Six onshore “neotectonic domains” are recognised, with an additional offshore domain proposed by analogy with the eastern United States (Clark *et al.*, 2011a, 2012) (Figure 2.4). Each domain relates to a distinct underlying crustal type and architecture, broadly considered to represent non-extended cratonic and non-cratonic crust, as well as extended crustal settings (Figure 2.4) (cf. Johnston *et al.*, 1994). Herein domains will be referred to using the abbreviation Dx, where *x* is the number of the domain (e.g., D1 is Domain 1).

The Australian landscape reveals a marked disparity in its crustal response to imposed tectonic stresses. For example, neotectonic faults in western and central Australia, which are founded in cratonic crust (Figure 2.4 – D1, D3), are typically spaced more than a scarp length apart, not associated with historic seismicity, and displace a low, undulating landscape by less than 10 m (e.g., Clark, 2010). In contrast, faults in non-cratonic crust of the Mount Lofty and Flinders Ranges (Figure 2.4 – D2) are relatively closely spaced, commonly associated

with historic seismicity, and displace Neogene strata by up to a few hundred metres (e.g., Sandiford, 2003b; Quigley *et al.*, 2006). Further east, within non-cratonic Australia (Figure 2.4 – D4), faults are commonly found in *en echelon* arrangements, with fault complexes extending for several hundred kilometres along strike (Beavis, 1960, 1962; Moye *et al.*, 1963). As little as 15 m (e.g., Canavan, 1988; Robson and Webb, 2011; McPherson *et al.*, 2012b) and as much as several hundred metres of neotectonic displacement has been documented on several of these features (Beavis, 1960, 1962; Moye *et al.*, 1963; Abell, 1985), and there is no clear relationship to historic seismicity. Faults within the extended continental crust fringing the Australian continental margin (Figure 2.4 – D5, D6), a remnant of the Cretaceous breakup of the supercontinent Gondwana (e.g., Veevers, 2000), have some of the larger throws of any Australian neotectonic faults (greater than 100 m; Holdgate *et al.*, 2003). Extensional structural architecture is largely preserved (e.g., Hill *et al.*, 1994), and neotectonic faults are often spatially associated with earthquake epicentres.

2.3.1 Variation in fault scarp length and vertical displacement

The neotectonic data compiled by Clark *et al.* (2012) contain two semi-quantitative variables useful for characterising fault behaviour – length and vertical displacement. The population distributions for Australian fault scarp length and vertical displacement data are presented in Figure 2.5a, b. Fault length is defined as the along-strike distance (tail to tail) of discrete geomorphic features (most often fault scarps) that are considered to represent one or more surface-rupturing earthquake events. Vertical displacement is the vertical separation across a topographic scarp or fold. Many of the fault-length values reported by Clark *et al.* (2012) might be expected to be underestimates, as vertical displacement tapers towards the tails of ruptures, resulting in these scarp sections being less discoverable in digital elevation data: the primary tool for their identification and characterisation (Clark, 2010; Clark *et al.*, 2011a, 2012). It is also reasonable to assume that relatively short scarps are under-represented as they are less discoverable. This factor may explain the positive skew in both length and height data (Figure 2.5a, b). Furthermore, the resolution and noise content of digital elevation data from various sources might be expected to affect the precision of both length and vertical displacement measurements.

Interpolated surfaces for Australian neotectonic data demonstrate the spatial variation in these fault parameters (Figure 2.5c, d), and these spatial patterns are borne out in the statistics for each of the neotectonic domains (Figure 2.5e, f). The cratonic domains (D1, D3) are characterised by the lowest vertical displacement values (Figure 2.5d, f). In view of the extremely low erosion rates in these parts of Australia (Bierman and Caffee, 2002; Belton *et al.*, 2004; Chappell, 2006), this is indicative of very low time-averaged rates of morphogenic seismicity. Scarp lengths in D3 are up to 100 km greater than in D1 (Figure 2.5e), raising the possibility that relief in the Proterozoic mobile belts is built in fewer, larger earthquakes. Five fault scarps have been subject to detailed paleoseismological investigation in this crustal type: (1) Meckering (Clark *et al.*, 2011a), (2) Hyden (Crone




Figure 2.5 Population distributions for Australian neotectonic (a) fault length and (b) vertical displacement data, with accompanying plots showing the spatial distribution of (c) fault length and (d) vertical displacement data across the Australian continent. Interpolated surfaces in parts (a) and (b) are constructed using the same method as Figure 2.3. Grey lines indicate neotectonic domain boundaries (cf. Figure 2.4). Data values have been scaled for gridding – values in parentheses in the legend indicate the full range of underlying data. Box and whisker plots of (e) fault length and (f) vertical displacement data, binned by neotectonic domain. Boxes denote 75th and 25th percentiles, central point indicates median value, and whiskers define 90th and 10th percentiles. Data table shows minimum, maximum, median, and number of data points (n) for each domain.

et al., 2003; Clark *et al.*, 2008), (3) Dumbleyung and (4) Lort River (Estrada, 2009), and (5) Lake Edgar (Clark *et al.*, 2011b). These features range from 30 to 40 km in length (cf. Clark, 2010) with maximum single-event vertical displacements in the order of 1.2–3.1 m. Recurrence data for morphogenic earthquake events is restricted to the most recent one to three events on each fault, and indicates inter-event intervals of ~ 10 –40 ka (Figure 2.6).

Non-cratonic (D2, D4) and extended (D5, D6) domains are characterised by comparatively large vertical displacements (Figure 2.5d, f), which scaling relationships (e.g., Wells and Coppersmith, 1994; Leonard, 2010; Leonard and Clark, 2011) imply must have built as the result of multiple seismic cycles. There is, therefore, less certainty as to whether scarp lengths are representative of single-event ruptures, or are the product of segmented rupture. A positively skewed fault-length data distribution (Figure 2.5a) might plausibly reflect segmented rupture behaviour. The longest fault scarp in non-cratonic eastern Australia (D4) that has been subject to paleoseismological investigation is the Cadell Fault (Figure 2.4). Evidence from abandoned fluvial and tectonic landforms (e.g., Bowler and Harford, 1966; Rutherford and Kenyon, 2005) is consistent with seismic rupture of the entire 80 km scarp length, potentially involving 2–4 m of uplift (Clark *et al.*, 2011a; McPherson *et al.*, 2012a). While the timing of individual seismic events is poorly constrained, the average recurrence interval within the 70–20 ka active period (assuming full-length rupture) may have been as little as ~ 8 ka. Within D2, single-event displacement values of 1.8 m (1.5 m vertical) have been recorded on the Alma and Williamstown-Meadows faults (Clark and McPherson, 2011). The latter fault, which has a mapped length of over 100 km, is associated with a 25 km long single-event scarp.

2.3.2 The influence of crustal type and character on seismic activity rates

Given that the compressive nature of the Australian stress field (Hillis and Reynolds, 2003) results in a predominance of dip-slip faulting (Leonard *et al.*, 2002), long-term qualitative seismic activity rates across the continent might be assessed in terms of neotectonic uplift of the landscape. With respect to the morphogenic earthquake record, this is a function of neotectonic fault slip rate and density.

Long-term slip rates estimated for faults in the cratonic western part of the continent (D1, D3) are typically in the order of ~ 1 m/Ma (Clark *et al.*, 2008, 2012; Hillis *et al.*, 2008), which is equal to or less than the extant erosion rates (Bierman and Caffee, 2002; Belton *et al.*, 2004; Chappell, 2006). These long-term slip rates are consistent with the low-relief landscape, and contrast with uplift rates estimated from the last several decades of seismicity in the SWSZ, which have been suggested to be in the order of 10 m/Ma (Braun *et al.*, 2009).

Long-term slip rates in non-extended, non-cratonic eastern Australia (D4) are less well constrained. The slip rate on the Cadell Fault, averaged over the life of the current stress field (i.e., ~ 10 Ma), has been estimated at ~ 10 m/Ma (Clark *et al.*, 2007). New data from the Lake George Fault (Pillans, 2012) within the Eastern Highlands (Figure 2.4) indicate a




Figure 2.6 Compilation of surface-breaking earthquake recurrence data for SCR settings (Clark *et al.*, 2012, modified after Crone *et al.*, 2003). Lort River and Dumbleyung data from Estrada (2009), Wilkatana, Burra and Mundi Mundi data from Quigley *et al.* (2006), Lake Edgar data from Clark *et al.* (2011b) and Cadell data from Clark *et al.* (2007). Australian examples are labelled with their relevant neotectonic domain (cf. Figure 2.4).

slip rate of ~ 50 m/Ma. Higher relief areas of eastern Australia are associated with erosion rates of up to 30–50 m/Ma (Weissel and Seidl, 1998; Heimsath *et al.*, 2000, 2001; Wilkinson *et al.*, 2005; Tomkins *et al.*, 2007). Several authors have suggested that as much as 200 m of relief has been added to the Eastern Highlands over the last ~ 10 Ma (Sandiford, 2003b; Holdgate *et al.*, 2008; Braun *et al.*, 2009). Such estimates are consistent with uplift rates estimated from strain rates derived from contemporary seismicity (Braun *et al.*, 2009); however, little unequivocal evidence exists to assign this uplift to active faults (cf. Holdgate *et al.*, 2006). The approximate equivalence of erosion rates and fault slip rates suggests that much of the extant relief may be inherited (e.g., Bishop *et al.*, 1982; Pickett and Bishop, 1992; van der Beek *et al.*, 2001).

Perhaps the most spectacular examples of neotectonism in southeast Australia are the inverted Mesozoic Otway and Gippsland Basins (D5) (see Figure 2.1) (Holdgate *et al.*, 2003, 2007; Sandiford, 2003a), which formed by extension of non-cratonic (D4) crust. Within the Gippsland Basin in particular, the Cretaceous basin deeps now form the topographic highs at elevations of 200–300 m above sea level. Preliminary cosmogenic radionuclide ages obtained on overlying folded alluvial sediments (Holdgate *et al.*, 2007) suggest uplift rates of 60 m/Ma and greater on the major relief-forming faults (McPherson *et al.*, 2009; Clark *et al.*, 2011a, 2012). The relatively high fault density, combined with high slip rates and frequent contemporary seismicity (e.g., Leonard, 2008), identify these basins as among the most actively deforming parts of the continent (compare with the Flinders Ranges – D2).

Slip rates on faults underlying folds in the passive margin basins that dominate the Northwest Shelf region (D6) (NWSSZ, Figure 2.1), are poorly constrained. While locally having resulted in the uplift of Miocene marine deposits to elevations of over 100 m above sea level (e.g., van de Graaff *et al.*, 1976), neotectonic uplift is typically more modest than in the eastern aulacogen and passive margin basins (e.g., D5).

Long-term (i.e., averaged over several million years) vertical slip rates are known from some of the range-bounding faults of the Flinders and Mount Lofty Ranges (D2). These typically vary between ~ 20 and 50 m/Ma (Bourman *et al.*, 1999; Belperio *et al.*, 2002; Sandiford, 2003b; Celerier *et al.*, 2005; Quigley *et al.*, 2006). Bedrock erosion rates have been recorded at up to 122 m/Ma, but average around 40 m/Ma (Bierman and Caffee, 2002; Chappell, 2006; Quigley *et al.*, 2007a, b), allowing that relief is being produced within the ranges. Several authors suggest that up to half of the ~ 800 m relief has been built in the current stress regime (Sandiford, 2003b; Quigley *et al.*, 2007c; Braun *et al.*, 2009). Seismic reflection data indicate that the structural architecture mapped at the surface, corresponding to the Paleozoic Adelaide Fold Belt which developed over the inversion axis of a Neoproterozoic rift basin (Jenkins and Sandiford, 1992; Paul *et al.*, 1999), extends to depths of 10–12 km beneath the ranges (Flöttmann and Cockshell, 1996). However, much of the seismicity recorded in the region occurs below this depth (Leonard, 2008), potentially in cratonic crust relating to the eastern margin of the Gawler Craton (cf. Figure 2.1). Therefore, it is unclear how the surface-faulting record might be related, if at all, to most of the instrumental seismicity (cf. Braun *et al.*, 2009).

In general, greater topographic expression associated with faults and fault systems occurring in extended crust relative to non-extended crust suggests a higher rate of seismic activity in the extended setting, consistent with observations worldwide (e.g., Johnston, 1994; Cloetingh *et al.*, 2008; Mooney *et al.*, 2012). Using the same reasoning, non-cratonic crust might be expected to have a higher rate of seismic activity than cratonic crust (cf. Figures 2.4 and 2.5), by virtue of there being no relief generation in cratonic crust. This distinction, together with the variation in fault character between domains, should be recognised in attempts to identify analogous systems worldwide.

2.4 Patterns in earthquake occurrence

The record of contemporary seismicity in Australia suggests that earthquake epicentres are spatially and temporally clustered (Denham, 1988; Leonard, 2008; Sinadinovski and McCue, 2010). As discussed in Section 2.1, concentrations of epicentres in the historic catalogue are dominated by four “seismogenic zones” (Figure 2.1) (Hillis *et al.*, 2008; Leonard, 2008; Sandiford and Egholm, 2008). However, whether the short record of contemporary seismicity is representative of time periods significantly longer than the observation window has not been statistically or empirically tested in the Australian context (cf. Kafka, 2002). The persistence of patterns in the short historic catalogue can be assessed at much longer timescales by comparison with the record of morphogenic seismicity. Evidence from the paleo-record (Crone *et al.*, 2003; Clark *et al.*, 2011a, 2012), which essentially captures events of $\sim M > 5.5$, suggests that large earthquake occurrence within Australia exhibits both spatial and temporal clustering (Section 2.3). Temporal patterns in large SCR earthquake occurrence may be inferred at the scale of a single fault (Crone *et al.*, 1997, 2003; Clark *et al.*, 2008), of groups of faults (Leonard and Clark, 2011), and at the domain scale (Holdgate *et al.*, 2003; Sandiford, 2003b; Paine *et al.*, 2004; Braun *et al.*, 2009; Clark *et al.*, 2011a, 2012).

For example, the distribution of fault scarps in cratonic southwest Western Australia (D1) (Figure 2.4), together with the seismogenic strain arguments referred to previously (cf. Leonard *et al.*, 2007; Leonard, 2008; Braun *et al.*, 2009), infer that seismicity in the SWSZ represents only the current locus of activity, rather than a zone of long-lived activity (cf. Sandiford and Egholm, 2008). As most fault scarps in other parts of Australia are not associated with historic seismicity (e.g., Crone *et al.*, 2003; Clark, 2010), a similar rule may apply.

There is no precedent in cratonic Australia (neotectonic domains 1 and 3 of Clark *et al.*, 2012) to indicate how long seismicity will persist. However, a large proportion of contemporary seismicity in the SWSZ is thought to relate to the Calingiri, Cadoux, and Meckering earthquakes (Leonard, 2008). If aftershock activity relating to surface-rupturing earthquakes is used as a measure of the longevity of activity in a region, then the work of Stein and Liu (2009) suggests that a millennial timescale might be applicable. Liu *et al.* (2011) and Liu and Wang (2012) present evidence for migration of the locus of seismicity

at a centennial timescale in active intraplate northern China (Kusky *et al.*, 2007; Wheeler, 2011) with no recurrence over the 2,000–3,000-year window captured by the Chinese record of historical seismicity. As crustal deformation rates in this 1,000 km \times 1,000 km region of China are in the order of 1–2 mm/yr (Liu *et al.*, 2011), such data might be considered as a minimum seismicity migration rate for Australia at the neotectonic domain scale.

There is some indication that the temporal clustering behaviour emerging from single-fault studies in non-cratonic Australia may be symptomatic of a larger picture of the more-or-less continuous tectonic activity from the late Miocene to Recent being punctuated by “pulses” of activity in specific deforming regions (e.g., Quigley *et al.*, 2010; Clark *et al.*, 2012). For example, major deformation episodes are constrained to the interval 6–4 Ma in southwest Victoria (D4) (Paine *et al.*, 2004) and 2–1 Ma in the Otway Ranges (D5) (Sandiford, 2003a). An episode of deformation ceased at 1.0 Ma in the offshore Gippsland Basin (D5) although it continued onshore until \sim 250 ka (Holdgate *et al.*, 2003). Holdgate *et al.* (2008) present evidence from the southeast highlands that resurrects the idea of a punctuated post-Eocene Kosciuszko Uplift event (Browne, 1967) that continued into the late Pliocene and possibly the Pleistocene.

The Mount Lofty and Flinders Ranges (D2) are perhaps an exception to this rule. The FRSZ has been associated with diffuse earthquake activity throughout the historic era (Greenhalgh and Singh, 1988; Greenhalgh *et al.*, 1994). While epicentres cannot in most cases be confidently associated with neotectonic faults, strain rates and uplift rates estimated from the seismic catalogue are approximately consistent with the number of neotectonic faults and their paleoseismologically derived slip rates (e.g., Sandiford, 2003b; Braun *et al.*, 2009). Hence, it is possible that seismicity is a long-lived (millions of years) process in this crustal setting.

While the suite of neotectonic fault behaviours may vary across Australia, as implied by the neotectonic domains model, one individual fault characteristic appears to be common to most Australian intraplate faults studied – active periods comprising a finite number of events are separated by typically much longer periods of quiescence (Crone *et al.*, 1997, 2003; Clark *et al.*, 2007, 2008, 2011a, 2012; Estrada, 2009) (Figure 2.6). Data and modelling from elsewhere in the world identify similar episodic behaviour on faults with low slip rates (e.g., Wallace, 1987; Ritz *et al.*, 1995; Marco *et al.*, 1996; Friedrich *et al.*, 2003) and suggest that the time between successive clusters of events (deformation phases) is highly variable but significantly longer than the inter-event times between successive earthquakes within an active phase (Marco *et al.*, 1996; Stein *et al.*, 1997; Chéry *et al.*, 2001; Chéry and Vernant, 2006; Li *et al.*, 2009). This characteristic has been referred to as Wallace-type behaviour (see Wallace, 1987), and may be conceptualised using a model similar to that proposed by Friedrich *et al.* (2003) (Figure 2.7).

The sparse data available in Australia suggests that an active period (e.g., t_1 , t_2 , t_3 in Figure 2.7) in cratonic central or western Australia (D1) might comprise as few as two or three events (e.g., Hyden – Crone *et al.*, 2003; Clark *et al.*, 2008), with interseismic intervals between large events in an active period in the order of 20–40 ka (Crone *et al.*,




Figure 2.7 Generalised fault-slip diagram for Australian SCR faults based upon data from the Cadell Fault (Clark *et al.*, 2007; McPherson *et al.*, 2012a). Three active periods of fault growth (earthquake occurrence) are denoted by t_1 , t_2 , and t_3 . These active periods are relatively short-lived, and each is composed of only a few ruptures (*c.* <6 events per active period). Inter-event times between successive ruptures within an active period may range up to several thousands to several tens of thousands of years. We adopt a characteristic earthquake rupture model based upon paleoseismic data from the Cadell (Clark *et al.*, 2007; McPherson *et al.*, 2012a) and Lake Edgar (Clark *et al.*, 2011b) faults. Long quiescent periods separate the active periods, and the length of the quiescent periods can range from many tens of thousands of years to greater than a million years.

1997, 2003; Clark *et al.*, 2008; Estrada, 2009). In non-cratonic eastern Australia (D4), recently acquired data on the Cadell Fault (Clark *et al.*, 2007; McPherson *et al.*, 2012a) indicate more frequent rupture, with up to six morphogenic events in the interval *c.* 70–25 ka. It is inferred that three uplift events of similar magnitude had occurred on the Cadell Fault prior to the diversion of the Murray and Goulburn Rivers at *c.* 45 ka (Bowler and Harford, 1966; Bowler, 1978), suggesting that these events are likely to have been spaced thousands of years apart, similar to the Meers and Cheraw faults in the western Central United States (Crone and Luza, 1990; Crone and Machette, 1995). This contrasts with the New Madrid seismic zone in the intraplate Central United States, where sequences of large earthquakes have occurred on average every 500 years for at least the last two seismic cycles in the current active period (Tuttle *et al.*, 2002). While Talwani and Shaeffer (2001) propose a similar recurrence for earthquakes large enough to produce liquefaction in the South Carolina Coastal Plain, it has not been determined whether their data reflect rupture on a single fault or multiple faults. Quiescent intervals can be sufficiently prolonged (hundreds of thousands to millions of years), in the western and central parts of Australia in particular, that most or all relief relating to an active period might be removed by erosion prior to the next active period (Crone *et al.*, 2003; Clark *et al.*, 2007, 2008, 2011a).

2.5 Maximum magnitude earthquake

Large earthquakes are so infrequent in SCRs such as Australia that the data distributions upon which recurrence and M_{\max} estimates are based are heavily skewed towards

magnitudes below M_W 5.0 (e.g., Leonard, 2008), and thus require significant extrapolation up to magnitudes at which the most damaging ground-shaking might be expected (Wheeler, 2009a, b). Australia is uniquely suited to assessing the validity of M_{\max} estimates derived from the instrumental record of seismicity by virtue of its extraordinary neotectonic and paleo-earthquake record spanning many tens of thousands of years (e.g., Clark *et al.*, 2011a).

Where a fault has been studied paleoseismologically, single-event scarp lengths and single-event displacements can provide two independent estimates of the characteristic magnitude (cf. Schwartz and Coppersmith, 1984) for that fault, with scarp length being considered the more reliable measure (e.g., Wells and Coppersmith, 1994; Hemphill-Haley and Weldon, 1999; Leonard, 2010). Where a fault has not been studied in detail the “characteristic” rupture of the entire scarp length during each morphogenic event, subject to appropriate caveats regarding segmentation and dip, might be assumed in order to estimate future large earthquake potential (e.g., Hemphill-Haley and Weldon, 1999; Stirling *et al.*, 2002; Wheeler, 2009a).

In most regions of Australia the neotectonic record is either incomplete, or under-explored (Sandiford, 2003b; Clark *et al.*, 2011a). As such, it is not possible to assert with confidence that future large earthquakes will be restricted to known scarps/faults with documented source characteristics. It is therefore desirable to aggregate characteristic event magnitudes from a group of faults with similar source characteristics from within a region of interest (e.g., neotectonic domain or aggregation of domains [Leonard *et al.*, 2012; Clark *et al.*, 2012]) to provide an estimate of M_{\max} . This can be achieved with varying levels of confidence, depending upon the estimated completeness of the paleo-record.

2.5.1 Scarp length as a proxy for paleo-earthquake magnitude

A relatively large area of 10 m resolution DEM data in the southwest corner of the Yilgarn Craton (D1) (Figure 2.4) allowed for mapping of fault scarps in unprecedented detail (Clark, 2010). Based upon an assessment of erosion and landscape modification rates, it was estimated that most scarps representing events of $M_W \geq 6.5$ that had occurred in the last ~ 100 ka were captured in the catalogue (Leonard and Clark, 2011). The scaling relations of Leonard (2010) were used to develop a paleo-seismicity catalogue comprising 65 events (Leonard and Clark, 2006, 2011), which was subsequently combined with the catalogue derived from instrumental seismicity. The data were found to exhibit typical truncated Gutenberg–Richter recurrence characteristics with a slope (b) of 0.9–1.0 between M_W 6.5 and 6.9. A rapid roll off in recurrence occurred above M_W 6.9, with an asymptotic value of $M_W 7.25 \pm 0.1$ considered to be the M_{\max} for non-extended cratonic crust typified by the Yilgarn Craton (Leonard and Clark, 2006, 2011). A less well constrained M_{\max} of $M_W 7.65 \pm 0.1$ was determined for scarps representing extended crust in the rift basins flanking the western margin of the continent (D6).

In most regions of Australia the neotectonic catalogue is far from complete, the preservation time of seismogenic features in the landscape is uncertain, and single-event scarp

Table 2.2 *Ninetieth (90th) percentile fault scarp length values for each neotectonic domain of Clark *et al.* (2012) and corresponding characteristic earthquake magnitude (M_W) values determined using the scaling relation of Leonard (2010) assuming a 45° dipping fault and a seismogenic depth of 15 km*

Domain	Length [90%] (km)	Magnitude (M_W)
1	51	7.3*
2	78	7.4
3	104	7.6
4	124	7.6
5	57	7.3
6	124	7.6
All data	101	7.6

*The cratonic non-extended relation of Clark *et al.* (2013) obtains a value for D1 of M_W 6.9.

length and slip data are largely absent. Without these data, key assumptions of the curve-fitting approach to M_{\max} estimation described above are not satisfied. However, scarp length variation across the continent (Figure 2.5) does imply variation in characteristic earthquake magnitude (and M_{\max}).

Across Australia there is a strong positive skew in the length data distribution, with 90% of scarps less than 101 km in length (Figure 2.5a; Table 2.2). Assuming a generic fault dip of 45° and a seismogenic depth of 15 km (Collins *et al.*, 2003), the 90th percentile value for length corresponds to a value of M_W 7.6 (cf. Leonard, 2010) (Table 2.2). Without paleoseismic data on the longest scarps it is difficult to assess the validity of this estimate as a value for M_{\max} . The longest scarp with paleoseismological evidence consistent with entire length rupture is the 80 km long ($\sim M_W$ 7.4) Cadell Fault scarp in eastern Australia (D4) (Clark *et al.*, 2007, 2011a; McPherson *et al.*, 2012a); however, rupture segmentation is plausible for intraplate faults of several tens of kilometres in length or greater (e.g., Machette *et al.*, 1991). The modal scarp length value of 53 km for all neotectonic features (Figure 2.5a) corresponds to an earthquake of magnitude $\sim M_W$ 7.3.

Fault scarp length data for all individual domains (Figure 2.5e) show a similar positive skew to the aggregated population data distribution (Figure 2.5a), which highlights the same uncertainties with respect to possible fault segmentation. The 90% values for fault length provide magnitude estimates in the range of M_W 7.3–7.6 (Table 2.2). The values for D1 and D3 are within error of the M_{\max} estimated by Leonard and Clark (2011). Within D2 the only known single-event scarp is ~ 28 km long (M_W 6.8–6.9) and relates to the greater than 100 km long Williamstown-Meadows Fault (Clark and McPherson, 2011).

However, paleoseismological evidence exists elsewhere in D2 for very large single-event displacements (>7 m – Quigley *et al.*, 2006; Reid, 2007; Clark *et al.*, 2011a, 2012), consistent with modelled event magnitudes of M_W 7.3–7.5 (Somerville *et al.*, 2008). Very long 90th percentile values for D3 and D4 require validation in terms of equivalence to M_{\max} as these scarps also tend to have accumulated significant neotectonic throw (tens of metres), implying multiple morphogenic events. The dominance of folding as opposed to discrete faulting in D5 has thus far prevented estimation of single-event rupture lengths and displacements (Clark *et al.*, 2011a).

As expected, overall indications are that the historic catalogue of seismicity significantly underestimates the large earthquake potential (and, by proxy, M_{\max}) in most regions of Australia. To a first order at least, the sub-division of the continent into domains on the basis of geology and tectonic history (Johnston, 1994; Clark *et al.*, 2011a, 2012) provides useful insights into variations in faulting character that can facilitate the interpretation of the neotectonic and historic catalogues. The use of fault-length data from the neotectonic catalogue provides reasonable preliminary estimates of M_{\max} when applied in conjunction with appropriate scaling relations (e.g., Leonard, 2010; Clark *et al.*, 2013). Analysis of the current neotectonic catalogue for Australia suggests that a range of M_{\max} values of $M_W \sim 7.0$ – 7.6 could reasonably encompass all geological and tectonic settings continent-wide (Clark *et al.*, 2011a, 2012; Leonard, 2012; Burbidge, 2012).

2.6 Implications for SCR analogue studies: factors important in earthquake localisation

Analogues between the Australian neotectonic domains (Clark *et al.*, 2011a, 2012) and SCR crust elsewhere in the world (cf. Johnston, 1994) are readily apparent. For example, poly-phase deformation of a compressional nature is a common feature in the post-rift evolution of many extended passive margins and rifts (D5/D6 analogues) (van Arsdale, 2000; Balasubrahmanyam, 2006; Cloetingh *et al.*, 2008). Archean cratonic nuclei fringed by Paleoproterozoic mobile belts (D1 analogues) make up a large portion of the geology of Peninsular India (Kroner and Cordani, 2003) and North America (Hoffman, 1989). Meso- and Neoproterozoic mobile belts involved in the accretion of the supercontinent Rodinia (D3 analogues) are found worldwide (e.g., Collins and Pisarevsky, 2005; Cawood and Buchan, 2007), as are Phanerozoic accretionary terranes associated with the amalgamation of the supercontinent Gondwana (D4 analogues) (e.g., Hoffman, 1989). The temporal clustering of morphogenic earthquake events seen in studies on Australian faults (Quigley *et al.*, 2010; Clark *et al.*, 2011a, 2012) also appears to be mirrored in the Central and Eastern United States (Crone and Luza, 1990; Crone *et al.*, 1997; Cox *et al.*, 2006). Inferences made regarding mechanisms responsible for localising intraplate seismicity in Australia might then be assessed in terms of their crustal and lithospheric setting, and tested on analogous crust elsewhere in the world.

2.6.1 Mechanical and thermal influences

It has been proposed that intraplate regions with higher seismic potential have pre-existing zones of weakness (Sykes, 1978; Talwani and Rajendran, 1991; Stuart *et al.*, 1997; Kenner and Segall, 2000; Dentith and Featherstone, 2003), intersecting faults (Talwani, 1988, 1999), elevated heat flow (Liu and Zoback, 1997; Celerier *et al.*, 2005; Hillis *et al.*, 2008; Holford *et al.*, 2011), crustal anomalies (Campbell, 1978; Kenner and Segall, 2000; Gangopadhyay and Talwani, 2003; Pandey *et al.*, 2008; Assumpção and Sacek, 2013) or can be identified by crustal boundaries inferred from potential field data (Langenheim and Hildenbrand, 1997; Lamontagne *et al.*, 2003; van Lanen and Mooney, 2007; Dentith *et al.*, 2009). It is likely that the variety of models reflects the range of mechanisms that are operating, and although all of the above mechanisms have demonstrated local applicability in the Australian context, counter-examples are abundant. This is particularly the case where models relying on thermal mechanisms for strain localisation have been proposed.

Sandiford and Egholm (2008) argue that enhanced seismicity along some parts of the Australian continental margin is a consequence of thermal weakening due to steady-state heat flow across the lithospheric thickness steps between oceanic and continental crust (e.g., SESZ, Figure 2.1) and extended continental and non-extended cratonic crust (e.g., SWSZ, Figure 2.1) (Fishwick *et al.*, 2008; Kennett *et al.*, 2013). While the hypothesis is intuitively appealing, in Australia and along the eastern seaboard of North America (Wheeler and Frankel, 2000), several lines of evidence suggest that the contribution of this mechanism to the continental seismicity budget over geological timescales is minor.

For example, east of the large lithospheric thickness step on the western boundary of the SWSZ (i.e., across the Darling Fault), fault scarps are randomly distributed (Clark, 2010) and the topography predicted if instrumental seismic moment release rates are extrapolated to million-year timescales (Braun *et al.*, 2009) is absent. Given extremely low rates of bedrock erosion (Bierman and Caffee, 2002; Belton *et al.*, 2004; Chappell, 2006), this finding implies that the locus of seismicity in the SWSZ is transitory, rather than responding to steady-state heat flow at the margin. Furthermore, very little seismicity, paleo- or instrumental, can be correlated with the dramatic transition from cratonic to non-cratonic lithospheric thickness in eastern Australia (i.e., the Tasman Line, Figures 2.1 and 2.3), and significant instrumental seismicity proximal to the eastern seaboard (the SESZ) is not associated with a large lithospheric thickness step (Fishwick *et al.*, 2008).

More generally, Holford *et al.* (2011) propose that the thermal properties of the crust and upper mantle exert a regional-scale (100–1000 km) modulating control on which parts of the Australian lithosphere undergo (seismogenic) failure and which parts experience relatively less deformation (cf. Celerier *et al.*, 2005; Sandiford and Quigley, 2009). Specifically, these authors invoke relatively high heat flow to explain localisation of seismic moment release and deformation in the Flinders Ranges Seismic Zone (FRSZ, Figure 2.1) compared to the flanking Murray Basin and Nullarbor Plain. The correlation is imperfect; the heat flow anomaly does not extend to the Gippsland and Otway Basins (Densley *et al.*, 2000;

Holdgate *et al.*, 2003; Sandiford, 2003a) (Figure 2.1), which are manifestly amongst the fastest deforming regions on the Australian continent (Sandiford, 2003a, b; Clark *et al.*, 2011a, 2012). Actively inverting basins of the Northwest Shelf (NWSSZ, Figure 2.1) are also not associated with significant heat flow anomalies (cf. He and Middleton, 2002). Furthermore, the heat flow anomaly is most pronounced in the Cooper/Eromanga Basin, a region sparse in seismicity (Figure 2.1) and devoid of any known neotectonic features or tectonic uplift.

At the sub-regional scale, a range of factors have been proposed to explain localisation of historic seismicity. However, it seems clear that, on timescales of thousands to tens of thousands of years, seismic potential is determined by factors at a scale much larger than a single fault or region, and might instead relate more strongly to continental-scale lithospheric and crustal architecture, and the age of that architecture, as first proposed by Johnston *et al.* (1994).

2.6.2 Structural architectural influences

In the context of structural architecture, intraplate seismicity worldwide is considered to be concentrated at rifted margins (Stein *et al.*, 1989; Wheeler, 1995, 1996; Sandiford and Egholm, 2008; Cloetingh *et al.*, 2008; Etheridge *et al.*, 1991; Talwani and Schaeffer, 2001), interior rifts [aulacogens] (Johnston, 1994; Wheeler, 1995; Gangopadhyay and Talwani, 2003; Schulte and Mooney, 2005; Sinha and Mohanty, 2012) and at the margins of cratons (e.g., Lenardic *et al.*, 2000; Mazzotti, 2007; Sloan *et al.*, 2011; Craig *et al.*, 2011; Mooney *et al.*, 2012). A consequence of the relatively cold geotherm characterising cratonic areas is that there is no decoupling between crustal and mantle deformation (Braun *et al.*, 2009). This implies that, over geological timescales, deformation of the upper crust must be spatially uniform (e.g., Clark, 2010), resulting in little strain localisation, and hence minimal topography. Predictably, the historic record of seismicity in Australia is a poor guide as to where localisation of seismic activity occurs over geological timescales, with significant concentrations of seismicity occurring within cratons (e.g., SWSZ, Figure 2.1), and far from extended crust or craton boundaries (e.g., parts of the SESZ; see also Adams *et al.* [1992] and Rajendran *et al.* [1996]).

The major tenet of the Johnston *et al.* (1994) SCR model – that extended crust is more seismically active than non-extended crust, and that within the non-extended class non-cratonic crust is more active than cratonic crust – appears to hold true for the Australian neotectonic record if the uplift rate implied by vertical neotectonic fault displacement is taken as a proxy for seismic activity (Figure 2.4d, f). Australian neotectonic data (Clark *et al.*, 2011a, 2012) permit further sub-division of the extended and non-extended crustal classes proposed by Johnston *et al.* (1994). In extended crust domains (e.g., D5, D6), the age of major rifting appears to be important in terms of the record of neotectonic activity. Paleozoic intracratonic rifts (e.g., the Fitzroy Trough [Drummond *et al.*, 1991]; cf. Figure 2.1) and passive margin components (e.g., Perth Basin [Crostella and Backhouse,

2000; Norvick, 2004]) preserve less evidence for neotectonic strain localisation than those rifted in the Mesozoic (e.g., Gippsland and Otway basins). As such, the long-term seismic potential of aulacogens impinging along the eastern margin of the North American continent (e.g., Ottawa Rift, Saguenay graben, Reelfoot Rift, Southern Oklahoma aulacogen; Wheeler and Frankel [2000]) might be assessed based upon the proportion of Mesozoic extensional reactivation of these largely Paleozoic structures. The distinction becomes even clearer when one considers most Precambrian rifts (included in D1 and D3). The major exception to this rule is the FRSZ (D2), which formed as a Precambrian rift, but was subsequently extensively reactivated under compression in the Paleozoic (Flöttmann and James, 1997). The Appalachian Orogen (Wheeler, 1996), including the St. Lawrence seismic zone (Vlahovic *et al.*, 2003) at its northern margin, occupies a similar crustal setting between Precambrian mobile belts (D3 equivalent) and Phanerozoic accretionary terranes (D4 equivalent).

Considerable uncertainty remains as to the long-term seismic potential of non-extended, non-cratonic crust (D4), which typically comprises Phanerozoic accretionary terranes (e.g., Hoffman, 1989; Glen, 2005). Within the Australian continent, much of this crustal type is associated with little positive relief, and so might be thought of as having low seismic potential (e.g., Sandiford, 2003b). However, the eastern part of D4 is punctuated by the Eastern Highlands. While it is widely considered that most of the relief of the highlands relates to the opening of the Tasman Sea in the Cretaceous (e.g., Norvick and Smith, 2001), debate remains as to how much relief has been added from the Neogene to Recent (Holdgate *et al.*, 2008, 2011; VandenBerg, 2010). The few known faults are poorly documented, but appear to have similar characteristics to those documented from the relief-poor western parts of the domain (compare the Lake George Fault [Abell, 1981, 1985] to the Cadell Fault [Clark *et al.*, 2007; McPherson *et al.*, 2012a]). Braun *et al.* (2009) hypothesise that deformation in eastern Australia is determined by the lithospheric strength (or rigidity) and hence controlled by lithospheric structures, in contrast to the western part of the continent where deformation is localised by crustal structures, because the underlying lithospheric mantle is almost ubiquitously strong.

A fundamental implication of the neotectonic domains model presented by Clark *et al.* (2011a, 2012), building upon the pioneering work of Johnston *et al.* (1994), is that intraplate fault characteristics (and by inference, seismicity) are not universal in their applicability in analogue studies. Careful choice of subject regions, structures, or faults within analogous crust of similar stress field character is required to extrapolate meaningfully to imperfectly characterised areas.

2.7 Conclusions

Australia has a short recorded history of seismicity, spanning only a couple of centuries (Leonard, 2008). As a consequence, there is significant uncertainty as to whether patterns evident in the contemporary seismic record are representative of the longer term. This uncertainty can, in part, be overcome by validation against Australia's rich record of

morphogenic earthquakes (Quigley *et al.*, 2010; Clark *et al.*, 2012). Long-term patterns in large earthquake occurrence, both temporal and spatial, can be deduced from the landscape record and used to inform contemporary earthquake hazard science. Seismicity source parameters such as large earthquake recurrence and magnitude vary across the Australian continent, and can be interpreted in a framework of large-scale neotectonic domains defined on the basis of geology and crustal setting (Clark *et al.*, 2012). Temporal and spatial clustering of earthquakes is apparent at the scale of a single fault, and at the 1,000 km scale of a domain. The utility of the domains approach, which ties seismicity characteristics to crustal architecture and geology, is that behaviours might be extrapolated from well-characterised regions to poorly known analogous regions, both within Australia and worldwide.

Acknowledgements

This paper is published with the permission of the CEO of Geoscience Australia. The authors would like to thank Marcelo Assumpção and Pradeep Talwani for providing review comments that improved the manuscript. Mark Leonard and David Burbidge are also thanked for their constructive reviews.

References

- Abell, R. S. (1981). Notes to accompany the 1:25 000 scale geological field compilation sheets of the CANBERRA 1:100 000 sheet area. *Bureau of Mineral Resources, Geology and Geophysics Record* 1981/6.
- Abell, R. S. (1985). Geology of the Lake George Basin, NSW *Bureau of Mineral Resources, Geology and Geophysics Record* 1985/4.
- Adams, J., R. Wetmiller, H. Hasegawa, and J. Drysdale (1991). The first surface faulting from a historical earthquake in North America. *Nature*, 352, 617–619.
- Adams, J., J. Percival, R. Wetmiller, J. Drysdale, and P. Robertson (1992). Geological controls on the 1989 Ungava surface rupture: a preliminary interpretation. *Geological Survey of Canada Paper*, 92C, 147–155.
- Allen, T. I. (2012). Ground motion prediction equations. In *The 2012 Australian Earthquake Hazard Map*, ed. D. R. Burbidge. *Geoscience Australia Record*, 2012/71, pp. 44–53.
- Allen, T. I., and G. M. Atkinson (2007). Comparison of earthquake source spectra and attenuation in eastern North America and southeastern Australia. *Bulletin of the Seismological Society of America*, 97, 1350–1354.
- Allen, T. I., M. Leonard, and C. D. N. Collins (2012). Revised Australian Earthquake Catalogue. In *The 2012 Australian Earthquake Hazard Map*, ed. D. R. Burbidge. *Geoscience Australia Record*, 2012/71, pp. 5–16.
- Assumpção, M., and V. Sacek (2013). Intra-plate seismicity and flexural stresses in central Brazil. *Geophysical Research Letters*, 40, 1–5.
- Atkinson, G. M., and R. F. Mereu (1992). The shape of ground motion attenuation curves in southeastern Canada. *Bulletin of the Seismological Society of America*, 82, 2014–2031.

- Atkinson, G. M., and M. Morrison (2009). Observations on regional variability in ground-motion amplitudes for small-to-moderate earthquakes in North America. *Bulletin of the Seismological Society of America*, 99, 2393–2409.
- Atkinson, G. M., and D. J. Wald (2007). “Did You Feel It?” intensity data: a surprisingly good measure of earthquake ground motion. *Seismological Research Letters*, 78, 362–368.
- Bakun, W. H., and A. McGarr (2002). Differences in attenuation among the stable continental regions. *Geophysical Research Letters*, 29, 2121, doi:10.1029/2002GL015457.
- Balasubrahmanyam, M. N. (2006). *Geology and Tectonics of India: An Overview*. International Association for Gondwana Research, IAGR Memoir 9.
- Bathgate, J., H. Glanville, and C. Collins (2010). The Kalgoorlie earthquake of 20 April 2010 and its aftershock sequence. *Australian Earthquake Engineering Society 2010 Conference*, Perth, Paper 54.
- Beavis, F. C. (1960). The Tawonga Fault, northeast Victoria. *Proceedings of the Royal Society of Victoria*, 72, 95–100.
- Beavis, F. C. (1962). The geology of the Kiewa area. *Proceedings of the Royal Society of Victoria*, 75, 349–410.
- Belperio, A. P., N. Harvey, and R. P. Bourman (2002). Spatial and temporal variability in the Holocene sea-level record of the South Australian coastline. *Sedimentary Geology*, 150, 153–169.
- Belton, D. X., R. W. Brown, B. P. Kohn, D. Fink, and K. A. Farley (2004). Quantitative resolution of the debate over antiquity of the central Australian landscape: implications for the tectonic and geomorphic stability of cratonic interiors. *Earth and Planetary Science Letters*, 219, 21–34.
- Bent, A. L. (1994). The 1989 (Ms 6.3) Ungava, Quebec, earthquake: a complex intraplate event. *Bulletin of the Seismological Society of America*, 84, 1075–1088.
- Bierman, P. R., and M. Caffee (2002). Cosmogenic exposure and erosion history of Australian bedrock landforms. *Geological Society of America Bulletin*, 114, 787–803.
- Bishop, P., P. Hunt, and P. W. Schmidt (1982). Limits to the age of the Lapstone Monocline, N.S.W.; a palaeomagnetic study. *Journal of the Geological Society of Australia*, 29, 319–326.
- Bourman, R. P., A. P. Belperio, C. V. Murray-Wallace, and J. H. Cann (1999). A last interglacial embayment fill at Normanville, South Australia, and its neotectonic implications. *Transactions of the Royal Society of South Australia*, 123, 1–15.
- Bowler, J. M. (1978). Quaternary climate and tectonics in the evolution of the Riverine Plain, southeastern Australia. In *Landform Evolution in Australasia*, ed. J. L. Davies, and M. A. J. Williams. Canberra: ANU Press, pp. 70–112.
- Bowler, J. M., and L. B. Harford (1966). Quaternary tectonics and the evolution of the Riverine Plain near Echuca, Victoria. *Journal of the Geological Society of Australia*, 13, 339–354.
- Bowman, J. R. (1988). Constraints on locations of large intraplate earthquakes in the Northern Territory, Australia, from observations at the Warramunga seismic array. *Geophysical Research Letters*, 15, 1475–1478.
- Bowman, J. R. (1992). The 1988 Tennant Creek, Northern Territory, earthquakes: a synthesis. *Australian Journal of Earth Sciences*, 39, 651–669.
- Bowman, J. R., and B. L. N. Kennett (1991). Propagation of *Lg* waves in the north Australian craton: influence of crustal velocity gradients. *Bulletin of the Seismological Society of America*, 81, 592–610.

- Bowman, J. R., G. Gibson, and T. Jones (1990). Aftershocks of the 1988 January 22 Tennant Creek, Australia intraplate earthquakes: evidence for a complex thrust-fault geometry. *International Geophysical Journal*, 100, 87–97.
- Braun, J., D. Burbidge, F. Gesto, *et al.* (2009). Constraints on the current rate of deformation and surface uplift of the Australian continent from a new seismic database and low-T thermochronological data. *Australian Journal of Earth Sciences*, 56, 99–110.
- Browne, W. R. (1967). Geomorphology of the Kosciusko block and its north and south extensions. *Proceedings of the Linnean Society of New South Wales*, 92, 117–144.
- Burbidge, D. R. (2004). Thin plate neotectonic models of the Australian plate. *Journal of Geophysical Research*, 109, B10405, doi:10.1029/2004JB003156.
- Burbidge, D. R. (editor) (2012). *The 2012 Australian Earthquake Hazard Map*, Geoscience Australia, Record 2012/71.
- Campbell, D. L. (1978). Investigation of the stress-concentration mechanism for intraplate earthquakes. *Geophysical Research Letters*, 5, 477–479.
- Campbell, K. W. (2011). Ground motion simulation using the hybrid empirical method: issues and insights. *Geotechnical Geological and Earthquake Engineering*, 14, 81–95.
- Canavan, F. (1988). Deep lead gold deposits of Victoria. *Geological Society of Victoria Bulletin*, 62.
- Cawood, P. A., and C. Buchan (2007). Linking accretionary orogenesis with supercontinent assembly. *Earth Science Reviews*, 82, 217–256.
- Celerier, J., M. Sandiford, D. L. Hansen, and M. Quigley (2005). Modes of active intraplate deformation, Flinders Ranges, Australia. *Tectonics*, 24, doi:10.1029/2004TC001679.
- Chappell, J. (2006). Australian landscape processes measured with cosmogenic nuclides. In *Regolith Geochronology and Landscape Evolution*, ed. B. Pillans. Perth: CRC LEME, pp. 19–26.
- Chéry, J., S. Carretier, and J. F. Ritz (2001). Postseismic stress transfer explains time clustering of large earthquakes in Mongolia. *Earth and Planetary Science Letters*, 194, 277–286.
- Chéry, J., and P. Vernant (2006). Lithospheric elasticity promotes episodic fault activity. *Earth and Planetary Science Letters*, 243, 211–217.
- Choy, G. L., and J. R. Bowman (1990). Rupture process of a multiple main shock sequence; analysis of teleseismic, local, and field observations of the Tennant Creek, Australia, earthquakes of January 22, 1988. *Journal of Geophysical Research B, Solid Earth and Planets*, 95, 6867–6882.
- Chung, W.-Y., A. C. Johnston, and J. Pujol (1992). *A Global Study of Stable Continental Intraplate Earthquakes: Focal Mechanisms, Source-Scaling Relations, and Seismotectonics*. Final Report, EPRI RP-2556–54, Electric Power Research Institute, Palo Alto, California.
- Clark, D. (2004). Seismicity along the northern foreland of the Albany-Fraser Orogen, Geoscience Australia (Earthquake Hazard Section) unpublished report.
- Clark, D. (2010). Identification of Quaternary scarps in southwest and central west Western Australia using DEM-based hill shading: application to seismic hazard assessment and neotectonics. *International Journal of Remote Sensing*, 31, 6297–6325.
- Clark, D. J., and M. Leonard (2003). Principal stress orientations from multiple focal plane solutions: new insight in to the Australian intraplate stress field. In *Evolution and Dynamics of the Australian Plate*, ed. R. R. Hillis, and D. Muller. Geological Society

- of Australia Special Publication 22 and Geological Society of America Special Paper 372, 91–105.
- Clark, D., and A. McPherson (2011). Large earthquake recurrence in the Adelaide region: a palaeoseismological perspective. *Australian Earthquake Engineering Society 2011 Conference*, 18–20 November, Barossa Valley, South Australia, Paper 11.
- Clark, D., M. Dentith, and M. Leonard (2003). Linking earthquakes to geology: contemporary deformation controlled by ancient structure. *Geological Society of Australia, Abstracts*, 72, 90.
- Clark, D., R. Van Dissen, M. Cupper, C. Collins, and A. Prendergast (2007). Temporal clustering of surface ruptures on stable continental region faults: a case study from the Cadell Fault scarp, south eastern Australia. *Australian Earthquake Engineering Society Conference*, Wollongong, Australian Earthquake Engineering Society.
- Clark, D., M. Dentith, K. H. Wyrwoll, *et al.* (2008). The Hyden fault scarp, Western Australia: paleoseismic evidence for repeated Quaternary displacement in an intracratonic setting. *Australian Journal of Earth Sciences*, 55, 379–395.
- Clark, D., A. McPherson, and C. D. N. Collins (2011a). *Australia's Seismogenic Neotectonic Record: A Case for Heterogeneous Intraplate Deformation*, *Geoscience Australia Record*, 2011/11.
- Clark, D. J., M. Cupper, M. Sandiford, and K. Kiernan (2011b). Style and timing of late Quaternary faulting on the Lake Edgar Fault, southwest Tasmania, Australia: implications for hazard assessment in intracratonic areas. In *Geological Criteria for Evaluating Seismicity Revisited: Forty Years of Paleoseismic Investigations and the Natural Record of Past Earthquakes*, ed. F. A. Audemard, M. A. Michetti, and J. P. McCalpin. Geological Society of America Special Publication 479, pp. 109–131.
- Clark, D., A. McPherson, and R. Van Dissen (2012). Long-term behaviour of Australian Stable Continental Region (SCR) faults. *Tectonophysics*, 566–567, 1–30.
- Clark, D., A. McPherson, T. Allen, and M. De Kool (2013). Co-seismic surface deformation relating to the March 23, 2012 MW 5.4 Ernabella (Pukatja) earthquake, central Australia: implications for cratonic fault scaling relations. *Bulletin of the Seismological Society of America*, 104, doi:10.1785/0120120361.
- Cleary, J. R. (1967). The seismicity of the Gunning and surrounding areas, 1958–1961. *Journal of the Geological Society of Australia*, 14, 23–29.
- Cloetingh, S., F. Beekman, P. A. Ziegler, J.-D. Van Wees, and D. Sokoutis (2008). Post-rift compressional reactivation potential of passive margins and extensional basins. In *The Nature and Origin of Compression in Passive Margins*, ed. H. Johnson, A. G. Doré, R. W. Gatliff *et al.* Geological Society, London, Special Publication 306, pp. 27–70.
- Cloetingh, S., and R. Wortel (1986). Stress in the Indo-Australian plate. *Tectonophysics*, 132, 49–67.
- Close, D., and L. Seeber (2007). Shallow seismicity in stable continental regions. *Seismological Research Letters*, 78, 554–562.
- Coblentz, D. D., M. Sandiford, R. M. Richardson, Z. Shaohua, and R. Hillis (1995). The origins of the intraplate stress field in continental Australia. *Earth and Planetary Science Letters*, 133, 299–309.
- Coblentz, D. D., S. Zhou, R. R. Hillis, R. M. Richardson, and M. Sandiford (1998). Topography boundary forces and the Indo-Australian intraplate stress field. *Journal of Geophysical Research*, 103, 919–938.

- Collins, A. S., and S. A. Pisarevsky (2005). Amalgamating eastern Gondwana: the evolution of the Circum-Indian Orogens. *Earth Science Reviews*, 71, 229–270.
- Collins, C. D. N., B. J. Drummond, and M. G. Nicoll (2003). Crustal thickness patterns in the Australian continent. In *Evolution and Dynamics of the Australian Plate*, ed. R. R. Hillis, and R. D. Müller. Geological Society of Australia Special Publication 22 and Geological Society of America Special Paper 372, pp. 121–128.
- Cox, R. T., J. Cherryhomes, J. B. Harris, and D. Larsen (2006). Paleoseismology of the southeastern Reelfoot rift in western Tennessee and implications for intraplate fault zone evolution. *Tectonics*, 25, 1–17.
- Craig, T. J., J. A. Jackson, K. Priestly, and D. McKenzie (2011). Earthquake distribution patterns in Africa: their relationship to variations in lithospheric and geologic structure, and their rheological implications. *Geophysical Journal International*, 185, 403–434.
- Crone, A. J., and K. V. Luza (1990). Style and timing of Holocene surface faulting on the Meers Fault, southwestern Oklahoma. *Geological Society of America Bulletin*, 102(1), 1–17.
- Crone, A. J., and M. N. Machette (1995). Holocene faulting on the Cheraw Fault, southeastern Colorado: another hazardous late Quaternary fault in the stable continental interior. *Eos Supplement*, 76, F362.
- Crone, A. J., M. N. Machette, and J. R. Bowman (1992). Geologic investigations of the 1988 Tennant Creek, Australia, earthquakes: implications for paleoseismicity in stable continental regions. *Bulletin U.S. Geological Survey*, 2032-A.
- Crone, A. J., M. N. Machette, and J. R. Bowman (1997). Episodic nature of earthquake activity in stable continental regions revealed by palaeoseismicity studies of Australian and North American Quaternary faults. *Australian Journal of Earth Sciences*, 44, 203–214.
- Crone, A. J., P. M. de Martini, M. N. Machette, K. Okumura, and J. R. Prescott (2003). Paleoseismicity of two historically quiescent faults in Australia: implications for fault behavior in stable continental regions. *Bulletin of the Seismological Society of America*, 93, 1913–1934.
- Crostella, A., and J. Backhouse (2000). Geology and petroleum exploration of the central and southern Perth Basin, Western Australia. *Western Australia Geological Survey, Report*, 57, 85.
- Cummins, P., C. Collins, and A. Bullock (2009). Monitoring of earthquakes in the Flinders Ranges, South Australia, using a temporary seismometer deployment. Geoscience Australia Publication (digital), https://www.ga.gov.au/products/servlet/controller?event=GEOCAT_DETAILS&catno=69525.
- Dawson, J., P. Cummins, P. Tregoning, and M. Leonard (2008). Shallow intraplate earthquakes in Western Australia observed by Interferometric Synthetic Aperture Radar. *Journal of Geophysical Research*, 113, doi:10.1029/2008JB005807.
- Denham, D. (1988). Australian seismicity: the puzzle of the not so stable continent. *Seismological Research Letters*, 59, 235–240.
- Denham, D., G. R. Small, J. R. Cleary, *et al.* (1975). Australian earthquakes (1897–1972), *Search*, 6, 34–36.
- Denham, D., J. Weekes, and C. Krayshok (1981). Earthquake evidence for compressive stress in the southeast Australian crust. *Journal of the Geological Society of Australia*, 28, 323–332.

- Denham, D., G. Gibson, R. S. Smith, and R. Underwood (1982). Source mechanisms and strong ground motion from the 1982 Wonnangatta and the 1966 Mount Hotham earthquakes. *Australian Journal of Earth Sciences*, 32, 37–46.
- Densley, M. R., R. R. Hillis, and J. E. P. Redfearn (2000). Quantification of uplift in the Carnarvon Basin based on interval velocities. *Australian Journal of Earth Sciences*, 47, 111–122.
- Dent, V. F. (2008). Graphical representation of some recent Australian earthquake swarms, and some generalisations on swarm characteristics. *Australian Earthquake Engineering Society Conference Proceedings*, Ballarat, Paper 23.
- Dent, V. F. (2009). The Beacon, WA earthquake swarm commencing January 2009. *Australian Earthquake Engineering Society Conference*, Newcastle, poster presentation.
- Dentith, M. C., and W. E. Featherstone (2003). Controls on intra-plate seismicity in south-western Australia. *Tectonophysics*, 376, 167–184.
- Dentith, M. C., D. J. Clark, and W. E. Featherstone (2009). Aeromagnetic mapping of Precambrian geological structures that controlled the 1968 Meckering Earthquake (Ms 6.8): implications for intraplate seismicity in Western Australia. *Tectonophysics*, 475, 544–553.
- Dickinson, J. A., M. W. Wallace, G. R. Holdgate, S. J. Gallagher, and L. Thomas (2002). Origin and timing of the Miocene-Pliocene unconformity in southeast Australia. *Journal of Sedimentary Research*, 72, 288–303.
- Doyle, H. A. (1971). Seismicity and structure in Australia. *Bulletin of the Royal Society of New Zealand*, 9, 149–152.
- Doyle, H. A., and R. Underwood (1965). Seismological stations in Australia. *Australian Journal of Earth Sciences*, 28, 40–43.
- Doyle, H. A., I. B. Everingham, and D. J. Sutton (1968). Seismicity of the Australian continent. *Journal of the Geological Society of Australia*, 15, 295–312.
- Drummond, B. J., M. J. Sexton, T. J. Barton, and R. D. Shaw (1991). The nature of faulting along the margins of the Fitzroy Trough, Canning Basin, and implications for the tectonic development of the Trough. *Exploration Geophysics*, 22, 111–115.
- Dyksterhuis, S., and R. D. Müller (2008). Cause and evolution of intraplate orogeny in Australia. *Geology*, 36, 495–498.
- Estrada, B. (2009). Neotectonic and palaeoseismological studies in the southwest of Western Australia. Ph.D. thesis (unpublished), School of Earth and Environment, The University of Western Australia, Perth.
- Etheridge, M. A., H. W. S. McQueen, and K. Lambeck (1991). The role of intraplate stress in Tertiary (and Mesozoic) deformation of the Australian continent and its margins: a key factor in petroleum trap formation. *Exploration Geophysics*, 22, 123–128.
- Everingham, I. B. (1982). Atlas of isoseismal maps of Australian earthquakes. *Bureau of Mineral Resources Bulletin*, 214.
- Everingham, I. B., and R. S. Smith (1979). Implications of fault-plane solutions for Australian earthquakes on 4 July 1977, 6 May 1978 and 25 November 1978. *BMR Journal of Australian Geology and Geophysics*, 4, 297–301.
- Fishwick, S., M. Heintz, B. L. N. Kennett, A. M. Reading, and K. Yoshizawa (2008). Steps in lithospheric thickness within eastern Australia, evidence from surface wave tomography. *Tectonics*, 27, doi:10.1029/2007TC002116.
- Flöttmann, T., and C. D. Cockshell (1996). Palaeozoic basins of southern South Australia: new insights into their structural history from regional seismic data. *Australian Journal of Earth Sciences*, 43, 45–55.

- Flöttmann, T., and P. James (1997). Influence of basin architecture on the style of inversion and fold-thrust belt tectonics: the southern Adelaide Fold-Thrust Belt, South Australia. *Journal of Structural Geology*, 19, 1093–1110.
- Frankel, A. (1994). Implications of felt area–magnitude relations for earthquake scaling and the average frequency of perceptible ground motion. *Bulletin of the Seismological Society of America*, 84, 462–465.
- Fredrich, J., R. McCaffrey, and D. Denham (1988). Source parameters of seven large Australian earthquakes determined by body waveform inversion. *Geophysical Journal*, 95, 1–13.
- Friedrich, A. M., B. P. Wernicke, N. A. Niemi, R. A. Bennett, and J. L. Davis (2003). Comparison of geodetic and geologic data from the Wasatch region, Utah, and implications for the spectral character of Earth deformation at periods of 10 to 10 million years. *Journal of Geophysical Research*, 108 (B4), doi:10.1029/2001JB000682.
- Gangopadhyay, A., and P. Talwani (2003). Symptomatic features of intraplate earthquakes. *Seismological Research Letters*, 74, 863–883.
- Gaull, B. A., H. Kagami, H. Taniguchi, *et al.* (1990). Preliminary results of the microzonation of the Perth Metropolitan area using microtremor spectral ratios. *Bureau of Mineral Resources, Geology and Geophysics Record*, 1990/44.
- Gibson, G., V. Wesson, and R. Cuthbertson (1981). Seismicity of Victoria to 1980. *Journal of the Geological Society of Australia*, 28, 341–356.
- Gibson, G., V. Wesson, and T. Jones (1994). The Eugowra NSW earthquake swarm of 1994, *Australian Earthquake Engineering Society Conference Proceedings*, Canberra, ACT, pp. 67–76.
- Glen, R. A. (2005). The Tasmanides of Eastern Australia. In *Terrane Processes at the Margins of Gondwana*, ed. A. P. M. Vaughan, P. T. Leat, and R. J. Pankhurst. Geological Society, London, Special Publication 246, pp. 23–96.
- Gordon, F. R., and J. D. Lewis (1980). The Meckering and Calingiri earthquakes October 1968 and March 1970. *Western Australia Geological Survey Bulletin*, 126.
- Greenhalgh, S. A., and R. Singh (1988). The seismicity of the Adelaide Geosyncline, South Australia. *Bulletin of the Seismological Society of America*, 78, 243–263.
- Greenhalgh, S. A., D. Love, K. Malpas, and R. McDougall (1994). South Australian seismicity, 1980–1992. *Australian Journal of Earth Sciences*, 41, 483–495.
- Hanks, T. C., and A. C. Johnston (1992). Common features of the excitation and propagation of strong ground motion for North American earthquakes. *Bulletin of the Seismological Society of America*, 82, 1–23.
- He, S., and M. Middleton (2002). Heat flow and thermal maturity modelling in the Northern Carnarvon Basin, North West Shelf, Australia. *Marine and Petroleum Geology*, 19, 1073–1088.
- Heimsath, A. M., J. Chappell, W. E. Dietrich, K. Nishiizumi, and R. C. Finkel (2000). Soil production on a retreating escarpment in southeastern Australia. *Geology*, 28, 787–790.
- Heimsath, A. M., J. Chappell, W. E. Dietrich, K. Nishiizumi, and R. C. Finkel (2001). Late Quaternary erosion in southeastern Australia: a field example using cosmogenic isotopes. *Quaternary International*, 83–85, 169–185.
- Hemphill-Haley, M. A., and R. J. Weldon (1999). Estimating prehistoric earthquake magnitude from point measurements of surface rupture. *Bulletin of the Seismological Society of America*, 89, 1264–1279.

- Hill, K. A., G. T. Cooper, M. J. Richardson, and C. J. Lavin (1994). Structural framework of the eastern Otway Basin: inversion and interaction between two major structural provinces. *Exploration Geophysics*, 25, 79–87.
- Hillis, R. R., and S. D. Reynolds (2000). The Australian stress map. *Journal of the Geological Society of London*, 157, 915–921.
- Hillis, R. R., and S. D. Reynolds (2003). In situ stress field of Australia. In *Evolution and Dynamics of the Australian Plate*, ed. R. R. Hillis, and D. Muller. Geological Society of Australia Special Publication 22 and Geological Society of America Special Paper 372, pp. 49–58.
- Hillis, R. R., M. Sandiford, S. D. Reynolds, and M. C. Quigley (2008). Present-day stresses, seismicity and Neogene-to-Recent tectonics of Australia's 'passive' margins: intraplate deformation controlled by plate boundary forces. In *The Nature and Origin of Compression in Passive Margins*, ed. H. Johnson, A. G. Doré, R. W. Gatliff *et al.* Geological Society, London, Special Publication 306, 71–90.
- Historical Records of Australia (1914). The Library Committee of the Commonwealth Parliament, Vol. 1, Series 1, p. 50.
- Hoffman, P. F. (1989). Precambrian geology and tectonic history of North America. In *The Geology of North America – Vol. A, The Geology of North America: An Overview*, ed. A. W. Bally, and A. R. Palmer. Boulder, Colorado: Geological Society of America, pp. 447–512.
- Holdgate, G. R., M. W. Wallace, S. J. Gallagher, *et al.* (2003). Plio-Pleistocene tectonics and eustasy in the Gippsland Basin, southeast Australia: evidence from magnetic imagery and marine geological data. *Australian Journal of Earth Sciences*, 50, 403–426.
- Holdgate, G. R., M. W. Wallace, S. J. Gallagher, *et al.* (2006). Cenozoic fault control on 'deep lead' palaeoriver systems, Central Highlands, Victoria. *Australian Journal of Earth Sciences*, 53, 445–468.
- Holdgate, G. R., I. Cartwright, D. T. Blackburn, *et al.* (2007). The Middle Miocene Yallourn coal seam: the last coal in Australia. *International Journal of Coal Geology*, 70, 95–115.
- Holdgate, G. R., M. W. Wallace, S. J. Gallagher, B. E. Wagstaff, and D. Moore (2008). No mountains to snow on: major post-Eocene uplift of the East Victoria Highlands; evidence from Cenozoic deposits. *Australian Journal of Earth Sciences*, 55, 211–234.
- Holdgate, G. R., M. W. Wallace, S. J. Gallagher, B. E. Wagstaff, and D. Moore (2011). Paleogene basalts prove early uplift of Victoria's Eastern Uplands. *Australian Journal of Earth Sciences*, 58, 95–99.
- Holford, S. P., R. R. Hillis, M. Hand, and M. Sandiford (2011). Thermal weakening localizes intraplate deformation along the southern Australian continental margin. *Earth and Planetary Science Letters*, 305, 207–214.
- Insurance Council of Australia (2013). *Historical Disaster Statistics*, <http://www.insurancecouncil.com.au/industry-statistics-data/disaster-statistics/historical-disaster-statistics>, retrieved 16 December 2013.
- Jenkins, R. J. F., and M. Sandiford (1992). Observations on the tectonic evolution of the southern Adelaide Fold Belt. *Tectonophysics*, 214, 27–36.
- Johnston, A. C. (1994). Seismotectonic interpretations and conclusions from the stable continental region seismicity database. In *The Earthquakes of Stable Continental Regions: Vol. 1, Assessment of Large Earthquake Potential*, ed. A. C. Johnston, K. J. Coppersmith, L. R. Kanter, and C. A. Cornell. Palo Alto, CA: Electric Power Research Institute, 4-1–4-103.

- Johnston, A. C., K. J. Coppersmith, L. R. Kanter, and C. A. Cornell (1994). *The Earthquakes of Stable Continental Regions: Vol. 1, Assessment of Large Earthquake Potential*. Electric Power Research Institute Report, TR102261V1.
- Jones, T. D., G. Gibson, K. F. McCue, *et al.* (1991). Three large intraplate earthquakes near Tennant Creek, Northern Territory, on 22 January 1988. *BMR Journal of Australian Geology and Geophysics*, 12, 339–343.
- Kafka, A. L. (2002). Statistical analysis of the hypothesis that seismicity delineates areas where future large earthquakes are likely to occur in the Central and Eastern United States. *Seismological Research Letters*, 73, 992–1003.
- Keep, M., and D. W. Haig (2010). Deformation and exhumation in Timor: distinct stages of a young orogeny. *Tectonophysics*, 483, 93–111.
- Keep, M., J. Hengesh, and B. Whitney (2012). Natural seismicity and tectonic geomorphology reveal regional transpressive strain in northwestern Australia. *Australian Journal of Earth Sciences*, 59, 341–354.
- Kenner, S. J., and P. Segall (2000). A mechanical model for intraplate earthquakes: application to the New Madrid seismic zone. *Science*, 289, 2329–2332.
- Kennett, B. L. N., A. Fitchner, S. Fishwick, and K. Yoshizawa (2013). Australian Seismological Reference Model (AuSREM): mantle component. *Geophysical Journal International*, 192, 871–887.
- Kroner, A., and U. Cordani (2003). African, southern Indian and South American cratons were not part of the Rodinia supercontinent: evidence from field relationships and geochronology. *Tectonophysics*, 375, 325–352.
- Kusky, T. M., B. F. Windley, and M.-G. Zhai (2007). Tectonic evolution of the North China Block: from orogen to craton to orogen. In *Mesozoic Sub-Continental Lithospheric Thinning Under Eastern Asia*, Geological Society, London, Special Publication 280, 1–34.
- Lamontagne, M., P. Keating, and S. Perreault (2003). Seismotectonic characteristics of the Lower St. Lawrence seismic zone, Quebec: insights from geology, magnetics, gravity, and seismics. *Canadian Journal of Earth Sciences*, 40, 317–336.
- Langenheim, V. E., and T. G. Hildenbrand (1997). Commerce geophysical lineament: its source, geometry, and relation to the Reelfoot rift and New Madrid seismic zone. *Geological Society of America Bulletin*, 109, 580–595.
- Langston, C. A. (1987). Depth of faulting during the 1968 Meckering, Australia, earthquake sequence determined from waveform analysis of local seismograms. *Journal of Geophysical Research*, 92(B11), 11561–11574.
- Lenardic, A., L. Moresi, and H. Muhlhaus (2000). The role of mobile belts for the longevity of deep cratonic lithosphere: the crumple zone model. *Geophysical Research Letters*, 27, 1235–1238.
- Leonard, M. (2002). The Burrakin WA earthquake sequence Sept 2000 – June 2002. *Australian Earthquake Engineering Society Conference: Total Risk Management in the Privatised Era*, Adelaide, Paper 22.
- Leonard, M. (2003). Respite leaves Burakin quaking in anticipation. *Ausgeo News*, 70, 5–7.
- Leonard, M. (2008). One hundred years of earthquake recording in Australia. *Bulletin of the Seismological Society of America*, 98, 1458–1470.
- Leonard, M. (2010). Earthquake fault scaling: self-consistent relating of rupture length, width, average displacement, and moment release. *Bulletin of the Seismological Society of America*, 100, 1971–1988.
- Leonard, M. (2012). Earthquake recurrence parameterisation. In *The 2012 Australian Earthquake Hazard Map*, ed. D. R. Burbidge. *Geoscience Australia Record*, 2012/71, 27–43.

- Leonard, M., and D. Clark (2006). Reconciling neotectonic and seismic recurrence rates in SW WA. *Proceedings of the Australian Earthquake Engineering Society Conference*, Paper 19, 19–24.
- Leonard, M., and D. Clark (2011). A record of stable continental region earthquakes from Western Australia spanning the late Pleistocene: insights for contemporary seismicity. *Earth and Planetary Science Letters*, 309, 207–212.
- Leonard, M., I. D. Ripper, and Y. Li (2002). Australian earthquake fault plane solutions. *Geoscience Australia Record*, 2002/19.
- Leonard, M., D. Darby, and G. Hu (2007). GPS-geodetic monitoring of the South West Seismic Zone of Western Australia: progress after two observation epochs in 2002 and 2006. *Proceedings of the Australian Earthquake Engineering Society Conference*, Wollongong, Paper 28.
- Leonard, M., A. McPherson, and D. Clark (2012). Source zonation. In *The 2012 Australian Earthquake Hazard Map*, ed. D. R. Burbidge. *Geoscience Australia Record*, 2012/71, 17–26.
- Lewis, J. D., N. A. Daetwyler, J. A. Bunting, and J. S. Moncrieff (1981). The Cadoux earthquake. *Western Australia, Geological Survey Report*, 1981/11, 133.
- Li, Q., M. Liu, and S. Stein (2009). Spatiotemporal complexity of continental intraplate seismicity: insights from geodynamic modeling and implications for seismic hazard estimation. *Bulletin of the Seismological Society of America*, 99, 52–60.
- Liu, L., and M. D. Zoback (1997). Lithospheric strength and intraplate seismicity in the New Madrid seismic zone. *Tectonics*, 16, 585–595.
- Liu, M., S. Stein, and H. Wang (2011). 2000 years of migrating earthquakes in North China: how earthquakes in midcontinents differ from those at plate boundaries. *Lithosphere*, doi: 10.1130/L129.1.
- Liu, M., and H. Wang (2012). Roaming earthquakes in China highlight midcontinental hazards. *Eos, Transactions, American Geophysical Union*, 93, 453–454.
- Love, D. (1996). *Seismic Hazard and Microzonation of the Adelaide Metropolitan Area*, South Australia, Department of Mines and Energy, Report Book 96/27.
- Machette, M. N., S. F. Personius, A. R. Nelson, D. P. Schwartz, and W. R. Lund (1991). The Wasatch fault zone, Utah: segmentation and history of Holocene earthquakes. *Journal of Structural Geology*, 13, 137–149.
- Machette, M. N., A. J. Crone, and J. R. Bowman (1993). Geologic investigations of the 1986 Marryat Creek, Australia, earthquake: implications for paleoseismicity in stable continental regions. *Bulletin U.S. Geological Survey*, 2032-B.
- Malpas, K. L. (1991). Seismic risk in South Australia. BSc (Hons) thesis (unpublished). Department of Earth Sciences, Flinders University of South Australia, Adelaide.
- Marco, S., M. Stein, and A. Agnon (1996). Long term earthquake clustering: a 50,000-year paleoseismic record in the Dead Sea Graben. *Journal of Geophysical Research*, 101, 6179–6191.
- Mazzotti, S. (2007). Geodynamic models for earthquake studies in intraplate North America. In *Continental Intraplate Earthquakes: Science, Hazard, and Policy Issues*, ed. S. Stein, and S. Mazzotti. Geological Society of America Special Paper 425, pp. 17–33.
- McCue, K. (1990). Australia's large earthquakes and recent fault scarps. *Journal of Structural Geology*, 12, 761–766.
- McCue, K. (1996). Atlas of isoseismal maps of Australian earthquakes: Part 3. *AGSO Record*, 1995/44.
- McCue, K. (2004). Australia: historical earthquake studies. *Annals of Geophysics*, 47, 387–397.

- McCue, K., V. Wesson, and G. Gibson (1990). The Newcastle, New South Wales, earthquake of 28 December 1989. *BMR Journal of Australian Geology and Geophysics*, 11, 559–567.
- McCue, K. F. (1975). *Seismicity and Seismic Risk in South Australia*. University of Adelaide Report, ADP 137.
- McCue, K. F. (1980). Magnitudes of some early earthquakes in Southeastern Australia. *Search*, 11(3), 78–80.
- McCue, K. F. (1989). *Australian Seismological Report 1985*, Bureau of Mineral Resources, Geology & Geophysics Report 285.
- McCue, K. F., and M. O. Michael-Leiba (1993). Australia's deepest known earthquake. *Seismology Research Letters*, 64(34), 201–206.
- McPherson, A., D. Clark, and T. Barrows (2009). Long-term slip rates for onshore faults in the Gippsland Basin, south-eastern Australia. *7th International Conference on Geomorphology (ANZIAG)*, Melbourne, Australia.
- McPherson, A., D. Clark, M. Cupper, C. D. N. Collins, and G. Nelson (2012a). The Cadell Fault: a record of long-term fault behaviour in south-eastern Australia. *Proceedings Second Australian Regolith Geoscientists Association Conference/22nd Australian Clay Minerals Society Conference*, Mildura, Victoria, February 7–9, 2012, pp. 7–16.
- McPherson, A., D. Clark, M. MacPhail, and M. Cupper (2012b). Neotectonism on the eastern Australian passive margin: evidence from the Lapstone Structural Complex. *Proceedings 15th Australia New Zealand Geomorphology Group Conference*, Bundanoon, NSW, December 2–7, 2012.
- Michael-Leiba, M. O. (1987). Temporal variation in seismicity of the southwest seismic zone, Western Australia: implications for earthquake risk assessment. *BMR Journal of Australian Geology and Geophysics*, 10, 133–137.
- Michael-Leiba, M. (1989). Macroseismic effects, locations and magnitudes of some early Tasmanian earthquakes. *BMR Journal of Australian Geology and Geophysics*, 11, 89–99.
- Michael-Leiba, M., D. Love, K. McCue, and G. Gibson (1994). The Uluru (Ayers Rock), Australia, earthquake of 28 May 1989. *Bulletin of the Seismological Society of America*, 84, 209–214.
- Mogi, K. (1963). Some discussions on aftershocks, foreshocks and earthquake swarms: the fracture of a semi-infinite body caused by an inner stress origin and its relation to the earthquake phenomena, 3. *Bulletin of the Earthquake Research Institute, Tokyo University*, 41, 615–658.
- Mooney, W. D., J. Ritsema, and Y. K. Hwang (2012). Crustal seismicity and the earthquake catalog maximum moment magnitude (M_{cmax}) in stable continental regions (SCRs): correlation with the seismic velocity of the lithosphere. *Earth and Planetary Science Letters*, 357–358, 78–83.
- Moye, D. G., K. R. Sharp, and D. H. Stapledon (1963). *Geology of the Snowy Mountains Region*. Snowy Mountains Hydro-electric Authority, Cooma (unpublished: held in Geological Survey of New South Wales Library).
- Norvick, M. S. (2004). Tectonic and stratigraphic history of the Perth Basin. *Geoscience Australia, Record*, 2004/16.
- Norvick, M. S., and M. A. Smith (2001). Mapping the plate tectonic reconstruction of southern and southeastern Australia and implications for petroleum systems. *APEA Journal*, 41, 15–35.
- Paine, M. D., D. A. Bennetts, J. A. Webb, and V. J. Morand (2004). Nature and extent of Pliocene strandlines in southwestern Victoria and their application to Late Neogene tectonics. *Australian Journal of Earth Sciences*, 51, 407–422.

- Pandey, O. P., K. Chandrakala, G. Parthasarathy, P. R. Reddy, and G. K. Reddy (2008). Upwarped high velocity mafic crust, subsurface tectonics and causes of intra plate Latur-Killari (M 6.2) and Koyna (M 6.3) earthquakes, India: a comparative study. *Journal of Asian Earth Sciences*, doi: 10.1016/j.jseas.2008.11.014.
- Paul, E., T. Flottmann, and M. Sandiford (1999). Structural geometry and controls on basement-involved deformation in the northern Flinders Ranges, Adelaide Fold Belt, South Australia. *Australian Journal of Earth Sciences*, 46, 343–354.
- Pickett, J. W., and P. Bishop (1992). Aspects of landscape evolution in the Lapstone Monocline area, New South Wales. *Australian Journal of Earth Sciences*, 39, 21–28.
- Pillans, B. (2012). How old is Lake George? *Australian and New Zealand Geomorphology Group 15th Conference*, Bundanoon, NSW, December 2–7, 2012, Program and Abstracts 79.
- Quigley, M. C., M. L. Cupper, and M. Sandiford (2006). Quaternary faults of south-central Australia: palaeoseismicity, slip rates and origin. *Australian Journal of Earth Sciences*, 53, 285–301.
- Quigley, M., M. Sandiford, A. Alimanovic, and L. K. Fifield (2007a). Landscape responses to intraplate tectonism: quantitative constraints from ^{10}Be abundances. *Earth and Planetary Science Letters*, 261, 120–133.
- Quigley, M., M. Sandiford, K. Fifield, and A. Alimanovic (2007b). Bedrock erosion and relief production in the northern Flinders Ranges, Australia. *Earth Surface Processes and Landforms*, 32, 929–944.
- Quigley, M. C., M. Sandiford, and M. L. Cupper (2007c). Distinguishing tectonic from climatic controls on range-front sedimentation, Flinders Ranges, South Australia. *Basin Research*, doi: 10.1111/j.1365–2117.2007.00336.x.
- Quigley, M., D. Clark, and M. Sandiford (2010). Late Cenozoic tectonic geomorphology of Australia. In *Australian Landscapes*, ed. P. Bishop and B. Pillans. Geological Society, London, Special Publication 346, 243–265.
- Rajendran, C. P. (2000). Using geological data for earthquake studies: a perspective from peninsular India. *Current Science*, 79, 1251–1258.
- Rajendran, C. P., K. Rajendran, and B. John (1996). The 1993 Killari (Latur), central India, earthquake: an example of fault reactivation in the Precambrian crust. *Geology*, 24, 651–654.
- Rajendran, K., C. P. Rajendran, M. Thakkar, and M. P. Tuttle (2001). The 2001 Kutch (Bhuj) earthquake: coseismic surface features and their significance. *Current Science*, 80, 1397–1405.
- Reid, B. (2007). A Neotectonic study of the Willunga and Milendella Faults, South Australia. B.Sc. (Hons) thesis, School of Earth Sciences, The University of Melbourne, Melbourne.
- Reinecker, J., O. Heidbach, and B. Mueller (2003). The 2003 release of the world stress map (available online at www.world-stress-map.org).
- Revets, S. A., M. Keep, and B. L. N. Kennett (2009). NW Australian intraplate seismicity and stress regime. *Journal of Geophysical Research*, 114, B10305.
- Reynolds, S. D., D. D. Coblenz, and R. R. Hillis (2002). Tectonic forces controlling the regional intraplate stress field in continental Australia: results from new finite element modeling. *Journal of Geophysical Research, Solid Earth*, 107(B7) 2131, 10.1029/2001JB000408.
- Richardson, R. M. (1992). Ridge forces, absolute plate motions, and the intraplate stress field. *Journal of Geophysical Research*, 97, 11739–11749.

- Ripper, I. D. (1963). Local and regional earthquakes recorded by the Tasmania Seismic Net. B.Sc thesis, University of Tasmania, Hobart.
- Ritz, J. F., E. T. Brown, D. L. Bourles, *et al.* (1995). Slip rates along active faults estimated with cosmic-ray-exposure dates: application to the Bogd Fault, Gobi-Altai, Mongolia. *Geology*, 23, 1019–1022.
- Robson, T. C., and J. A. Webb (2011). Late Neogene tectonics in northwestern Victoria: evidence from the Late Miocene–Pliocene Loxton Sand. *Australian Journal of Earth Sciences*, 58, 579–586.
- Rutherford, I. D., and C. Kenyon (2005). Geomorphology of the Barmah-Millewa forest. *Proceedings of the Royal Society of Victoria*, 117, 23–39.
- Rynn, J. M. W., D. Denham, S. A. Greenhalgh, *et al.* (1987). Atlas of isoseismal maps of Australian earthquakes, Part 2. *Bureau of Mineral Resources Bulletin*, 222.
- Sandiford, D., and G. Gibson (2012). The Moe/Thorpedale earthquake: preliminary investigation. *Proceedings of the Australian Earthquake Engineering Society Conference*, Tweed Heads, Australia, December 7–9, 2012, Paper 25.
- Sandiford, M. (2003a). Geomorphic constraints on the late Neogene tectonics of the Otway Ranges. *Australian Journal of Earth Sciences*, 50, 69–80.
- Sandiford, M. (2003b). Neotectonics of southeastern Australia: linking the Quaternary faulting record with seismicity and in situ stress. In *Evolution and Dynamics of the Australian Plate*, ed. R. R. Hillis, and D. Muller. Geological Society of Australia Special Publication 22 and Geological Society of America Special Paper 372, 101–113.
- Sandiford, M., and D. L. Egholm (2008). Enhanced intraplate seismicity along continental margins: some causes and consequences. *Tectonophysics*, 457, 197–208.
- Sandiford, M., and M. C. Quigley (2009). Topo-Oz: insights into the various modes of intraplate deformation in the Australian continent. *Tectonophysics*, 474, 405–416.
- Sandiford, M., M. Wallace, and D. Coblenz (2004). Origin of the in situ stress field in southeastern Australia. *Basin Research*, 16, 325–338.
- Sandiford, M., D. D. Coblenz, and W. P. Schellart (2005). Evaluating slab-plate coupling in the Indo-Australian plate. *Geology*, 33, 113–116.
- Schulte, S. M., and W. D. Mooney (2005). An updated global earthquake catalogue for stable continental regions: reassessing the correlation with ancient rifts. *Geophysical Journal International*, 161, 707–721.
- Schwartz, D. P., and K. J. Coppersmith (1984). Fault behaviour and characteristic earthquakes: examples from the Wasatch and San Andreas Fault Zones. *Journal of Geophysical Research*, 89(B7), 5681–5698.
- Seeber, L., G. Ekstrom, S. K. Jain, *et al.* (1996). The 1993 Killari earthquake in central India: a new fault in Mesozoic basalt flows? *Journal of Geophysical Research*, 101, 8543–8560.
- Sheard, M. J. (1995). Quaternary volcanic activity and volcanic hazards. In *The Geology of South Australia*, Vol. 2, ed. J. F. Drexel, and W. V. Preiss. *Geological Survey of South Australia, Bulletin*, 54, 264–268.
- Sinadinovski, C., and K. McCue (2010). Testing the hypothesis that earthquake hazard is uniform across Australia. *Proceedings Australian Earthquake Engineering Society Conference*, Perth, Western Australia, Paper 42.
- Sinadinovski, C., T. Jones, D. Stewart, and N. Corby (2002). Earthquake history, regional seismicity and the 1989 Newcastle earthquake. In *Earthquake Risk in Newcastle and Lake Macquarie*, ed. T. Dhu, and T. Jones. Geoscience Australia Record, 2002/15.

- Sinha, S., and S. Mohanty (2012). Spatial variation of crustal strain in the Kachchh region, India: implication on the Bhuj earthquake of 2001. *Journal of Geodynamics*, 61, 1–11.
- Sloan, R. A., J. A. Jackson, D. McKenzie, and K. Priestley (2011). Earthquake depth distributions in central Asia, and their relations with lithospheric thickness, shortening and extension. *Geophysical Journal International*, 185, 1–29.
- Smith, D. E., and R. Kolenkiewicz (1990). Tectonic motion and deformation from satellite laser ranging to LAGEOS. *Journal of Geophysical Research*, 95, 22,013–22,041.
- Somerville, P., P. Quijada, H. K. Thio, M. Sandiford, and M. Quigley (2008). Contribution of identified active faults to near fault seismic hazard in the Flinders Ranges. *Australian Earthquake Engineering Society Meeting*, Paper 45.
- Somerville, P., R. W. Graves, N. F. Collins, S. G. Song, and S. Ni (2010). *Ground Motion Models for Australian Earthquakes*. Final report to Geoscience Australia, 30 March 2010, URS Corporation, Pasadena, CA.
- Stein, R. S., A. A. Barka, and J. H. Dieterich (1997). Progressive failure on the North Anatolian fault since 1939 by earthquake stress triggering. *Geophysical Journal International*, 128, 594–604.
- Stein, S., and M. Liu (2009). Long aftershock sequences within continents and implications for earthquake hazard assessment. *Nature*, 462, 87–89.
- Stein, S. S., S. Cloetingh, N. H. Sleep, and R. Wortel (1989). Passive margin earthquakes, stresses and rheology. In *Earthquakes at North American Passive Margins: Neotectonics and Postglacial Rebound*, ed. S. Gregersen and P. Basham. NATO ASI Series C, Boston, MA: Kluwer Academic, pp. 231–259.
- Stirling, M., D. Rhoades, and K. Berryman (2002). Comparison of earthquake scaling relations derived from data of the instrumental and preinstrumental era. *Bulletin of the Seismological Society of America* 92, 812–830.
- Stuart, W. D., T. G. Hildenbrand, and R. W. Simpson (1997). Stressing the New Madrid Seismic Zone by a lower crustal detachment fault. *Journal of Geophysical Research*, 102, 27623–27633.
- Sutton, D. J., K. F. McCue, and A. Bugeja (1977). Seismicity of the southeast of South Australia. *Australian Journal of Earth Sciences*, 24, 357–364.
- Sykes, L. R. (1978). Intraplate seismicity, reactivation of preexisting zones of weakness, alkaline magmatism, and other tectonism postdating continental fragmentation. *Reviews of Geophysics and Space Physics*, 16, 621–688.
- Talwani, P. (1988). The intersection model for intraplate earthquakes. *Seismological Research Letters*, 59, 305–310.
- Talwani, P. (1999). Fault geometry and earthquakes in continental interiors. *Tectonophysics*, 305, 371–379.
- Talwani, P., and K. Rajendran (1991). Some seismological and geometric features of intraplate earthquakes. *Tectonophysics*, 186, 19–41.
- Talwani, P., and W. T. Schaeffer (2001). Recurrence rates of large earthquakes in the South Carolina coastal plain based on paleoliquefaction data. *Journal of Geophysical Research*, 106, 6621–6642.
- Tomkins, K. M., G. S. Humphreys, M. T. Wilkinson, *et al.* (2007). Contemporary versus long-term denudation along a passive plate margin: the role of extreme events. *Earth Surface Processes and Landforms*, 32, 1013–1031.
- Tregonning, P. (2003). Is the Australian Plate deforming? A space geodetic perspective. In *Evolution and Dynamics of the Australian Plate* ed. R. R. Hillis, and D. Muller. Geological Society of Australia Special Publication 22 and Geological Society of America Special Publication 372, pp. 41–48.

- Tuttle, M. P., E. S. Schweig, J. D. Sims, *et al.* (2002). The earthquake potential of the New Madrid seismic zone. *Bulletin of the Seismological Society of America*, 92, 2080–2089.
- van Arsdale, R. (2000). Displacement history and slip rate on the Reelfoot fault of the New Madrid seismic zone. *Engineering Geology*, 55, 219–226.
- van de Graaff, W. J. E., P. D. Denman, and R. M. Hocking (1976). Emerged Pleistocene marine terraces on Cape Range, Western Australia. *Western Australia Geological Survey Annual Report 1975*, 62–70.
- van der Beek, P., A. Pulford, and J. Braun (2001). Cenozoic landscape development in the Blue Mountains (SE Australia): lithological and tectonic controls on rifted margin morphology. *The Journal of Geology*, 109, 35–56.
- van Lanen, X., and W. D. Mooney (2007). Integrated geologic and geophysical studies of North American continental intraplate seismicity. In *Continental Intraplate Earthquakes: Science, Hazard, and Policy Issues*, ed. S. Stein, and S. Mazzotti. Geological Society of America Special Paper 425, 101–112.
- VandenBerg, A. H. M. (2010). Paleogene basalts prove early uplift of Victoria's Eastern Uplands. *Australian Journal of Earth Sciences*, 57, 291–315.
- Veevers, J. J. (2000). *Billion-Year Earth History of Australia and Neighbours in Gondwanaland*. Sydney, Australia: GEMOC Press.
- Vlahovic, G., C. Powell, and M. Lamontagne (2003). A three-dimensional P wave velocity model for the Charlevoix seismic zone, Quebec, Canada. *Journal of Geophysical Research*, 108, 2439, doi:10.1029/2002JB002188.
- Vogfjörð, K. S., and C. A. Langston (1987). The Meckering earthquake of 14th October 1968: a possible downward propagating rupture. *Bulletin of the Seismological Society of America*, 77, 1558–1578.
- Wald, D. J., V. Quitoriano, B. Worden, M. Hopper, and J. W. Dewey (2011). USGS “Did You Feel It?” Internet-based macroseismic intensity maps. *Annals of Geophysics*, 54, 688–707.
- Wallace, R. E. (1987). Grouping and migration of surface faulting and variations in slip rates on faults in the Great Basin province. *Bulletin of the Seismological Society of America*, 77, 868–876.
- Weissel, J. K., and M. A. Seidl (1998). Inland propagation of erosional escarpments and river profile evolution across the southeast Australian passive continental margin. In *Rivers Over Rock: Fluvial Processes in Bedrock Channels*, ed. K. J. Tinkler and E. E. Wohl. American Geophysical Union, Geophysical Monograph 107, pp. 189–206.
- Wells, D. L., and K. J. Coppersmith (1994). New empirical relationships among magnitude, rupture length, rupture width, rupture area, and surface displacement. *Bulletin of the Seismological Society of America*, 84, 974–1002.
- Wheeler, R. L. (1995). Earthquakes and the cratonward limit of Iapetan faulting in eastern North America. *Geology*, 23, 105–108.
- Wheeler, R. L. (1996). Earthquakes and the southeastern boundary of the intact Iapetan margin in Eastern North America. *Seismological Research Letters*, 67, 77–83.
- Wheeler, R. L. (2009a). *Methods of M_{\max} Estimation East of the Rocky Mountains*. U.S. Geological Survey Open-File Report 2009–1018.
- Wheeler, R. L. (2009b). *Sizes of the Largest Possible Earthquakes in the Central and Eastern United States: Summary of a Workshop, September 8–9, 2008, Golden, Colorado*. U.S. Geological Survey Open-File Report 2009–1263.
- Wheeler, R. L. (2011). Reassessment of stable continental regions of Southeast Asia. *Seismological Research Letters*, 82, 971–983.

- Wheeler, R. L., and A. Frankel (2000). Geology in the 1996 USGS Seismic-hazard Maps, Central and Eastern United States. *Seismological Research Letters*, 71, 273–282.
- Wilkinson, M. T., J. Chappell, G. S. Humphreys, *et al.* (2005). Soil production in heath and forest, Blue Mountains, Australia: influence of lithology and palaeoclimate. *Earth Surface Processes and Landforms*, 30, 923–934.
- Zoback, M. D. (1992). First- and second-order patterns of stress in the lithosphere: the world stress map project. *Journal of Geophysical Research*, 97, 11703–11728.