

# NMMB331 - HW2

Jan Oupický

## 1

Let's assume  $L : \mathbb{F}_2^n \rightarrow \mathbb{F}_2^m$ . Then

$$\text{Im}(L^*)^\perp = \{y \in \mathbb{F}_2^n \mid \forall x \in \text{Im}(L^*) : \langle x, y \rangle = 0\} = \{y \in \mathbb{F}_2^n \mid \forall x \in \mathbb{F}_2^m : \langle L^*(x), y \rangle = 0\}$$

From lecture we know that  $\forall x, y : \langle L^*(x), y \rangle = \langle x, L(y) \rangle \implies$

$$\text{Im}(L^*)^\perp = \{y \in \mathbb{F}_2^n \mid \forall x \in \mathbb{F}_2^m : \langle x, L(y) \rangle = 0\}$$

We know that for a fixed  $v \in \mathbb{F}_2^m$  if  $v$  satisfies  $\forall x \in \mathbb{F}_2^m : \langle x, v \rangle = 0$  then  $v = \underline{0}$ . Using this fact we can see that  $\text{Im}(L^*)^\perp$  is exactly those  $y \in \mathbb{F}_2^n$  such that  $L(y) = 0$ . That is the definition of  $\text{Ker}(L)$ .

## 2

Let  $G, F$  be EA-equivalent vectorial boolean functions i.e.:

$$G = A_1 \circ F \circ A_2 + A_3 \text{ where for } i = 1, 2, 3: A_i(x) = L_i(x) + b \quad (1)$$

and  $L_i$  is a linear permutation. As in the lecture, choose  $u, v$  and we will show that  $|\hat{G}(u, v)|$  corresponds to  $|\hat{F}(a, b)|$  for exactly one pair  $(a, b)$ . Note that we can write:

$$G(x) = L_1 \circ F(L_2(x) + b_2) + b_1 + L_3(x) + b_3$$

Denote  $z = L_2(x) + b_2$  then  $x = L_2^{-1}(z - b_2) = L_2^{-1}(z) + L_2^{-1}(b_2)$  ( $L_2$  is a linear permutation).

$$\begin{aligned} \hat{G}(u, v) &= \sum_{x \in \mathbb{F}_2^n} \chi(u \cdot G(x) + v \cdot x) = \sum_{x \in \mathbb{F}_2^n} \chi(u \cdot (L_1 \circ F(z) + b_1 + L_3(x) + b_3) + v \cdot x) = \\ &= \chi(u \cdot (b_1 + b_3)) \sum_{z \in \mathbb{F}_2^n} \chi(u \cdot (L_1 \circ F(z) + L_3 \circ L_2^{-1}(z) + L_3 \circ L_2^{-1}(b_2))) + v \cdot (L_2^{-1}(z) + L_2^{-1}(b_2))) = \end{aligned}$$

Denote  $c = \chi(u \cdot (b_1 + b_3 + L_3 \circ L_2^{-1}(b_2)) + v \cdot L_2^{-1}(b_2))$ . By definition  $c = \pm 1$ . So we can ommit it.

$$\begin{aligned}
|\hat{G}(u, v)| &= \sum_{z \in \mathbb{F}_2^n} \chi(u \cdot (L_1 \circ F(z) + L_3 \circ L_2^{-1}(z)) + v \cdot L_2^{-1}(z)) = \\
&\sum_{z \in \mathbb{F}_2^n} \chi(u \cdot L_1 \circ F(z) + u \cdot L_3 \circ L_2^{-1}(z) + v \cdot L_2^{-1}(z)) = \\
&\sum_{z \in \mathbb{F}_2^n} \chi(L_1^*(u) \cdot F(z) + (L_3 \circ L_2^{-1})^*(u) \cdot z + (L_2^{-1})^*(v) \cdot z) = \\
&\sum_{z \in \mathbb{F}_2^n} \chi(L_1^*(u) \cdot F(z) + ((L_3 \circ L_2^{-1})^*(u) + (L_2^{-1})^*(v)) \cdot z) \implies \\
|\hat{G}(u, v)| &= |\hat{F}(a, b)| \text{ where} \\
a &= L_1^*(u) \\
b &= (L_3 \circ L_2^{-1})^*(u) + (L_2^{-1})^*(v)
\end{aligned}$$

$a, b$  are determined uniquely since  $L_i$ s (and equivalently  $L_i^*$ s) are permutations.