

Relativity Assignments

Pranav Satheesh

July 17, 2021

1 Einstein's velocity addition

In this problem, you will derive the velocity addition rule in relativity using Lorentz transformations. Consider a particle A in the frame B. Her velocity w.r.t frame B is V_{AB} . In another frame C moving w.r.t B with a velocity V_{CB} . Show that the velocity of A w.r.t to the frame C using **Lorentz transformation**:

$$V_{AC} = \frac{V_{AB} + V_{BC}}{1 + (V_{AB}V_{BC}/c^2)} \quad (1)$$

(**Hint** : You can use consider the velocity to be $u = \frac{dx}{dt}$ in the frame B and then write the lorentz transformation equations for dx and dt to the frame C where these become $d\bar{x}$ and $d\bar{t}$)

2 Superluminal Motion

Astronomers observed radio galaxies moving with velocities exceeding the velocity of c! M87 is an example in the Virgo cluster The distance to this galaxy, M87, is about $D = 62$ million light years. One can use this distance to convert angular separations into linear separations across the line of sight.




Figure 1: Blobs

1. Now, look at one of the blobs: the innermost one, which appears most clearly in 1996 and 1997. Approximately calculate the velocity of the blob. Hold on a minute! Is this against special relativity postulate 2?
2. This calculation is not correct. The clouds are actually moving straight towards us with a velocity below c . As the cloud rapidly approaches, the distance light has to travel reduces and hence it would appear to reach us sooner. This effect can be understood from the second diagram. The cloud starts at N and moves at an angle θ w.r.t the line of sight of the observer O . Let \vec{V} be the velocity of the cloud and let t_{obs} be the time the observer sees the light emitted from the cloud at time t . From the distance travelled by light calculate the relation between t_{obs} and t .




Figure 2: Apparent motion of the cloud

3. Now we use the t_{obs} relation, figure out the transverse speed V_T observed by the observer using $V_T = dx/dt_{obs}$.
4. Can you now plot V_T for different θ and see at what angles you see an apparent motion greater than c ?

3 Relativistic Doppler Effect

In this exercise you will derive the Doppler effect using Lorentz transformations.

Consider the source to be at rest and the receiver is moving away with some velocity v and $v > 0$. Use subscript s for source and r for receiver.

1. Consider the first pulse received by the receiver at $t_s = 0$ and $x_s = 0$. λ_s is the wavelength of the wave. Now at time t_{rs} the receiver receives the second pulse. Then show that

$$ct_{rs} = \lambda_s + vt_{rs}$$

2. We now need to make a lorentz transformation to reciever frame and change t_{rs} to t_r . Show that after doing Lorentz transformation

$$t_r = \frac{t_{rs}}{\gamma}$$

3. Now subsitiute this in the previous relation and obtain the fampus relation between the frequencies.

$$f_r = \sqrt{\frac{1 - \frac{v}{c}}{1 + \frac{v}{c}}} f_s$$