

# QSPR ensemble modelling of the 1:1 and 1:2 complexation of $\text{Co}^{2+}$ , $\text{Ni}^{2+}$ , and $\text{Cu}^{2+}$ with organic ligands: relationships between stability constants

Vitaly Solov'ev · Alexandre Varnek ·  
Aslan Tsivadze

Received: 24 January 2014 / Accepted: 1 April 2014 / Published online: 16 April 2014  
© Springer International Publishing Switzerland 2014

**Abstract** Quantitative structure–property relationship (QSPR) modeling of stability constants for the metal:ligand ratio 1:1 ( $\log K$ ) and 1:2 ( $\log \beta_2$ ) complexes of 3 transition metal ions with diverse organic ligands in aqueous solution was performed using ensemble multiple linear regression analysis and substructural molecular fragment descriptors. The modeling was performed on the sets containing 396 and 132 ( $\text{Co}^{2+}$ ), 613 and 233 ( $\text{Ni}^{2+}$ ), 883 and 257 ( $\text{Cu}^{2+}$ )  $\log K$  and  $\log \beta_2$  values, respectively. The models have been validated in external fivefold cross-validations procedure as well as on the external test set containing new ligands recently reported in the literature. Predicted  $\log K$  and  $\log \beta_2$  values were calculated as arithmetic means of several hundred individual models (consensus models) using their applicability domains in averaging. The root mean squared error of predictions varies from 0.94 to 1.2 ( $\log K$ ) and from 1.2 to 1.4 ( $\log \beta_2$ ) which is close to observed experimental systematic errors. Linear correlations between experimental  $\log K$  values for pair of metal ions were evaluated. For all metal ions and ligands forming both 1:1 and 1:2 complexes the following ratio is observed:  $\log \beta_2 / \log K = 1.8 \pm 0.1$ ,  $n = 492$ .

**Electronic supplementary material** The online version of this article (doi:10.1007/s10822-014-9741-3) contains supplementary material, which is available to authorized users.

V. Solov'ev (✉) · A. Tsivadze  
Institute of Physical Chemistry and Electrochemistry,  
Russian Academy of Sciences, Leninskiy Prospekt, 31,  
119071 Moscow, Russia  
e-mail: solovev-vp@mail.ru

A. Varnek  
Laboratoire de Chimoinformatique, UMR 7140 CNRS,  
Université de Strasbourg, 1, rue Blaise Pascal, 67000 Strasbourg,  
France

**Keywords** QSPR modeling of stability constants · Complexes of  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$ , and  $\text{Cu}^{2+}$  with organic ligands in water · Multiple linear regression analysis, substructural molecular fragment descriptors · Stability of complexes with different stoichiometry

## Introduction

Binding of metal ions (M) by organic ligands (L) in solutions is of great importance in many chemical [1–3] and biological [4–11] processes. The thermodynamic quantities such as equilibrium constants and free energy are the most general and significant characteristics used in chemical speciation calculations of metal binding [12–18]. In the past decade, quantitative structure–property relationships (QSPR) linking the stability constant with the structure of organic ligands were reported [19–40] and reviewed [41, 42]. Developed QSPR models help to assess a selectivity of ligand for metal ions [40] which is a key parameter in biomedical, biological, and environmental applications, and in metal ion separations [43]. Technological importance of the  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$  metal ions is well known, they are often widely used in materials for high-tech applications [44, 45] and objects of ion sequestering in biological systems [46]. Earlier, QSPR modeling of the stability constant  $\log K$  for the 1:1 (M:L) complexes of  $\text{Co}^{2+}$  [20, 47, 48],  $\text{Ni}^{2+}$  [20, 25, 47, 48],  $\text{Cu}^{2+}$  [20, 25, 47–49] were reported on relatively small data sets and particular classes of ligands including from 6 ( $\text{Cu}^{2+}$ ) [49] to 85 ( $\text{Ni}^{2+}$ ) and 96 ( $\text{Cu}^{2+}$ ) [25] ligands. These models were either not properly validated [49] or their validation have been performed on a single test sets [20, 25]. The models applicability domains (ADs) issue was never discussed in these studies [20, 25, 47–49]. In most of cases, QSPR

modeling of the  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$  complexation was performed on the 1:1 complexes. At the same time, estimation of stability of the complexes with different stoichiometry is important for quantitative description of industrial and biological processes [50, 51], and for design of new materials [44]. In particular, models for the stability constant ( $\log\beta_2$ ) of the 1:2 complexes could be an obvious step in stability assessment of multiligand and metallosupramolecular complexes.

In this paper, we describe QSPR modeling of both the stability constants for the 1:1 and 1:2 (M:L) complexes of  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$  with different classes of organic ligands in water at 298 K and an ionic strength 0.1 M. The models were obtained using ensemble multiple linear regression analysis (MLR) and substructural molecular fragment (SMF) descriptors. The modeling was performed on the biggest known data sets of the complexation containing 396 and 132 ( $\text{Co}^{2+}$ ), 613 and 233 ( $\text{Ni}^{2+}$ ), 883 and 257 ( $\text{Cu}^{2+}$ ) organic ligands for  $\log K$  and  $\log\beta_2$ , respectively. The models have been validated in external 5-fold cross-validations procedure and on an additional test set of the 23 complexes of  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$  with 13 organic ligands taking into account ADs of models. The root mean squared error (RMSE) of predictions varies from 0.94 to 1.2 ( $\log K$ ) and from 1.2 to 1.4 ( $\log\beta_2$ ) which is close to observed systematic experimental errors. Linear relationships linking  $\log K$  for pairs of metal ions and known ligands were evaluated. Sufficiently general linear relationship between  $\log\beta_2$  and  $\log K$  has been suggested. Developed QSPR models were applied for assessments of the stability constant  $\log K$  of the ML complexes of the ligands which form supramolecular metallogrid complexes.

## Materials and methods

### Data sets

The experimental values of the stability constants for the 1:1 and 1:2 (M:L) complexes of  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$  with various organic ligands in water were selected from the IUPAC Stability Constants Database (SC DB) (version 5.33, Academic Software) [52] at standard temperature 298 K and an ionic strength  $I = 0.1$  M. Some  $\log K$  and  $\log\beta_2$  values (around 15 %) were corrected to specified temperature and an ionic strength using the procedures included in SC DB. Debye–Hückel theory and the Davies equation for the mean ionic activity coefficient were applied to adjust the stability constant from  $I' = 0$ –0.3 M to an ionic strength  $I = 0.1$  M [52]. If an experimental enthalpy ( $\Delta H$ ) value of an equilibrium is available, and it is assumed that  $\Delta H$  is independent of temperature, the van't

Hoff equation in the integration form was used to assess  $\log K$  ( $\log\beta_2$ ) at 298 K from  $\log K$  ( $\log\beta_2$ ) measures at any other temperature  $T'$  ( $T' = 283$ –308 K) [52].














2D structures of ligands, names of metal ions and corresponding  $\log K$  and  $\log\beta_2$  values resulted from searching in SC DB were converted into structure-data files (SDF) served as an input in the MLR module of the ISIDA/QSPR program [53]. Chemical editor EdChemS [54] and the SDF manager EdiSDF [23, 55–57] were used to prepare data sets. If several values of  $\log K$  ( $\log\beta_2$ ) were available for a particular ligand, the most recent data or the data consistent with respect to different experimental methods were chosen. It should be noted that discrepancies in experimental  $\log K$  values reported for the same equilibrium by different authors may attain rather high values (more than 2.8  $\log K$  units, Table 1). At the pretreatment stage, some specific ligands were excluded if they bring fragments occurred less than in 3 ligands for different types of SMF descriptors and given metal ion. Finally, 396 and 132 ( $\text{Co}^{2+}$ ), 613 and 233 ( $\text{Ni}^{2+}$ ), 883 and 257 ( $\text{Cu}^{2+}$ ) organic ligands were selected for the modeling for the 1:1 and 1:2 complexes, respectively. For the studied complexes, the  $\log K$  values vary in the ranges of 0.8–19.1 ( $\text{Co}^{2+}$ ), 0.4–20.5 ( $\text{Ni}^{2+}$ ) and 1.2–23.0 ( $\text{Cu}^{2+}$ ). The  $\log\beta_2$  values lie in the ranges of 2.8–15.4 ( $\text{Co}^{2+}$ ), 3.0–20.6 ( $\text{Ni}^{2+}$ ) and 6.9–22.3 ( $\text{Cu}^{2+}$ ) (Fig. 1).

The datasets for three metal ions contain totally 1,237 ( $\log K$ ) and 423 ( $\log\beta_2$ ) organic ligands. The names of the ligands and experimental stability constant values are given as supporting information in SDFs. As a rule, an organic ligand has several functional groups. Studied molecules include derivatives of carboxylic and polycarboxylic acids; polyamines, (thio)ethers, various amino acids and amino-carboxylates; derivatives of phosphonous, phosphoric, (di)phosphonic and phosphinic acids; cyclic and acyclic polydentate ligands with the terminal carboxy and phosphoryl groups separated by various cyclic or acyclic spacers; various (di)sulfonic acids; ternary amines with phosphono and carboxy groups; mono- and dipodands of ternary amines; amide, (thio)phenol, imidazole, adenosine, inosine, uridine, uracil, cytidine, thymidine, adenine and guanosine derivatives; crown-ethers, thia-, and aza-crowns with neutral and acidic lariat groups, cryptands; pyridines; cyclodextrins; purine, phenanthroline, hydrazide derivatives, etc. (see supporting information: SD Files).

### Machine learning methods


Quantitative structure–property relationship modeling was performed using MLR Analysis of the ISIDA/QSPR program [53] with combined forward [40] and backward [19] stepwise variable selection techniques.

**Table 1** The log*K* values for the 1:1 (M:L) complexes of several studied ligands with Co<sup>2+</sup>, Ni<sup>2+</sup> and Cu<sup>2+</sup> in water at temperature 298 K and an ionic strength 0.1 M. The demonstration of a discrepancy in the experimental log*K* values

No.	Ligand	Metal ion	Log <i>K</i>	References
1		Co <sup>2+</sup>	4.15	[81]
			4.80	[82]
2		Co <sup>2+</sup>	10.23	[83]
			9.72	[84]
3		Co <sup>2+</sup>	4.33	[85]
			4.85	[86]
4		Co <sup>2+</sup>	3.1	[87]
			5.65	[86]
5		Co <sup>2+</sup>	5.1	[88]
			4.40	[89]
6		Ni <sup>2+</sup>	3.46	[90]
			4.43	[91]
7		Ni <sup>2+</sup>	5.49	[92]
			6.82	[93]
8		Ni <sup>2+</sup>	8.32	[94]
			10.06	[95]
9		Ni <sup>2+</sup>	2.83	[96]
			3.69	[97]
10		Ni <sup>2+</sup>	4.58	[91]
			3.27	[98]
11		Cu <sup>2+</sup>	6.67	[99]
			4.28	[100]
12		Cu <sup>2+</sup>	8.38	[101]
			7.23	[102]
13		Cu <sup>2+</sup>	3.98	[103]
			2.61	[104]

**Table 1** continued

No.	Ligand	Metal ion	LogK	References
14		$\text{Cu}^{2+}$	4.22	[87]
			6.80	[90]
15		$\text{Cu}^{2+}$	4.10	[87]
			6.96	[90]

**Fig. 1** Distribution of experimental values of the stability constants for the 1:1 (M:L) (left) and 1:2 (right) complexes of  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$ , and  $\text{Cu}^{2+}$  with organic ligands in water at temperature 298 K and an ionic strength 0.1 M

### Ensemble multiple linear regression analysis

Multiple linear regression analysis is applied to build linear relationships between independent variables (SMF descriptors:  $X_i$ ,  $i = 1, 2, \dots$ ) and a dependent variable (here target property is  $Y = \log K$  or  $\log \beta_2$ ):  $Y = c_0 + \sum c_i X_i$ , where every descriptor value (SMF count  $x_{ij}$ ,  $j = 1, 2, \dots, n$ ; here  $n$  is the number of ligands) is associated with observed property value ( $y_j$ ,  $j = 1, 2, \dots, n$ ),  $c_i$  is descriptor contribution, and  $c_0$  is the independent term which is omitted in a part of models. The singular value decomposition method [58, 59] is used to fit contributions  $c_i$  and to minimize the sum of squared residuals which are squared differences between the property values calculated by the model ( $y_{j,calc}$ ) and observed values ( $y_{j,exp}$ ) in the training set. Ensemble modeling was performed generating up to 840 MLR models each of them corresponds to particular type of the SMF descriptors and MLR equation ( $c_0 = 0$  or  $c_0 \neq 0$ ) and applied variable selection technique. Three sub-algorithms FVS-1, FVS-2 and FVS-3 for forward stepwise variable selection [40, 60] and the algorithm for backward stepwise variable selection [40, 61] have been applied. The efficiency of the FVS procedure was compared with an implementation of Genetic Algorithm (GA) on the QSPR modeling of different types of anti-HIV activities [61]. The results show similar predictive performance of computationally expensive GA-based approaches and FVS calculations. The leave-one-out (LOO) cross-validation correlation coefficient  $Q$  served as a criterion of model selection: the acceptable models were characterized by  $Q^2 > 0.5$ .

### Consensus modeling

The  $\log K$  ( $\log \beta_2$ ) values for test set compounds were predicted by consensus models (CMs). CMs combines predictions issued from great number of selected individual models originated from different types of the SMF descriptors and variable selection algorithms [19, 23, 34, 60, 62]. Thus for each compound from the test set, the target property is computed as an arithmetic mean of values obtained by individual models excluding those leading to outlying values according to Thompson's rule [63]. If a test compound is identified as being outside combined AD of individual model, the prediction by given model for a given compound is not included in CM.

### Applicability domain of models

Ensemble of three approaches for model AD has been applied: Bounding Box, Fragment Control [57] and "Quorum Control" [35]. The bounding box method considers as AD a multi-dimension descriptor space confined by minimal and maximal values of counts of SMF descriptors involved in an individual model. Fragment control rejects a prediction for a test

compound containing SMF fragments which don't occur in the initial SMF pool generated for the training set. "Quorum Control" discards a test compound if CM includes less than  $n_Q$  % of the total number of selected individual models ("no consensus without quorum"). Here  $n_Q = 15$  % was used.

### Validation of models

In order to validate CM, the external 5-fold cross validation (5-CV) was applied [21, 60]. In this procedure, an entire dataset is divided in 5 non-overlapping pairs of training and test sets. Predictions are prepared for all molecules ( $n$ ) of the initial dataset, since each of them belongs to one of the test sets. The descriptor selection and model acceptance procedures were performed only on the training folds. The compounds of test sets were not used for the model development. Predictive ability of CMs was additionally validated on the external test set of the 23 ( $M^{2+}$ )L complexes of  $Co^{2+}$ ,  $Ni^{2+}$  and  $Cu^{2+}$  with 13 new organic ligands (see Supporting information, Additional\_Test\_Set.SDF) which were not studied in the initial data sets. Predictive performance of CM has been estimated using coefficient of determination ( $R_{det}^2$ ) and RMSE for a combination of all five test sets

$$R_{det}^2 = 1 - \frac{\sum_{i=1}^n (Y_{exp,i} - Y_{pred,i})^2}{\sum_{i=1}^n (Y_{exp,i} - \langle Y \rangle_{exp})^2},$$

$$RMSE = \left[ \frac{\sum_{i=1}^n (Y_{exp,i} - Y_{pred,i})^2}{n} \right]^{1/2}$$

where  $Y_{exp}$  and  $Y_{pred}$  are, respectively, experimental and predicted values of the stability constant  $\log K$  ( $\log \beta_2$ ). The number of data points ( $n$ ), the standard deviation ( $s$ ) and the Pearson correlation coefficient ( $R_{corr}$ ) also describe linear relationships.

### Descriptors

The SMF [34, 56, 64, 65] as subgraphs of molecular graphs of the ligands were descriptors in QSPR models. Molecules were represented with implicit hydrogen atoms. Two classes of the SMF descriptors were applied: shortest topological paths with explicit representation of atoms and bonds, and terminal groups as shortest paths but defined by length and explicit identification of terminal atoms and bonds [34, 65]. Single, double and triple bonds were considered different in acyclic and cyclic non-aromatic motifs. For every class of the sequences, the minimal ( $2 \leq n_{min} \leq 4$ ) and maximal ( $6 \leq n_{max} \leq 15$ ) numbers of constituent atoms are defined. The notations  $IAB(n_{min}-n_{max})$  and  $IAB(n_{min}-n_{max})t$  represent SMF types of two classes, which include all intermediate shortest paths with  $n$  atoms:  $n_{min} \leq n \leq n_{max}$ . 60 types of the sequences of two classes have been generated varying


the values of  $n_{min}$  and  $n_{max}$ . SMF descriptors of each particular type were used as an initial descriptors' pool in QSPR modeling to build several QSPR models using different variable selection technique. Concatenated fragments always occurring in the same combination in compounds of the training set were considered as one extended fragment.

## Results and discussion

### QSPR modeling

Quantitative structure–property relationship modeling was performed for the complexation of 1,237 ( $\log K$ ) and 423


( $\log \beta_2$ ) organic ligands. 840 (360), 120 (360) and 160 (360) individual structure–property models were built for the 1:1 (1:2) complexes of  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$ , respectively, only the most stable models ( $Q^2 > 0.5$ ) were selected to include in CMs. Obtained CMs demonstrate a reasonable predictive ability of  $\log K$  in 5-CV (Fig. 2): RMSE is 0.94 ( $\text{Co}^{2+}$ ), 1.1 ( $\text{Ni}^{2+}$ ) and 1.2 ( $\text{Cu}^{2+}$ ), whereas squared determination coefficient  $R_{det}^2$  changes from 0.907 ( $\text{Cu}^{2+}$ ) to 0.929 ( $\text{Co}^{2+}$ ). RMSE of predicted  $\log \beta_2$  values varied from 1.2 to 1.4 and  $R_{det}^2$  varies from 0.780 to 0.838 as a function of metal ion (Fig. 2). Notice that similar performance for  $\log K$  predictions (RMSE = 1.1 ( $\text{Cu}^{2+}$ ) and 1.3 ( $\text{Ni}^{2+}$ ) [25]) was reported for the models built on small and structurally homogeneous datasets. On the Regression



**Fig. 2** Predicted versus experimental values of the stability constants for the 1:1 (M:L) (left) and 1:2 (right) complexes of  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$  with organic ligands in water at temperature 298 K and an ionic

strength 0.1 M. The predicted data represent a combination of the five external test sets of the 5-CV procedure





**Fig. 3** Percentage of compounds versus absolute prediction error  $|\log K_{\text{exp}} - \log K_{\text{pred}}|$  for the 1:1 (M:L) (left) and 1:2 (right) complexes of  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$ , and  $\text{Cu}^{2+}$  with organic ligands in water at 298 K and ionic strength 0.1 M.; the series of the lines (a) corresponds to CMs;

Error Curves' plots (Fig. 3), the curves corresponding to  $\log K$  lay higher than those of  $\log \beta_2$  relative to the compared lines of “no model” as arithmetic means of experimental constant values of all ligands for given metal ion (Fig. 3). Absolute prediction error of  $\log K$  is below 1.0 for 70 % ( $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$ ) and 60 % ( $\text{Cu}^{2+}$ ) of the ligands (Fig. 3). Thus, RMSE values for  $\log K$  are appreciably lower than those for  $\log \beta_2$  in line with the observed trend in experimental errors of stability constants [52, 66–69]. The RMSE values are close to observed experimental systematic errors (Table 1) showing that predictive performance of QSPR models is appropriate.

Particular types of the SMF descriptors enabled to build the individual models with high squared LOO cross-validation correlation coefficient ( $Q^2 = 0.91\text{--}0.98$ ), and low standard deviation ( $s = 0.37\text{--}1.1$ ) for all training sets of 5-CV. In most of cases, these are shortest topological paths and terminal groups containing from 2 to 12 atoms (Table 2).

The SMF types providing the best predictions in 5-CV have been used to build and validate the models on the entire modeling data set. The final CMs include the 601 (190) ( $\text{Co}^{2+}$ ), 117 (254) ( $\text{Ni}^{2+}$ ) and 160 (188) ( $\text{Cu}^{2+}$ ) individual models for  $\log K$  ( $\log \beta_2$ ). Predictive ability of CMs was validated on the additional external test set of the 23 ( $\text{M}^{2+}$ )L complexes of  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$  with 13 new organic ligands (see Supporting information, Additional\_Test\_Set.SDF) [70–76] which were not studied in the initial data sets. New ligands include acetyl phosphate, acetonylphosphonate, (2-phosphonoethoxy)ethane, 9-[2-(2-phosphonoethoxy)ethyl]adenine, 2,9-bis(hydroxymethyl)-1,10-phenanthroline, N,N'-bis(2-hydroxycyclohexyl)ethane-1,2-diamine, N,N'-bis(2-hydroxycyclohexyl)propane-1,3-diamine, 1,7-bis(2-hydroxycyclohexyl)-1,4,7-triazahexane, 1,4,8,11-tetraazacyclotetradecane-1,8-bis(methylphosphonic acid) and aminocarboxylate derivatives (Table 3). Statistical parameters of correlations (Fig. 4,  $R_{\text{det}}^2 = 0.918$  and  $\text{RMSE} = 1.97$ ) demonstrate an

the series of the lines (b) correspond to “no model”: arithmetic means of experimental constant values of all ligands for given metal ion is as the predicted value for any ligand

**Table 2** The statistical parameters of the best individual MLR models and optimal descriptor types according to the five training sets of the 5-CV procedure<sup>a</sup>

No.	SMF type <sup>b</sup>	<i>n</i>	<i>m</i>	<i>s</i>	$Q^2$
<b><math>\text{Co}^{2+}</math></b>					
1	IAB (3–10)	316–317	89–106	0.40–0.46	0.963–0.977
2	IAB (3–12)	316–317	96–109	0.37–0.44	0.960–0.977
3	IAB (2–8)	316–317	53–106	0.40–0.70	0.952–0.969
<b><math>\text{Ni}^{2+}</math></b>					
4	IAB (2–11)t	490–491	57–74	0.84–1.0	0.912–0.937
5	IAB (3–13)t	490–491	56–65	0.88–0.96	0.913–0.934
6	IAB (3–11)t	490–491	59–64	0.87–0.97	0.912–0.933
<b><math>\text{Cu}^{2+}</math></b>					
7	IAB (2–11)	706–707	71–160	0.61–1.1	0.907–0.966
8	IAB (2–12)	706–707	77–169	0.64–1.1	0.911–0.960
9	IAB (2–10)	706–707	70–161	0.64–1.1	0.906–0.964












<sup>a</sup> Statistical parameters of the MLR models: the number of data point in training set (*n*), the number of SMF variables (*m*), standard deviation (*s*), squared LOO cross-validation correlation coefficient ( $Q^2$ )

<sup>b</sup> SMF type: see the notation in Methods section: descriptors

acceptable agreement between the experimental and predicted  $\log K$  values, which is similar to that obtained in 5-CV on the modeling sets (Fig. 2).

The SMF descriptors enable detection of ligand moieties with important contributions into  $\log K$ . Thus, we detected several individual models involving common fragments whose contributions into  $\log K$  slightly vary from one model to another one. For instance, these are  $\text{O}=\text{C}-\text{C}-\text{N}_\text{c}-\text{C}_\text{c}-\text{C}_\text{c}-\text{O}_\text{c}-\text{C}_\text{c}-\text{C}_\text{c}-\text{N}_\text{c}-\text{C}-\text{C}=\text{O}$  ( $\text{Co}^{2+}$ ),  $\text{C}-\text{C}-\text{O}-\text{P}-\text{O}-\text{P}=\text{O}$  ( $\text{Co}^{2+}$ ),  $\text{N}_\text{c}-\text{C}_\text{c}-\text{C}_\text{c}-\text{N}_\text{c}-\text{C}_\text{c}-\text{C}_\text{c}-\text{O}$  ( $\text{Ni}^{2+}$ ),  $\text{C}_\text{ar}-\text{N}_\text{ar}-\text{C}_\text{ar}-\text{C}_\text{ar}-\text{N}_\text{ar}-\text{C}_\text{ar}$  ( $\text{Ni}^{2+}$ ),  $\text{O}-\text{N}-\text{C}=\text{O}$  ( $\text{Cu}^{2+}$ ),  $\text{O}-\text{N}=\text{C}-\text{C}=\text{O}$  ( $\text{Cu}^{2+}$ ) fragments which are involved in 13–126 models and contribute from 2.1 to 6.0  $\log K$  units into the stability (see Table 4). These fragments occur in derivatives of 1,7-di- and 1,4,7-triazacyclododecane with acetate pendant arms



**Table 3** Experimental and predicted stability constant values  $\log K$  of the 1:1 (M:L) complexes of  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$  for the additional test set<sup>a</sup>

No.	Ligand	Cation	LogK		$N_m^c$
			<i>exp.</i>	<i>pred.</i> <sup>b</sup>	
1		$\text{Co}^{2+}$	1.83 <sup>d</sup>	3.18 (0.29)	110
2		$\text{Co}^{2+}$	2.19 <sup>d</sup>	2.46 (0.80)	256
3		$\text{Co}^{2+}$	2.24 <sup>e</sup>	1.85 (0.89)	214
4		$\text{Co}^{2+}$	2.21 <sup>e</sup>	5.32 (0.71)	168
5		$\text{Co}^{2+}$	6.36 <sup>f</sup>	4.31 (0.74)	206
6		$\text{Ni}^{2+}$	1.75 <sup>d</sup>	4.1 (1.2)	54
7		$\text{Ni}^{2+}$	2.14 <sup>d</sup>	3.52 (0.53)	34
8		$\text{Ni}^{2+}$	2.18 <sup>e</sup>	2.51 (0.36)	38
9		$\text{Ni}^{2+}$	2.41 <sup>e</sup>	4.34 (0.49)	31
10		$\text{Ni}^{2+}$	7.42 <sup>f</sup>	7.46 (0.58)	51
11		$\text{Ni}^{2+}$	7.77 <sup>g</sup>	8.42 (0.83)	44



No.	Ligand	Cation	LogK		
			<i>exp.</i>	<i>pred.</i> <sup>b</sup>	<i>N<sub>m</sub></i> <sup>c</sup>
12		Cu <sup>2+</sup>	2.86 <sup>d</sup>	5.36 (0.65)	54
13		Cu <sup>2+</sup>	3.36 <sup>d</sup>	4.89 (0.79)	59
14		Cu <sup>2+</sup>	3.44 <sup>e</sup>	3.00 (0.22)	40
15		Cu <sup>2+</sup>	3.98 <sup>e</sup>	4.60 (0.31)	37
16		Cu <sup>2+</sup>	21.43 <sup>h</sup>	18.54 (0.92)	122
17		Cu <sup>2+</sup>	19.47 <sup>h</sup>	17.63 (0.77)	77
18		Cu <sup>2+</sup>	24.9 <sup>i</sup>	22.7 (1.4)	43
19		Cu <sup>2+</sup>	11.47 <sup>g</sup>	9.8 (1.7)	68
20		Cu <sup>2+</sup>	12.67 <sup>g</sup>	11.7 (1.4)	67
21		Cu <sup>2+</sup>	16.74 <sup>g</sup>	12.2 (1.5)	60

**Table 3** continued

No.	Ligand	Cation	LogK		$N_m^c$
			exp.	pred. <sup>b</sup>	
22		$\text{Cu}^{2+}$	5.13 <sup>j</sup>	7.66 (0.91)	60
23		$\text{Cu}^{2+}$	5.64 <sup>j</sup>	7.78 (0.52)	65
$R_{det}^2$ 0.918					
RMSE 1.97					

<sup>a</sup> Experimental data are given at 298 K and an ionic strength ( $I$ ) 0.1 M excepting: for the ligand 18  $I = 1.0$  M, for the ligands 22 and 23  $I = 0.2$  M

<sup>b</sup> Predicted stability constant values  $\log K_{pred}$  are computed by the MLR CMs, standard deviations are given in parentheses

<sup>c</sup> The number of individual models in CM using AD for the given ligand

<sup>d</sup> Ref. [70]

<sup>e</sup> Ref. [71]


<sup>f</sup> Ref. [72]

<sup>g</sup> Ref. [73]

<sup>h</sup> Ref. [74]

<sup>i</sup> Ref. [75]

<sup>j</sup> Ref. [76]



**Fig. 4** Experimental versus predicted stability constant values  $\log K$  of the 1:1 (M:L) complexation for the additional external test set

( $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$ ), diphosphoric acid ( $\text{Co}^{2+}$ ), 1,10-phenanthroline ( $\text{Ni}^{2+}$ ), hydroxamic acid and 2-hydroxyiminopropanamide ( $\text{Cu}^{2+}$ ) (Table 4). As we earlier demonstrated [34,

62, 77], SMF and their contributions could become convenient building blocks for compounds' design.

It should be noticed that developed in this work QSPR models outperform previously reported ones ( $\text{Co}^{2+}$  [20, 47, 48],  $\text{Ni}^{2+}$  [20, 25, 47, 48],  $\text{Cu}^{2+}$  [20, 25, 47–49]) in respect to both diversity and the size of the ligand sets, and rigorous cross-validation procedure.










#### Relationships between stability constants

Analysis of available experimental data allowed us to detect several linear correlations between experimental  $\log K$  values for one same ligand with different metal ions and between  $\log K$  and  $\log \beta_2$  for one same metal ion.

For known ligand, the stability constant  $\log K$  for one metal ion can be predicted by means of the stability constant for a second metal ion using the equations:

$$\log K_{\text{CoL}} = 0.867 \log K_{\text{NiL}}, \quad (1)$$


**Table 4** Selected SMF and their mean contributions into  $\log K$  according to the sets of individual MLR models<sup>a</sup>

No.	SMF <sup>b</sup>	$\langle a_i \rangle$	$N_{model}$	$N_{mol}$	Ligand <sup>c</sup>
Co <sup>2+</sup> + L, 396 ligands					
1	O=C–C–N <sub>c</sub> –C <sub>c</sub> –C <sub>c</sub> –O <sub>c</sub> –C <sub>c</sub> –C <sub>c</sub> –N <sub>c</sub> –C–C=O	6.05 (0.35)	30	3	
2	O–N–C=O	2.11 (0.41)	126	11	
3	C–C–O–P–O–P=O	2.16 (0.19)	39	10	
Ni <sup>2+</sup> + L, 613 ligands					
4	N <sub>c</sub> –C <sub>c</sub> –C <sub>c</sub> –N <sub>c</sub> –C <sub>c</sub> –C <sub>c</sub> –O <sub>c</sub>	3.36 (0.69)	35	13	
5	C <sub>ar</sub> –N <sub>ar</sub> –C <sub>ar</sub> –C <sub>ar</sub> –N <sub>ar</sub> –C <sub>ar</sub>	4.65 (0.56)	25	4	
6	O–N–C=O	2.67 (0.14)	13	12	
Cu <sup>2+</sup> + L, 883 ligands					
7	O–N–C=O	6.50 (0.60)	54	7	
8	O–N=C–C=O	5.63 (0.61)	35	5	
9	C–N–C–C=N–O	3.62 (0.45)	35	6	

<sup>a</sup>  $\langle a_i \rangle$  is fragment contribution (arithmetic mean) and its standard deviation (in parentheses) according to the  $N_{model}$  individual models and the  $N_{mol}$  ligands

<sup>b</sup> Substructural molecular fragments (SMF): shortest topological paths with explicit representation of atoms and bonds; bonds: '–' and '=' are single and double in chain, N<sub>c</sub>–C<sub>c</sub> is single bond in non-aromatic cycle; C<sub>ar</sub>–N<sub>ar</sub> is aromatic bond

<sup>c</sup> Example of ligand including corresponding SMF which bonds are accented by thickness




**Fig. 5** The linear relationships  $\log K_{CoL}$  versus  $\log K_{NiL}$  (a),  $\log K_{CoL}$  versus  $\log K_{CuL}$  (b) and  $\log K_{NiL}$  versus  $\log K_{CuL}$  (c) for the complexation  $M^{2+} + L = (M^{2+})L$  of the  $Co^{2+}$ ,  $Ni^{2+}$  and  $Cu^{2+}$  ( $M^{2+}$ ) ions with diverse organic ligands (L) in water at 298 K and an ionic strength 0.1 M

$$\log K_{CoL} = 0.618 \log K_{CuL}, \quad (2)$$

$$\log K_{NiL} = 0.731 \log K_{CuL}. \quad (3)$$

These linear relationships were derived using the  $\log K$  values for 272, 221 and 339 ligands, respectively (Fig. 5). The standard deviations of the  $\log K$  estimations vary from 0.7 to 1.1 (Fig. 5). For given ligand, the order of stability of complexes is  $(Co^{2+})L < (Ni^{2+})L < (Cu^{2+})L$ . The  $Cu^{2+}$  cation forms most stable complexes compared to



**Fig. 6**  $\log \beta_2$  versus  $\log K$  for the complexations of the  $Co^{2+}$ ,  $Ni^{2+}$  and  $Cu^{2+}$  ions with organic ligands in water at 298 K and an ionic strength 0.1 M

those of  $Co^{2+}$  and  $Ni^{2+}$ . The coefficient of Eq. 1 is more close to one than those of Eqs. 2 and 3 indicate larger chemical similarity between  $Co^{2+}$  and  $Ni^{2+}$  than that for other metal pairs.

Figure 6 shows, that a linear relationship between  $\log \beta_2$  and  $\log K$  is observed for every metal ion. The squared correlation coefficient varies from 0.826 ( $Cu^{2+}$ ) to 0.954 ( $Co^{2+}$ ), and the standard deviation changes from 0.63 ( $Co^{2+}$ ) to 1.26 ( $Cu^{2+}$ ). The relationships (Fig. 6) give possibility to estimate  $\log \beta_2$  ( $\log K$ ) from the knowledge of  $\log K$  ( $\log \beta_2$ ) for given ligand. For all ligands forming both

**Table 5** Predicted stability constant values  $\log K$  of the 1:1 (M:L) complexes of  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$  with the ligands which can form metallosupramolecular grid complexes in water

No.	Ligand <sup>a</sup>	LogK		
		( $\text{Co}^{2+}$ )L	( $\text{Ni}^{2+}$ )L	( $\text{Cu}^{2+}$ )L
1		9.5	10.6	12.6
2		9.6	11.1	14.4
3		7.5	8.8	10.0

<sup>a</sup> The ligands are **33**, **28** and **11** from Ref. [44]

1:1 and 1:2 complexes with the  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$  ions, the following ratio is observed:

$$\log\beta_2/\log K = 1.8 \pm 0.1; n = 492. \quad (4)$$

Thereby, stepwise stability constant  $\log K_2 = \log\beta_2 - \log K$  constitutes 80 % of  $\log K$ , and the free energy of coordination of second ligand molecule is weaker by 20 % in comparison with coordination of first ligand molecule to metal ion.

Linear correlation between stability constants of ML and  $\text{ML}_2$  complexes may have an interesting interpretation in the context of metal-containing supramolecular architectures (grids). Thus, one may suggest that L forms stable metallogrid complexes  $\text{M}_n\text{L}_n$ , if stability of its 1:1 complex is large enough. To illustrate this suggestion, developed QSPR models were applied to assess  $\log K$  for ( $\text{Co}^{2+}$ )L, ( $\text{Ni}^{2+}$ )L and ( $\text{Cu}^{2+}$ )L complexes of the 44 ligands [44] which form metallogrid complexes (see their structures in Supporting information: Metallogrid\_forming\_ligands.SDF). Three ligands forming most stable ML complexes are shown in Table 5. One may suggest that they, indeed, may form metallogrid complexes with  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$  in water. There are some experimental proves of these suggestion: the ( $\text{Co}^{2+}$ )<sub>4</sub>L<sub>4</sub> complex of ligand 1 (Table 5) has been formed at the air-aqueous solution interface [78], and grid complex of  $\text{Co}^{2+}$  with ligand 2 was synthesized in the  $\text{CD}_3\text{CN}$  and  $\text{D}_2\text{O}$  solution [79]. For ligand 3, the synthesis of the ( $\text{Zn}^{2+}$ )<sub>4</sub>L<sub>4</sub> complex in MeCN was reported [80].

## Conclusions

This article reports predictive QSPR models for the stability constants  $\log K$  and  $\log\beta_2$  of the 1:1 and 1:2 (M:L) complexes of transition metal ions ( $\text{M} = \text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{Cu}^{2+}$ ) with different classes of organic ligands (L) in aqueous solution at 298 K and an ionic strength 0.1 M. The models have been built using ensemble MLR and SMF descriptors on big data sets of 396 ( $\text{Co}^{2+}$ ), 613 ( $\text{Ni}^{2+}$ ), 883 ( $\text{Cu}^{2+}$ ) ligands for  $\log K$  and 132 ( $\text{Co}^{2+}$ ), 233 ( $\text{Ni}^{2+}$ ), 257 ( $\text{Cu}^{2+}$ ) ligands for  $\log\beta_2$ . They have reasonable prediction performance: RMSE is 0.94–1.2 ( $\log K$ ) and 1.2–1.4 ( $\log\beta_2$ ) which is close to observed experimental systematic errors. The  $\log K$  and  $\log\beta_2$  values were predicted by CMs as arithmetic means of several hundreds of individual models taking into account their combined ADs. The SMFs enable detection of ligand moieties with important contributions into stability constants and they can be used as building blocks of new ligands. Detected relationships between stability constants for pair of metal ions demonstrate that the order of stability of complexes is ( $\text{Co}^{2+}$ )L < ( $\text{Ni}^{2+}$ )L < ( $\text{Cu}^{2+}$ )L, and the  $\log K_{\text{CoL}}$  and  $\log K_{\text{NiL}}$  values average 62 and 73 % of the  $\log K_{\text{CuL}}$  value, respectively. For all ligands forming both 1:1 and 1:2 complexes the following ratio is observed:  $\log\beta_2/\log K = 1.8 \pm 0.1$ ,  $n = 492$ . We believe that this relationship could be useful in design of new ligands for metallosupramolecular complexes.

**Acknowledgments** The authors thank Profs. G. Pettit and L. Pettit from Academic Software for providing with the SCDB-to-SDF program.

## References

- McCleverty JA, Meyer TJ (eds) (2003) Comprehensive coordination chemistry II: from Biology to Nanotechnology. Applications of coordination chemistry, vol. 9. Elsevier, Amsterdam
- Duca G (2012) Homogeneous catalysis with metal complexes: fundamentals and applications. Springer, Berlin
- Kumar S, Dhar DN, Saxena PN (2009) Applications of metal complexes of Schiff bases—a review. *J Sci Ind Res* 68(March): 181–187
- Schühle DT, Peters JA, Schatz J (2011) Metal binding calixarenes with potential biomimetic and biomedical applications. *Coord Chem Rev* 255:2727–2745
- Mewis RE, Archibald SJ (2010) Biomedical applications of macrocyclic ligand complexes. *Coord Chem Rev* 254(15–16):1686–1712
- Delgado R, Felix V, Lima LMP, Price DW (2007) Metal complexes of cyclen and cyclam derivatives useful for medical applications: a discussion based on thermodynamic stability constants and structural data. *Dalton Trans* 26:2734–2745
- Anderegg G, Arnaud-Neu F, Delgado R, Felcman J, Popov K (2005) Critical evaluation of stability constants of metal complexes of complexones for biomedical and environmental applications. *Pure Appl Chem* 77(8):1445–1495
- Que LJ, Tolman WB (eds) (2003) Comprehensive Coordination Chemistry II. Bio-coordination Chemistry, vol 8. Elsevier, San Diego
- Bhattacharya PK (2005) Metal ions in biochemistry. Alpha Science International, Harrow
- Sigel A, Sigel H (eds) (2000) Metal ions in biological systems. Manganese and its role in biological processes, vol 37. CRC Press, New York
- Tretyakov YD, Martynenko LI, Grigoryev AN, Tsivadze AY (2001) Inorganic chemistry. Chemistry of elements. Book 1 (Rus). Himia, Moscow
- Hancock RD (1997) Approaches to predicting stability constants: a critical review. *Analyst* 122(4):51R–58R
- Hancock RD, Martell AE (1989) Ligand design for selective complexation of metal ions in aqueous solution. *Chem Rev* 89(8):1875–1914
- Martell AE, Hancock RD, Motekaitis RJ (1994) Factors affecting stabilities of chelate, macrocyclic and macrobicyclic complexes in solution. *Coord Chem Rev* 133(JUL):39–65
- Dimmock PW, Warwick P, Robbins RA (1995) Approaches to predicting stability constants. *Analyst* 120(8):2159–2170
- Popov KI, Wanner H (2005) Stability constants data sources: critical evaluation and application for environmental speciation. In: Nowack B, VanBriesen JM (eds) Biogeochemistry of chelating agents. ACS Symposium Series 910. American Chemical Society, Washington, pp 50–73
- Pattammattel A, Deshapriya IK, Chowdhury R, Kumar CV (2013) Metal-enzyme frameworks: role of metal ions in promoting enzyme self-assembly on  $\alpha$ -zirconium(IV) phosphate nanoplates. *Langmuir* 29:2971–2981
- Daniele PG, Foti C, Gianguzza A, Prenesti E, Sammartano S (2008) Weak alkali and alkaline earth metal complexes of low molecular weight ligands in aqueous solution. *Coord Chem Rev* 252(1011):1093–1107
- Solov'ev VP, Varnek AA (2004) Structure–property modeling of metal binders using molecular fragments. *Russ Chem Bull* 53(7):1434–1445
- Toropov AA, Toropova AP, Nesterova AI, Nabiev OM (2004) QSPR modeling of complex stability by correlation weighing of the topological and chemical invariants of molecular graphs. *Russ J Coord Chem* 30(9):611–617
- Tetko IV, Solov'ev VP, Antonov AV, Yao XJ, Fan BT, Hoonakker F, Fourches D, Lachiche N, Varnek A (2006) Benchmarking of linear and non-linear approaches for quantitative structure–property relationship studies of metal complexation with organic ligands. *J Chem Inf Model* 46(2):808–819
- Svetlitski R, Lomaka A, Karelson M (2006) QSPR modelling of lanthanide-organic complex stability constants. *Sep Sci Technol* 41(1):197–216
- Solov'ev VP, Kireeva NV, Tsivadze AY, Varnek AA (2006) Structure–property modelling of complex formation of strontium with organic ligands in water. *J Struct Chem* 47(2):298–311
- Varnek A, Fourches D, Kireeva N, Klimchuk O, Marcou G, Tsivadze A, Solov'ev V (2008) Computer-aided design of new metal binders. *Radiochim Acta* 96(8):505–511
- Cabaniss SE (2008) Quantitative structure–property relationships for predicting metal binding by organic ligands. *Environ Sci Technol* 42(14):5210–5216
- X-y Wang, H-l Wang (2008) QSPR study of crown ether complexes with K<sup>+</sup> by density functional theory. *Xiangtan Daxue Ziran Kexue Xuebao* 30(3):86–93
- Ghasemi J, Saaidpour S (2008) QSPR modeling of stability constants of diverse 15-crown-5 ethers complexes using best multiple linear regression. *J Incl Phenom Macrocycl Chem* 60(3–4):339–351
- Raos N, Miličević A (2009) Estimation of stability constants of coordination compounds using models based on topological indices. *Arch Ind Hyg Toxicol* 60(1):123–128
- Ghasemi JB, Ahmadi S, Ayati M (2010) QSPR modeling of stability constants of the Li-hemispherands complexes using MLR: a theoretical host-guest study. *Macrocycles* 3(4):234–242
- Ghasemi JB, Rofouei MK, Salahinejad M (2011) A quantitative structure–property relationships study of the stability constant of crown ethers by molecular modelling: new descriptors for lariat effect. *J Incl Phenom Macrocycl Chem* 70:37–47
- Li Y, Su L, Zhang X, Huang X, Zhai H (2011) Prediction of association constants of cesium chelates based on uniform design optimized support vector machine. *Chemom Intell Lab Syst* 105(1):106–113
- Mousavi A (2011) Predicting mercury(II) binding by organic ligands: a chemical model of therapeutic and environmental interests. *Environ Forensic* 12(4):327–332
- Garkani-Nejad Z, Ahmadvand M (2011) Simultaneous estimation of stability constants of Mg, Ba, Ca, and Sr complexes using a small subset of molecular descriptors. *J Coord Chem* 64(14):2466–2479
- Solov'ev V, Sukhno I, Buzko V, Polushin A, Marcou G, Tsivadze A, Varnek A (2012) Stability constants of complexes of Zn<sup>2+</sup>, Cd<sup>2+</sup>, and Hg<sup>2+</sup> with organic ligands: QSPR consensus modeling and design of new metal binders. *J Incl Phenom Macrocycl Chem* 72(3–4):309–321
- Solov'ev VP, Tsivadze AY, Varnek AA (2012) New approach for accurate QSPR modeling of metal complexation: application to stability constants of complexes of lanthanide ions Ln<sup>3+</sup>, Ag<sup>+</sup>, Zn<sup>2+</sup>, Cd<sup>2+</sup> and Hg<sup>2+</sup> with organic ligands in water. *Macrocycles* 5(4–5):404–410
- Solov'ev V, Marcou G, Tsivadze AY, Varnek A (2012) Complexation of Mn<sup>2+</sup>, Fe<sup>2+</sup>, Y<sup>3+</sup>, La<sup>3+</sup>, Pb<sup>2+</sup>, and UO<sub>2</sub><sup>2+</sup> with organic ligands: QSPR ensemble modeling of stability constants. *Ind Eng Chem Res* 51(41):13482–13489
- Ahmadi S (2012) Application of GA-MLR method in QSPR modeling of stability constants of diverse 15-crown-5 complexes with sodium cation. *J Incl Phenom Macrocycl Chem* 74(1–4):57–66

38. Daraei H, Irandoust M, Ghasemi JB, Kurdian AR (2012) QSPR probing of Na<sup>+</sup> complexation with 15-crown-5 ethers derivatives using artificial neural network and multiple linear regression. *J Incl Phenom Macrocycl Chem* 72(3–4):423–435
39. Mousavi A (2013) A chemical model for predicting zinc(II) binding by organic ligands as hypothetical therapeutic agents. *Med Chem Res* 22(1):234–239
40. Solov'ev VP, Kireeva N, Tsivadze AY, Varnek A (2013) QSPR ensemble modelling of alkaline-earth metal complexation. *J Incl Phenom Macrocycl Chem* 76(1–2):159–171
41. Varnek A, Solov'ev V (2009) Quantitative structure–property relationships in solvent extraction and complexation of metals. In: Sengupta AK, Moyer BA (eds) *Ion exchange and solvent extraction, a series of advances*, vol 19., CRC Press/Taylor and Francis Group, Boca Raton, pp 319–358
42. Ghasemi JB, Salahinejad M, Rofouei MK (2011) Review of the quantitative structure–activity relationship modelling methods on estimation of formation constants of macrocyclic compounds with different guest molecules. *Supramol Chem* 23(9):615–631
43. Buist D, Williams NJ, Reibenspies JH, Hancock RD (2010) Control of metal ion size-based selectivity through chelate ring geometry. metal ion complexing properties of 2,2'-biimidazole. *Inorg Chem* 49(11):5033–5039
44. Hardy JG (2013) Metallosupramolecular grid complexes: towards nanostructured materials with high-tech applications. *Chem Soc Rev* 42:7881–7899
45. Schneider H-J (2009) Binding mechanisms in supramolecular complexes. *Angew Chem Int Ed* 48:3924–3977
46. Carolan AN, Mroz AE, El Ojaimi M, VanDerveer DG, Thummel RP, Hancock RD (2012) Metal-ion-complexing properties of 2-(pyrid-2'-yl)-1,10-phenanthroline, a more preorganized analogue of terpyridyl. A crystallographic, fluorescence, and thermodynamic study. *Inorg Chem* 51(5):3007–3015
47. Toropov AA, Toropova AP (2001) QSPR modeling of stability of complexes of adenosine phosphate derivatives with metals absent from the complexes of the teaching access. *Russ J Coord Chem* 27(8):574–578
48. Toropov AA, Toropova AP (2002) QSPR modeling of complex stability by optimization of correlation weights of the hydrogen bond index and the local graph invariants. *Russ J Coord Chem* 28(12):877–880
49. Grgas B, Nikolić S, Paulić N, Raos N (1999) Estimation of stability constants of copper(II) chelates with N-alkylated amino acids using topological indices. *Croat Chem Acta* 72(4):885–895
50. Gorden AEV, Xu J, Raymond KN, Durbin P (2003) Rational design of sequestering agents for plutonium and other actinides. *Chem Rev* 103(11):4207–4282
51. Bianchi A, Calabi L, Corana F, Fontana S, Losi P, Maiocchi A, Paleari L, Valtancoli B (2000) Thermodynamic and structural properties of Gd(III) complexes with polyamino-polycarboxylic ligands: basic compounds for the development of MRI contrast agents. *Coord Chem Rev* 204:309–393
52. IUPAC Stability Constants Database (2012) Academic Software. <http://www.acadsoft.co.uk/>. Accessed 26 Apr 2013
53. Solov'ev VP, Varnek AA (2008–2013) ISIDA (In silico design and data analysis) program. <http://infochim.u-strasbg.fr/spip.php?rubrique53> or <http://vpsolovev.ru/programs/>. Accessed 12 Dec 2013
54. Solov'ev VP, Varnek AA (1999–2013) EdChemS (Editor of chemical structures). <http://infochim.u-strasbg.fr/spip.php?rubrique51> or <http://vpsolovev.ru/programs/>. Accessed 12 Dec 2013
55. Solov'ev VP, Varnek AA (2013) EdiSDF (Editor of Structure—Data Files). <http://infochim.u-strasbg.fr/spip.php?rubrique52> or <http://vpsolovev.ru/programs/>. Accessed 12 Dec 2013
56. Varnek A, Fourches D, Hoonakker F, Solov'ev VP (2005) Substructural fragments: an universal language to encode reactions, molecular and supramolecular structures. *J Comput Aided Mol Des* 19(9–10):693–703
57. Varnek A, Fourches D, Horvath D, Klimchuk O, Gaudin C, Vayer P, Solov'ev V, Hoonakker F, Tetko IV, Marcou G (2008) ISIDA-platform for virtual screening based on fragment and pharmacophoric descriptors. *Curr Comput Aided Drug Des* 4(3):191–198
58. Lawson CL, Hanson RJ (1974) *Solving least squares problems*. Prentice Hall, Englewood Cliffs
59. Forsythe GE, Malcolm MA, Moler CB (1977) *Computer methods for mathematical computations*. Prentice Hall, Englewood Cliffs
60. Varnek A, Kireeva N, Tetko IV, Baskin II, Solov'ev VP (2007) Exhaustive QSPR studies of a large diverse set of ionic liquids: how accurately can we predict melting points? *J Chem Inf Model* 47(3):1111–1122
61. Horvath D, Bonachera F, Solov'ev V, Gaudin C, Varnek A (2007) Stochastic versus stepwise strategies for quantitative structure–activity relationship generation—how much effort may the mining for successful QSAR models take? *J Chem Inf Model* 47(3):927–939
62. Varnek A, Solov'ev VP (2005) “In silico” design of potential anti-HIV actives using fragment descriptors. *Comb Chem High Throughput Screen* 8(5):403–416
63. Muller PH, Neumann P, Storm R (1979) *Tafeln der mathematischen Statistik*. VEB Fachbuchverlag, Leipzig
64. Solov'ev VP, Varnek AA, Wipff G (2000) Modeling of ion complexation and extraction using substructural molecular fragments. *J Chem Inf Comput Sci* 40(3):847–858
65. Solov'ev V, Oprisiu I, Marcou G, Varnek A (2011) Quantitative structure–property relationship (QSPR) modeling of normal boiling point temperature and composition of binary azeotropes. *Ind Eng Chem Res* 50(24):14162–14167
66. Martell AE, Smith RM (1989) *Critical stability constants*, vol 1–6. Plenum Press, New York
67. Christensen JJ, Izatt RM (1983) *Handbook of metal ligand heats and related thermodynamic quantities*. Marcel Dekker Inc., New York
68. Izatt RM, Pawlak K, Bradshaw JS, Bruening RL (1991) Thermodynamic and kinetic data for macrocycle interaction with cations and anions. *Chem Rev* 91(8):1721–2085
69. Solov'ev VP, Vnuk EA, Strakhova NN, Raevsky OA (1991) Thermodynamics of complexation of the macrocyclic polyethers with salts of alkali and alkaline-earth metals (Rus.). VINITI, Moscow
70. Sigel H, DaCosta CP, Song B, Carloni P, Gregan F (1999) Stability and structure of metal ion complexes formed in solution with acetyl phosphate and acetonylphosphonate: quantification of isomeric equilibria. *J Am Chem Soc* 121(26):6248–6257
71. Fernandez-Botello A, Griesser R, Holy A, Moreno V, Sigel H (2005) Acid-base and metal-ion-binding properties of 9-[2-(2-Phosphonoethoxy)ethyl]adenine (PEEA), a relative of the antiviral nucleotide analogue 9-[2-(Phosphonomethoxy)ethyl]adenine (PMEA). An exercise on the quantification of isomeric complex equilibria in solution. *Inorg Chem* 44(14):5104–5117
72. Gephart RT III, Williams NJ, Reibenspies JH, De Sousa AS, Hancock RD (2008) Metal ion complexing properties of the highly preorganized LIGAND 2,9-bis(Hydroxymethyl)-1,10-phenanthroline: a crystallographic and thermodynamic study. *Inorg Chem* 47(22):10342–10348
73. Hancock RD, de Sousa AS, Walton GB, Reibenspies JH (2007) Metal-ion selectivity produced by C-alkyl substituents on the bridges of chelating ligands: the importance of short H–H nonbonded van der Waals Contacts in controlling metal-ion selectivity. A Thermodynamic, molecular mechanics, and crystallographic study. *Inorg Chem* 46(11):4749–4757



74. Koteck J, Kálmán FK, Hermann P, Brücher E, Binnemans K, Lukeš I (2006) Study of thermodynamic and kinetic stability of transition metal and lanthanide complexes of DTPA analogues with a phosphorus acid pendant arm. *Eur J Inorg Chem* 2006(10):1976–1986
75. Kálmán FK, Baranyai Z, Tóth I, Bányai I, Király R, Brücher E, Aime S, Sun X, Sherry AD, Kovács Z (2008) Synthesis, potentiometric, kinetic, and NMR studies of 1,4,7,10-tetraazacyclododecane-1,7-bis(acetic acid)-4,10-bis(methylenephosphonic acid) (DO2A2P) and its complexes with Ca(II), Cu(II), Zn(II) and lanthanide(III) Ions. *Inorg Chem* 47(9):3851–3862
76. Nagy NV, Van Doorslaer S, Szabó-Plánka T, Van Rompaey S, Hamza A, Fülöp F, Tóth GK, Rockenbauer A (2012) Copper(II)-binding ability of stereoisomeric cis- and trans-2-aminocyclohexanecarboxylic acid-L-phenylalanine dipeptides. A combined CW/pulsed EPR and DFT study. *Inorg Chem* 51(3):1386–1399
77. Varnek A, Fourches D, Solov'ev VP, Baulin VE, Turanov AN, Karandashev VK, Fara D, Katritzky AR (2004) "In silico" Design of new uranyl extractants based on phosphoryl-containing podands: QSPR Studies, generation and screening of virtual combinatorial library, and experimental tests. *J Chem Inf Comput Sci* 44(4):1365–1382
78. Weissbuch I, Baxter PNW, Cohen S, Cohen H, Kjær K, Howes PB, Als-Nielsen J, Hanan GS, Schubert US, Lehn JM, Leiserowitz L, Lahav M (1998) Self-assembly at the air-water interface. In-situ preparation of thin films of metal ion grid architectures. *J Am Chem Soc* 120(19):4850–4860
79. Breuning E, Ziener U, Lehn J-M, Wegelius E, Rissanen K (2001) Two-level self-organisation of arrays of [2 × 2] grid-type tetranuclear metal complexes by hydrogen bonding. *Eur J Inorg Chem* 6:1515–1521
80. Bark T, Düggeli M, Stoeckli-Evans H, von Zelewsky A (2001) Designed molecules for self-assembly: the controlled formation of two chiral self-assembled polynuclear species with predetermined configuration. *Angew Chem Int Ed* 40(15):2848–2851
81. Israeli M, Pettit LD (1975) Complex formation between unsaturated  $\alpha$ -aminoacids and silver(I) and some divalent transition metal ions. *J Inorg Nucl Chem* 37(4):999–1003
82. Khalil MM, Attia AE (2000) Potentiometric studies on the formation equilibria of binary and ternary complexes of some metal ions with dipicolinic acid and amino acids. *J Chem Eng Data* 45(6):1108–1111
83. Motekaitis RJ, Murase I, Martell AE (1971) New multidentate ligands—XI: synthesis and chelating tendencies of ethylenediamine-N, N'-di(methylenephosphonic) acid, ethylenediamine-N, N', N'-tetra(methylenephosphonic) acid and ethylenediamine-N, N'-di(methylenephosphonic) acid. *J Inorg Nucl Chem* 33(10):3353–3365
84. Markhaeva VP, Nikolaeva LS, Ditzel F, Baier L, Parshikova EA, Golskikh VA, Bodoev NV (2001) The study of the complexation of 1,2-diaminoethane-N, N'-bis(methylenephosphonic) acid with Ni<sup>2+</sup> and Co<sup>2+</sup> cations. *Zh Neorg Khim (Rus)* 46(1):85–91
85. Sovago I, Kiss T, Gergely A (1993) Critical survey of the stability constants of complexes of aliphatic amino acids. *Pure Appl Chem* 65(5):1029–1080
86. Ismail NM (1997) Potentiometric studies on ternary metal complexes of some aliphatic acids and aminoacids. *J Indian Chem Soc* 74(5):396–398
87. Boraie AAA, Mohamed NFA (2002) Equilibrium studies of ternary systems involving divalent transition metal ions, aliphatic acids, and triazoles. *J Chem Eng Data* 47(4):987–991
88. Smith RM, Martell AE, Chen Y (1991) Critical evaluation of stability constants for nucleotide complexes with protons and metal ions and the accompanying enthalpy changes. *Pure Appl Chem* 63(7):1015–1080
89. Boraie AAA, Taha F, Mohamed AH, Ibrahim SA (2001) Medium effect and thermodynamic studies for the proton-ligand and metal-ligand formation constants of the ternary systems MII+adenosine-5'-triphosphate (ATP)+ asparagine. *J Chem Eng Data* 46(2):267–275
90. Khalil MM, Radalla AM (1998) Binary and ternary complexes of inosine. *Talanta* 46(1):53–61
91. Azab HA, Hassan A, El-Nady AM, Azkal RSA (1993) Ternary complexes of nickel(II) with AMP, ADP and ATP as primary ligands and some biologically important polybasic oxygen acids as secondary ligands. *Monatsh Chem* 124(3):267–276
92. Azab HA, Anwar ZM, Sokar M (2004) Metal ion complexes containing nucleobases and some zwitterionic buffers. *J Chem Eng Data* 49(1):62–72
93. Khalil MM, Fazary AE (2004) Potentiometric studies on binary and ternary complexes of di- and trivalent metal ions involving some hydroxamic acids, amino acids, and nucleic acid components. *Monatsh Chem* 135(12):1455–1474
94. Mulla F, Marsicano F, Nakani BS, Hancock RD (1985) Stability of ammonia complexes that are unstable to hydrolysis in water. *Inorg Chem* 24(19):3076–3080
95. Zhadanov BV, Polyakova IA, Tsirul'nikova NV, Sushitskaya TM, Temkina VY (1979) Study of acid dissociation and complexing properties of imino-N, N-bis(methylenephosphonic acid). *Koord Khim (Rus)* 5:1614–1619
96. Patel RN, Shrivastava RP, Singh N, Pandeya KB (2001) Equilibrium study on the mixed ligand mixed metal complex formation stability of copper(II), nickel(II) and zinc(II) with glycylvaline and imidazole. *Indian J Chem* 40A(4):361–367
97. Mukherjee GN, Sahu HK (1998) Multimetal multiligand complexes. Part I. Equilibrium study on the formation and stability of mixed ligand mixed metal complexes of cobalt-, nickel-, copper- and zinc(II) with aspartate and imidazole in aqueous solution. *J Indian Chem Soc* 75(3):143–147
98. Powell JE, Johnson DK (1969) Stability trends of some 1:1 and 2:1 malonato and 1,1-cyclobutanedicarboxylato cobalt, nickel, copper and zinc chelates. *J Chromatogr* 44:212–213
99. Brookes G, Pettit LD (1977) Complex formation and stereoselectivity in the ternary systems copper(II)-D/L-histidine-L-amino-acids. *J Chem Soc Dalton Trans* 19:1918–1924
100. Venkatnarayana G, Swamy S, Lingaiah P (1988) Ternary complexes of copper(II) with malonic acid and O, O; O, N and N, N donor ligands. *Indian J Chem* 27A(7):613–616
101. Shoukry MM, Khairy EM, El-Sherif AA (2002) Ternary complexes involving copper(II) and amino acids, peptides and DNA constituents. The kinetics of hydrolysis of  $\alpha$ -amino acid esters. *Transition Met Chem* 27(6):656–664
102. Anwar ZM, Azab HA (1999) Ternary complexes in solution. Comparison of the coordination tendency of some biologically important zwitterionic buffers toward the binary complexes of some transition metal ions and some amino acids. *J Chem Eng Data* 44(6):1151–1157
103. Coetzee CJ (1989) Determination of formation constants of copper(II) dicarboxylates with a solid state copper(II) ion-selective electrode. *Polyhedron* 8(9):1239–1242
104. Arena G, Cali R, Rizzarelli E, Sammartano S, Barbucci R, Campbell MJM (1978) Thermodynamic and spectroscopic properties of mixed complexes in aqueous solution. Copper(II) complexes of 2,2-bipyridyl and dicarboxylic acids. *J Chem Soc Dalton Trans* 9:1090–1094