

# Tautomerism in drug discovery

Alan R. Katritzky · C. Dennis Hall ·  
Bahaa El-Dien M. El-Gendy · Bogdan Draghici

Received: 1 April 2010 / Accepted: 17 April 2010 / Published online: 20 May 2010  
© Springer Science+Business Media B.V. 2010

**Abstract** The influence of tautomerism on the precise structure of drugs and thus of their potential to interact with biological systems is discussed from thermodynamic and kinetic aspects. The types of tautomerism encountered in the structure of drugs in current use are surveyed together with the effect of pH, solvent polarity, and temperature.

**Keywords** Tautomerism · Heterocycles · Energetics · Biological processes · Drugs

## Introduction

A tautomeric equilibrium is one between two or more isomeric structures of a single compound that are interconverted by movement of an atom (usually hydrogen) or group of atoms from one site to another within the molecular structure. By contrast, two or more isomers each possess the same atomic composition but in general do not interconvert easily. It must be emphasized however, that there is no clear dividing line between tautomerism and isomerism; tautomers are simply isomers that interconvert with a relatively low activation energy below ca 20 kcal/mol.

Keto-enol is probably the best known example of tautomerism. In this case interconversion of the tautomers

may be acid- or base-catalyzed. The relative stability of tautomers may be judged from bond energy differences (Fig. 1) which show that, in this instance of acetaldehyde/vinyl alcohol, the keto form is the more stable isomer by 18 k cal mol<sup>-1</sup>, which corresponds to an equilibrium constant of  $2.6 \times 10^{13}$  at 20 °C.

Tautomeric equilibria are profoundly dependent on the dielectric constant of the medium and the ability of solvents to hydrogen bond with each tautomer with more polar solvent favoring the more polar tautomer. The equilibrium composition is also influenced by temperature, with higher temperatures increasing the proportion of the least stable tautomer (the enol form) since  $\Delta G^\circ = -RT \ln K$ . Since temperature differences are relatively small in most biological systems, temperature effects on tautomerism are unlikely to be relevant to this article or important in drug design. The influence of medium polarity, however, may be highly significant since a biological system may involve an aqueous medium (blood or plasma) or an essentially non-protic medium such as a cell membrane or possibly an enzymatic reaction center.


Molecular structure and solvent polarity clearly define the equilibrium position of a tautomeric system but the rate of interconversion is controlled by the free energy of activation between the interconverting species. If the latter are not energetically equal (i.e.  $K \neq 1$ ) the free energy of activation from high (less stable) to low energy (more stable) species is inevitably lower than the energy for the reverse process. Thus it is important to take both kinetic and thermodynamic factors into consideration (vide infra) when assessing the influence of tautomeric structures on biological activity.

The vast majority of pharmaceutical molecules (drugs) contain heteroaromatic systems with 4-, 5- or 6-membered rings and many of these molecules are capable of existing

---

A. R. Katritzky (✉) · C. D. Hall · B. E.-D. M. El-Gendy ·  
B. Draghici  
Center for Heterocyclic Compounds, Department of Chemistry,  
University of Florida, Gainesville, FL 32611-7200, USA  
e-mail: katritzky@chem.ufl.edu

B. E.-D. M. El-Gendy  
Department of Chemistry, Faculty of Science, Benha University,  
Benha, Egypt



**Fig. 1** Bond energy differences in acetaldehyde-vinyl alcohol, an example of keto-enol tautomerism

as two or more tautomeric structures that usually involve migration of a proton (prototropy) from one site to another within a given molecule.

The importance of prototropy is illustrated dramatically by the change in base pairing brought about by the formation of the imino tautomer of adenine. Within DNA, adenine normally pairs with thymine but the imino form of adenine pairs with cytosine (Fig. 2), thus promoting a misreading of the code: such misreadings are believed to lead to mutations and hence genetic variations that may be the basis of evolution.

Clearly therefore, it is important to recognize the potential for tautomerization in heteroaromatic systems and to assess the role of individual tautomers in biological activity.

### Types of tautomerism in heteroaromatic systems

The vast majority of tautomeric equilibria in heteroaromatic molecules are prototropic, i.e. involve proton migration between (1) carbon and C, N, or O centers; (2) nitrogen and N or O centers, and (3) oxygen and O centers. Details of molecular structures falling into these three general categories have been documented extensively elsewhere [1–4] but several examples are shown in Fig. 3. In general, only those tautomeric systems in which a C–H bond is cleaved and a new C–H bond is formed are capable of isolation as separate tautomers. For example, the pairs of oxo-tautomers of many 2-hydroxy-furans, -pyrroles or -thiophenes [3, 5, 6] (Fig. 3a) are separable and interconvert sufficiently slowly in solution to enable measurement of their individual physical properties.

If a labile proton attached to carbon migrates to N or O, although it is usually difficult to isolate pure individual tautomers, mixtures enriched in one tautomer can often be

obtained [7–9]. However, if the migration is from N to O or O to O, proton exchange is so fast that one almost invariably obtains an equilibrium mixture, the position of which is dependent on the specific conditions (solvent/temperature). This does not mean, however, that one cannot observe individual tautomers by spectroscopic techniques—including multi-nuclear magnetic resonance, ir, uv, ms, etc.

Beyond prototropy, there are tautomeric systems involving cationic or anionic group transfers which, in some instances, manifest themselves as equilibrium mixtures of tautomers again dependent on solvent and temperature. An example of the former is the tautomeric equilibration of N-( $\alpha$ -aminoalkyl)-tetrazoles (Fig. 4), which is an intramolecular migration *via* ion pairs with a  $\Delta G^\circ(298\text{ K}) \approx 0.5\text{ kcal mol}^{-1}$  and a  $\Delta G^\ddagger$  values of 16.3 (low to high energy) and 15.8 kcal mol $^{-1}$  in the reverse direction [10, 11].


Tautomerism of this type, however, is not found in drugs currently widely used and will not be discussed further in this article.

### Energetics

#### Introduction

Thermodynamic and kinetic factors are important to the understanding of tautomeric equilibria and their impact on biology.

In general if the rate at which tautomers interconvert is much slower than the biological process, the most active tautomer would be consumed but the equilibrium would not shift quickly enough to consume both tautomers. Since interaction with the biological system must be second order (at least) the concentration of the active species would influence activity and in a system,






With equilibrium constant,  $K_1 \gg 1$  and  $T_1$  as the active species, the rate of reaction with the biological system, S, would be given by Eq. 1 where  $T_T$  is the total tautomer concentration.

$$\text{Rate} = k_2[S][T_1] = k_2[S][T_T]/(1 + K_1) \quad (1)$$

On the other hand, if the rate of tautomeric interconversion is much faster than the biological process they

**Fig. 2** Base pairing in nucleic acids: Differences for adenine between its normal amino and rare imino forms



**Fig. 3** The common classes of prototropic tautomerism**Fig. 4** Tautomeric equilibrium of N-( $\alpha$ -aminoalkyl)-tetrazoles

influence, the rate of interaction with the biological system again depends on the concentration of the active species but the equilibrium would shift so that all the tautomeric mixture would be consumed. Equation 1 again applies but if  $K_1 \ll 1$ ,  $\text{rate} = k_2[\text{S}][\text{T}_\text{T}]$  and if  $K_1 \gg 1$ ,  $\text{rate} = k_2K_{-1}[\text{T}_\text{T}]$ .

#### Energetics of biological functions

Various biological functions relative to the time scale of the life span of human kind ( $t_{1/2}$  ca.  $10^9$ s) are shown in Table 1 [12]. With these figures in mind, we consider the

energetics of tautomer inter-conversion in terms of both thermodynamics and kinetics.

#### Energetics of Tautomeric Interchange

As mentioned above (Fig. 3), tautomers that interchange by CH bond cleavage/CH bond formation can often be isolated as distinct, stable species and hence the energy barrier between the tautomers must exceed  $30 \text{ kcal mol}^{-1}$  equivalent to a  $t_{1/2}$  value  $\geq 10^4$ s (ca 3 h).<sup>1</sup> With CH to NH, OH, or SH tautomerism, however, the energy barriers are much lower and in the best-known example of keto-enol tautomerism (Scheme 1),  $\Delta G^\ddagger \approx 20 \text{ kcal mol}^{-1}$  giving  $t_{1/2} \approx 0.1 \text{ s}$ . A study of the equilibrium position for such systems is, however, available by multinuclear NMR in

<sup>1</sup> The indistinct line between slowly inter-converting tautomers and structural isomers is generally considered to fall somewhere in between CH/CH tautomerism ( $\Delta G^\ddagger \approx 25\text{--}35 \text{ kcal mol}^{-1}$ ) and *cis/trans* isomers ( $\Delta G^\ddagger \geq 60 \text{ kcal mol}^{-1}$  since the average energy for cleavage of a  $\pi$  bond is  $63 \text{ kcal mol}^{-1}$ ).

**Table 1** Time scale of biological functions

Function	$\Delta G^\ddagger$ at 37 °C (kcal mol <sup>-1</sup> )	$t_{1/2}$ (s)
Macromolecular synthesis (e.g. DNA)	ca 27	$10^2 \rightarrow 10^6$
Cell division	ca 23	$10^3$
Muscle contraction (e.g. heartbeat)	ca 18	1
Antibody complexes, lifetime	15 $\rightarrow$ 21	$10^{-2} \rightarrow 10^2$
Enzyme-substrate complexes	10 $\rightarrow$ 13	$10^{-4} \rightarrow 10^{-6}$
Reciprocal turnovers, enzymes	11 $\rightarrow$ 16	$10^{-5} \rightarrow 10^{-2}$
Photosynthesis; $\alpha$ -helix base pairing	10	$10^{-6}$
H-bonding	4	$10^{-10}$

solvents with a wide range of polarity. The rates of inter-conversion of tautomers can also be studied by variable temperature (VT) NMR. As the temperature is increased the NMR signals from individual tautomers coalesce and at  $T_c$ , the temperature of coalescence, for a two-site exchange involving equal proportions of each tautomer, the rate of exchange at  $T_c$  ( $k_{Tc}$ ) is given by Eq. 2 [13] where  $\Delta\nu$  is the maximum chemical shift difference (in Hz) of signals from each tautomer under “no exchange” conditions.

$$k_{Tc} = \pi\Delta\nu/\sqrt{2} \quad (2)$$

This calculation can also be extended to tautomeric systems with unequal populations of tautomers at equilibrium [14] and thus the rate constants (and hence the  $\Delta G^\ddagger$  values) at  $T_c$  in both directions may be determined. If a number of signals in any one molecular system are observed to coalesce, one may derive a series of  $k_{Tc}$  and  $T_c$  values from which an Arrhenius plot of  $\ln k_{Tc}$  vs.  $1/T_c$  may be generated. This applies to both  $^1\text{H}$  and  $^{13}\text{C}$  signals, or indeed, any nucleus that is observed to coalesce, and results in evaluation of  $E_A$  (and hence  $\Delta H^\ddagger$  and  $\Delta S^\ddagger$ ) over a wide range of  $T_c$  values thus obviating the inherent error of imprecise probe temperature measurements when using line shape analysis over a narrow temperature range. Knowledge of  $\Delta H^\ddagger$  and  $\Delta S^\ddagger$  can provide valuable information on the mechanism (intra- or intermolecular) of tautomerism. With NH to OH or OH to OH tautomerism, the rates are very much faster and therefore rarely amenable to NMR coalescence experiments.

A survey of the top 200 drugs (of year 2010) identified some 33 containing five- or six-membered heterocyclic rings. All examples of potential tautomerism within this


group involved shifts of hydrogen from N to N, N to O, O to O. These processes are all expected to be extremely fast and hence on the biological time scale, the rate of tautomerism would not impact biological processes such as DNA synthesis, cell division, muscle contraction, and probably antibody complexes. On the other hand, an understanding of the relative stability of potential tautomers may well influence enzyme activity, primary processes such as photosynthesis, and base pairing. We next discuss the various types of tautomerism found in current drug molecules.


### Drugs containing tautomeric systems with 5-membered rings

#### NH to N tautomerism

Nexium (1), a drug commonly used to inhibit gastric acid problems [15, 16] and the related compound Protonix (2) [17] both incorporate the benzimidazole unit, but there are no reported studies of their N to N prototropic tautomerism. The mode of action, however, depends on the basicity of the pyridine ring to control the proton pump of the gastrointestinal system and hence the tautomer composition is probably irrelevant (Fig. 5).

The tetrazole ring appears in a number of drugs affording N1 to N2 tautomerism, including Diovan (3) [18], Benicar (4) [19], Avapro (5) [20], Atacand (6) [21] and Hyzaar (7) [22].  $^{15}\text{N}$  CPMAS experiments showed that at room temperature (295 K), the four tetrazole nitrogen atoms gave a very broad signal compared to the imidazole signals, but became sharp at 253 K. These findings are

**Fig. 5** Structure of Nexium (1) and Protonix (2)




**Fig. 6** Structure of drugs 3–7

consistent with fast prototropic exchange as proposed by Harris et al. [23].

These tetrazoles are generally used as cardiovascular or hypertension drugs and the acidity of the tetrazole ring system ( $pK_a \sim 4.8$ ) [24] means they exist largely as anions in biological systems which effectively eliminates the relevance of tautomerism. Losartan potassium salt or Hyzaar (7), the parent drug was approved by FDA 1995 under the commercial name COZAAR. Early studies showed that losartan degrades by dimerisation [25] and the drugs act on very specific targets such as cell receptors of Angiotensin II [26] (Fig. 6).

Imidazolidine-2-imine derivatives include Catapres-TTS (8) and Alphagan P (9) [27] and NMR studies plus computational studies on the former [28] suggest the imino tautomer is more stable than the amino tautomer by ca. 7 kcal mol<sup>-1</sup> ( $K = 8 \times 10^4$  at 37 °C). Lack of experimental information on the relative stabilities or rates of interconversion of the tautomers however, precludes speculation on their importance to biological activity. Suffice it to note that either or both could be active. On the other hand, 2-aminothiazole derivatives exemplified by Mirapex (10) [29] exist, as expected, in the aminothiazole form (Fig. 7).



**Fig. 7** Structure of drugs 8–10

NH to O tautomerism

The examples of drugs containing this type of tautomerism all involve proton migration from N to amide carbonyl oxygen as found in thiazolidine-2,4-dione derivatives Avandia (11) [30], Actoplus Met (12) [31], or oxazolidine-2-one derivatives Skelaxin (13) [32], and Zomig (14) [33]. All of these compounds exist predominantly in the NH form, although the possible significance of the alternative tautomer is recognized for Avandia. Since proton migration would be fast, one or both tautomers could again be available for biological activity (Fig. 8).


## 6-Membered ring drugs containing tautomeric systems

NH to N tautomerism


Drugs with potential NH to N tautomerism include those used for the treatment of HIV 15 [34, 35], 16 [36], epilepsy (17) [37], and skin (18) [38, 39], lung and pancreatic cancers (19) [40–42]. They all involve amine (NH<sub>2</sub> or NH) to imine tautomerism a change that decreases the aromaticity of each heteroaromatic system. Thus the predominant tautomers in these cases contain the amino group (Fig. 9).

NH to O tautomerism

Seven topical drugs (20–26) that could exhibit potential NH to OH tautomerism are all examples of cyclic amide-enamide interconversion and in most cases exist as the amide tautomer which provide centers for the formation of



**Fig. 8** Structure of drugs 11–14



**Fig. 9** Structure of drugs 15–19

intermolecular hydrogen bonds. These drugs are used as sedatives (**20** [43], **22** [44, 45]) and for the treatment of myeloma (**20** [43], **21** [46, 47]), HIV (**22** [44, 45], **24** [48]), schizophrenia (**23** [49]), and Benign prostatic hyperplasia (BPH) (**25** [50, 51], **26** [52]). No studies of tautomerism within these molecular systems have been reported (Fig. 10).


NH to N and NH to O tautomerism within a single molecule

Examples (**27–33**) (Fig. 11) of drugs with potentially both NH to N and NH to O tautomerism, all contain aminopyridone units: (**27** [53, 54], **28** [55, 56], **29** [57–59], **31** [60–62], **32** [63–65]); these are all used as antiviral agents including the treatment of HIV. Shift of a proton from  $\text{NH}_2$  to a heterocyclic N greatly reduces the aromatic character of the pyrimidine ring whilst generating the energetically unfavorable imino tautomer. The potential for NH to O tautomerism is again exhibited in a cyclic amide group with its usual polarity and H-bonding capacity. Xeloda (**33**) [66]—an anti-cancer agent—contains a carbamic ester group which would compete with NH to N tautomerism.


AM1-RHF type semiempirical calculations of Viagra (**30**) [67] revealed that the pyrimidone tautomers (**30A** and **30B**) are more stable than the pyrimidinol tautomer (**30C**) (Fig. 12) [68].

### The influence of pH


The pKa values of most of the drugs mentioned are shown in Table 2 but it should be noted that not all of the values refer to the potentially tautomeric proton. The pKa of the tautomeric proton is, however, an important factor since, if it falls below normal physiological pH (7.2) the molecule will exist as the anion and the resultant delocalized anion would render tautomerism of the system redundant. This is the case for compounds **1–7**. The first two (Nexium and Protonix) each contain a benzimidazole unit that would be ionized at pH 7.2. Since both drugs are used to treat gastric problems, however, they would experience pH values of 1–2 under which conditions they would be protonated. However, in principle, Nexium could exist as two tautomers by virtue of the unsymmetrical structure imposed



**Fig. 10** Structure of drugs 20–26



**Fig. 11** Structure of drugs 27–33



**Fig. 12** Possible tautomers of Viagra



**Table 2** Drug's pKa

Entry	Drug	pKa	Entry	Drug	pKa
1	Nexium	4.00 (A) and 8.80 (L)	18	Aldara	7.30 (A)
2	Protonix	≈4.00 (A)	19	Tarceva (Erlotinib)	5.40 (A)
3	Diovan	NA	20	Thalidomide	3.75 (?)
4	Benicar	4.30 (L)	21	Revlimid	NA
5	Avapro	4.20 (L)	22	Kaletra	NA
6	Atacand	NA	23	Abilify	7.60 (A)
7	Hyzaar	NA	24	Efavirenz	10.20 (?)
8	Catapress TTS	9.50 (A)	25	Avodart	NA
9	Alphagan P	7.40 (A)	26	Propecia	9.41 (?)
10	Mirapex	9.60 (A)	27	Valaciclovir HCl (Valtrex)	1.90, 7.47, and 9.43
11	Avandia	6.10 (A) and 6.80 (L)	28	Acyclovir (Zovirax)	2.27 and 9.25
12	Actoplus MET	12.40 (L)	29	Valganciclovir HCl (Valcyte)	7.60 (A)
13	Skelaxin	NA	30	Viagra	NA
14	Zomig	9.64 (A)	31	Emtricitabin (Emtriva)	2.65 (A)
15	Abacavir	5.01 (A)	32	Lamivudine	4.30 (A)
16	Tenofovir DF	3.75 (L)	33	Capecitabine (Xeloda)	5.41 (?)
17	Lamictal	5.70 (A)			

NA data are not available

In Table 2 we have recorded reported values for the pK<sub>a</sub>s of the drugs. In some cases it is clear whether these values refer to proton loss (L) or proton addition (A). However, in other cases, it is not clear what the reported pK<sub>a</sub> values refer to, and in some cases these reported values appear to be very uncertain and unreliable (?). We have not commented in detail on these cases

by the methoxy-substituted benzimidazole unit whereas the tautomeric structures in Protonix are equivalent. Compounds **3–7**, used as cardiovascular or hypertension drugs, all contain a tetrazole unit with a pK<sub>a</sub> value of ca. 4.8. Thus, at normal physiological pH they would be present as anions and tautomerism would be irrelevant.

The pK<sub>a</sub> values of Catapress (**8**) and Alphagan P (**9**) at 9.5 and 7.4 respectively obviously refer to proton addition but tautomerism between the more stable neutral imino form and the neutral exocyclic amino structure is likely to be fast on most biological time scales so either structure could be active.

Compounds **11–14** all involve amide tautomerism with the NH form predominating but fast prototropy could render either amide or the imidol structure available for biological activity. The same applies to compounds **20–26**. Compounds **15–19** all involve NH (or NH<sub>2</sub>) to ring N tautomerism but since the imino form substantially reduces aromaticity in the hetero ring, the amino form greatly predominates. Fast prototropy could, however, again make either structure available for therapeutic activity. The pK<sub>a</sub> values (apart from Tenofovir DF, **16**, containing the –P(O)(OH)<sub>2</sub> group at pK<sub>a</sub> 3.75) all refer to proton addition to the ring N.

Compounds **27–33** all contain both NH to N and NH to O potential tautomerism within aminopyrimidone units. At physiological pH, all of these molecules probably remain

unprotonated and the pyrimidinone structures would predominate in fast prototropic equilibria.

## Conclusions

Tautomerism in drug molecules almost invariably involves prototropy but in all cases the influence of a single tautomer on therapeutic activity depends on the time scale of the tautomeric equilibrium relative to that of the biological process in question. Thus in judging the influence of tautomerism on biological activity, it is essential to consider both thermodynamic and kinetic factors. Fast interconversion of tautomers relative to a specific biological process means that both tautomers may be consumed. Conversely, slow interconversion relative to the biological process may result in one tautomer being the only active species.

## References

- Katritzky AR, Lagowski JM (1963) Adv Heterocycl Chem (Katritzky AR (eds) Academic Press, New York, N.Y.) 1: 311–338
- Katritzky AR, Lagowski JM (1963a) Adv Heterocycl Chem (Katritzky AR (eds) Academic Press, New York) 1: 339–437
- Katritzky AR, Lagowski JM (1963b) Adv Heterocycl Chem (Katritzky AR (eds) Academic Press, New York) 2: 1–26



4. Elguero J, Marzin C, Katritzky A, Linda P (1976) Adv Heterocycl Chem, Supplement 1: the tautomerism of heterocycles. Academic Press, New York
5. Katritzky AR (1970) *Chimia* 24:134–146
6. Gronowitz S, Hoffman RA (1960) *Arkiv foer Kemi* 15:499–512
7. Hino T, Nakagawa M, Hashizume T, Yamaji N, Miwa Y, Tsuneoka K, Akaboshi S (1971) *Tetrahedron* 27:775–787
8. Dickinson RP, Iddon B (1970) *J Chem Soc C* 14:1926–1928
9. Woerner FP, Reimlinger H, Merenyi R (1971) *Chem Ber* 104:2789–2792
10. Katritzky AR, Jozwiak A, Lue P, Yannakopoulou K, Palenik GJ, Zhang ZY (1990) *Tetrahedron* 46:633–640
11. Katritzky AR, El-Gendy B-EM, Draghici B, Hall CD, Steel PJ (2010) *Manuscript in preparation*
12. Cox BG (1994) Modern liquid phase kinetics. Oxford University Press, New York
13. Hore PJ (1995) Nuclear magnetic resonance. Oxford University Press, New York
14. Shanan-Atidi H, Bar-Eli KH (1970) *J Phys Chem* 74:961–963
15. McKeage K, Blick SKA, Croxtall JD, Lyseng-Williamson KA, Keating GM (2008) *Drugs* 68:1571–1607
16. Inoue M, Zhai H, Sakazaki H, Furuyama H, Fukuyama Y, Hirama M (2004) *Bioorg Med Chem* 14:663–665
17. Cheer SM, Prakash A, Faulds D, Lamb M (2003) *Drugs* 63:101–132
18. Th Moenius VR, Burtcher P, Zuger Ch (2000) *J Labelled Compd Radiopharm* 43:1245–1252
19. Becker A (2008) PCT Int Appl WO 2008/149155A1
20. Alkorta I, Rozas I, Elguero J (1998) *J Chem Soc Perkin Trans* 2:2671–2675
21. Mulvany MJ, Skov K, Eiskjær H, Hansen HE, Madsen JK, Kvist S, Mulvany MJ (2007) *Hypertension* 50:89–95
22. Wilson TW, Lacourciere Y, Barnes CC (1998) *CMAJ* 159:469–476
23. Bauer M, Harris RK, Rao RC, Apperley DC, Rodger CA (1998) *J Chem Soc, Perkin Trans* 2:475–482
24. Myznikov LV, Hrabalek A, Koldobskii GI (2007) *Chem Heterocycl Comp* 43:1–9
25. McCarthy KE, Wang Q, Tsai EW, Gilbert RE, Dominic P, Brooks MA (1998) *J Pharm Biomed Anal* 17:671–677
26. Cagigal E, Gonzalez L, Alonso RM, Jimenez RM (2001) *J Pharm Biomed Anal* 26:477–486
27. Fudenberg SJ, Batiste C, Katz LJ (2008) *Expert Opin Drug Saf* 7:795–799
28. Jackman IB LM, Jen T (1975) *J Am Chem Soc* 97:2811–2818
29. Varga LI, Ako-Agugua N, Colasante J, Hertweck L, Houser T, Smith J, Watty AA, Naga S, Raffa RB (2009) *J Clin Pharm Ther* 34:493–505
30. Pool CR, Roman RS, Brighwell MD, Tremper AW (1994) PCT Int Appl WO9405659A1
31. Seufert J (2006) *Curr Med Res Opin* 22:S39–S48
32. Poklis JL, Roper-Miller JD, Garside D, Winecker RE (2004) *J Anal Toxicol* 28:537–540
33. Rapoport AM, Bigal ME, Tepper SJ, Sheftell FD (2004) *Expert Rev Neurotherapeutics* 4:33–41
34. Daluge SM, Good SS, Faletto MB, Miller WH, St. Clair MH, Boone LR, Tisdale M, Parry NR, Reardon JE, Dornsife RE, Averett DR, Krenitsky TA (1997) *Antimicrob Agents Chemother* 41:1082–1093
35. Daluge SM (1991) U.S. Patent US5034394A
36. Deeks SG, Barditch-Crovo P, Lietman PS, Hwang F, Cundy KC, Rooney JF, Hellmann NS, Safran S, Kahn JO (1998) *Antimicrob Agents Chemother* 42:2380–2384
37. Binnie CD, Boas WVE, Land W, Meijer GS, Overweg JWA, Van Wieringen A (1985) *Br J Clin Pharmacol* 20: 285 (abstract)
38. Beutner KR, Geisse JK, Helman D, Fox TL, Ginkel A, Owens ML (1999) *J Am Acad Dermatol* 41:1002–1007
39. Sidky YA, Borden EC, Weeks CE, Reiter MJ, Hatcher JF, Bryan GT (1992) *Cancer Res* 52:3528–3533
40. Rusch V, Baselga J, Cordon-Cardo C, Orazem J, Hoda S, McIntosh J, Kurie J, Dmitrovsky E (1993) *Cancer Res* 53:2379–2385
41. Grandis JR, Tweardy DJ (1993) *Cancer Res* 53:3579–3584
42. Moyer JD, Barbacci EG, Iwata KK, Arnold L, Boman B, Cunningham A, Diorio C, Doty J, Morin MJ, Moyer MP, Neveu M, Pollack VA, Pustilnik LR, Reynolds MM, Sloan D, Theleman A, Miller P (1997) *Cancer Res* 57:4838–4848
43. Bartlett JB, Dredge K, Dalglish AG (2004) *Nat Rev Cancer* 4:314–322
44. Laffitte E, Revuz J (2004) *Expert Opin Drug Saf* 3:47–56
45. Murphy RL, Brun S, Hicks C, Eron JJ, Gulick R, King M, White AC, Benson C, Thompson M, Kessler HA, Hammer S, Bertz R, Hsu A, Japour A, Sun E (2001) *AIDS* 15:F1–F9
46. Dimopoulos M, Spencer A, Attal M, Prince HM, Harousseau JL, Dmoszynska A, San Miguel J, Hellmann A, Facon T, Foa R, Corso A, Masliak Z, Marta Olesnyckyj, Yu Z, Patin J, Zeldis JB, Knight RD (2007) *N Engl J Med* 357:2123–2132
47. Weber D, Chen C, Niesvizky R, Chen C, Niesvizky R, Wang M, Belch A, Stadtmayer EA, Siegel D, Borrello I, Rajkumar SV, Chanan-Khan AA, Lonial S, Yu Z, Patin J, Olesnyckyj M, Zeldis JB, Knight RD (2007) *N Engl J Med* 357:2133–2142
48. Young SD, Britcher SF, Tran LO, Payne LS, Lumma WC, Lyle TA, Huff JR, Anderson PS, Olsen DB, Carroll SS, Pettibone DJ, O'Brien JA, Ball RG, Balani SK, Lin JH, Chen I-WU, Schleif WA, Sardana VV, Long WJ, Byrnes VW, Emini EA (1995) *Antimicrob Agents Chemother* 39:2602–3605
49. Kane JM, Carson WH, Saha AR, McQuade RD, Ingenito GG, Zimbroff DL, Ali MW (2002) *J Clin Psychiatry* 63:763–771
50. Thomas LN, Douglas RC, Lazier CB, Too CKL, Rittmaster RS, Tindall DJ (2008) *Eur Urol* 53:244–252
51. Miller J, Tarter TH (2009) *Clin Interv Aging* 4:251–258
52. Gormley GJ, Stoner E, Rittmaster RS, Gregg H, Thompson DL, Lasseter KC, Vlasses PH, Steven EA (1990) *J Clin Endocrinol Metab* 70:1136–1141
53. Beutner KR, Friedman DJ, Forszpaniak C, Andersen PL, Wood MJ (1995) *Antimicrob Agents Chemother* 39:1546–1553
54. DeJesus E, Wald A, Warren T, Schacker TW, Trotter S, Shahmanesh M, Hill JL, Brennan CA (2003) *J Infect Dis* 188:1009–1016
55. Elion GB, Furman PA, Fyfe JA, de Miranda P, Beauchamp L, Schaeffer HJ (1977) *Proc Natl Acad Sci USA* 74:5716–5720
56. Schaeffer HJ, Beauchamp L, de Miranda P, Elion GB, Bauer DJ, Collins P (1978) *Nature* 272:583–585
57. McGavin JK, Goa KL (2001) *Drugs* 61:1153–1183
58. Singh N (2000) *Clin Liver Dis* 4:657–673
59. Van der Bij W, Speich R (2001) *Clin Infect Dis* 33(Suppl 1):S32–S37
60. Schinazi RF, McMillan A, Cannon D, Mathis R, Lloyd RM, Peck A, Sommadossi JP, St Clair M, Wilson J, Furman PA, Painter GR, Choi WB, Liotta DC (1992) *Antimicrob Agents Chemother* 36:2423–2431
61. Furman PA, Davis M, Liotta DC, Paff M, Frick LW, Nelson DJ, Dornsife RE, Wurster JA, Wilson LJ, Fyfe JA, Tuttle JV, Miller WH, Condreay L, Averett DR, Schinazi RF, Painter GR (1992) *Antimicrob Agents Chemother* 36:2686–2692
62. Tisdale M, Kemp SD, Parry NR, Larder BA (1993) *Proc Natl Acad Sci USA* 90:5653–5656
63. Lai CL, Chien RN, Leung NWY, Chang TT, Guan R, Tai DI, Ng KY, Wu PC, Debt JC, Barber J, Stephenson SL, Gray DF (1998) *N Engl J Med* 339:61–68
64. Dienstag JL, Schiff ER, Wright TL, Perrillo RP, Hann HWL, Goodman Z, Crowther L, Condreay LD, Woessner M, Rubin M, Brown NA (1999) *N Engl J Med* 341:1256–1263

65. Tassopoulos NC, Volpes R, Pastore G, Heathcote J, Buti M, Goldin R, Hawley S, Gray DF et al (1998) *Hepatology* 28(Suppl):43
66. Ishitsuka H, Miwa M, Ishikawa T, Ura M, Sawada N, Nishida M, Fukase Y, Yamamoto T, Mori K, Arasaki M, Umeda I, Shimma N (1995) *Proc Am Assoc Cancer Res* 36:407
67. Terrett NK, Bell AS, Brown D, Ellis P (1996) *Bioorg Med Chem Lett* 6:1819–1824
68. Turker L (2000) *ACH-Models Chem* 137:767–774