

# Molecular basis for the enantioselective binding of a novel class of cytochrome bc1 complex inhibitors

Ya-Jun Zheng\*

*DuPont Crop Protection, Stine-Haskell Research Center, P.O. Box 30, Newark, DE 19714, USA*

Received 11 October 2005; received in revised form 10 November 2005; accepted 10 November 2005

Available online 20 December 2005

## Abstract

The recently solved co-crystal structures of mitochondrial cytochrome bc1 complex with inhibitors have provided an important structural framework for the elucidation of modes of binding of various bc1 complex inhibitors. *N*-Phenyl triazolones, a novel class of bc1 complex ubiquinol oxidation ( $Q_o$ )-site inhibitors, were found to exhibit atropisomerism; in few cases, the atropisomers were resolved and shown to express different biological activities. However, the underlying mechanism for such differential binding of the enantiomers to bc1 complex is unknown. Here molecular docking is used to examine the binding modes of the *N*-phenyl triazolones fungicides. Our docking studies allow the molecular basis for the enantioselective binding of atropisomeric triazolones to be elucidated. Furthermore, the mode of binding of azoxystrobin has also been clarified.

© 2005 Elsevier Inc. All rights reserved.

**Keywords:** Fungicide; Strobilurin; Molecular docking; Atropisomeric; Scoring

## 1. Introduction

Mitochondrial cytochrome bc1 complex (ubiquinol–cytochrome *c* oxidoreductase) is a very fruitful target for the discovery of fungicides useful in controlling crop diseases [1–4]. Two distinct ubiquinone (or ubiquinol) binding sites are known: ubiquinol oxidation site ( $Q_o$  site) and ubiquinol reduction site ( $Q_i$  site) [5]. Several  $Q_o$ -site inhibitors (e.g., 2–5 in Scheme 1) and one  $Q_i$ -site inhibitor (6) of the bc1 complex have been commercialized [1–4].

Most of the commercialized  $Q_o$ -site inhibitors of the cytochrome bc1 complex could be considered as analogues of the naturally occurring  $Q_o$ -site inhibitor strobilurins and oudemansins [3]; famoxadone represents a structurally distinct  $Q_o$ -site inhibitor [2]. Lead optimization programs at several companies have resulted in a large number of strobilurin analogues and structure–activity relationships (SARs) were elucidated for those analogue series. The methoxyacrylate head group as in strobilurin A is normally considered as the pharmacophore (or toxiphore) for the inhibition of cytochrome bc1 complex. However, as shown in Scheme 1, the


methoxyacrylate can be replaced by isosteric groups as in kresoxim-methyl and metominostrobin without losing biological activity [3]. Scheme 2 lists several additional variations of this pharmacophore. Early SAR study also established the requirement of the E configuration of the central double bond for biological activity since the Z isomer is inactive [6]. Recently, X-ray crystallographic studies on cytochrome bc1 complexed with various inhibitors have been reported [7–11].

*N*-Phenyl triazolones (as exemplified by compound 12 in Scheme 3) are one novel class of strobilurin analogues [12]. The triazolone ring could be considered as a tied-back version of the methoxyacrylate pharmacophore. Like the biphenyls, such *N*-phenyl-triazolone strobilurin analogs are twisted with large inter-ring torsion angles [13]. In principle, they could exhibit atropisomerism. Indeed, when the phenyl ring is substituted at the ortho positions (positions 2 and 6), the torsion barrier around the inter-ring C–N bond is large enough to allow atropisomers to be stable even at room temperature [13]. The atropisomers of compound 12 have been resolved and their bioactivities were evaluated [12]. One enantiomer is found to be much more active than the other. However, the underlying mechanism for such enantioselective binding is unknown.

Here molecular docking is used to elucidate the mode of binding of compound 12 in the hope of uncovering the underlying mechanism of differential binding of the enantiomers.

\* Tel.: +1 302 451 4604; fax: +1 302 366 5738.

E-mail address: [ya-jun.zheng@usa.dupont.com](mailto:ya-jun.zheng@usa.dupont.com).



Scheme 1.

## 2. Theoretical procedures


Molecular docking was performed using QXP program [14], a part of the Flo+ software [15]. The MCDOCK and MCDOCK+ procedures were used. For each inhibitor, an initial 100 Monte Carlo steps were used to place the inhibitor inside the binding pocket; then 10,000 repeated cycles of Monte Carlo (MC) followed by energy minimization and additional scoring were used to yield an ensemble of 25 top scoring inhibitor poses. Simulations were repeated several times using different starting inhibitor conformations to assess convergence. The protein binding pocket surface mesh was generated using Flo+ with a previously described procedure [16]. The co-crystal structure of bc1 with bound azoxystrobin (3) was used [8c]. To assess the reliability of the docking software, the bound azoxystrobin was taken out and flexibly redocked into the binding site. Owing to the low resolution of the X-ray crystal structure, there were some uncertainties concerning the details of the bound conformation of azoxystrobin. For instance, the conformation of the methoxyacrylate head group could be either *s-cis* or *s-trans*. Previous SAR data were then used in conjunction with the docking experiments to elucidate the

binding mode of azoxystrobin. Once the performance of the docking software is established, we proceed to examine the binding of compound 12. A pdb file containing the docked compound 12 in the binding pocket is available upon request from the author.

## 3. Results and discussions

### 3.1. Mode of binding of azoxystrobin

For conjugated unsaturated compounds as shown in Scheme 4, there are two possible conformation, *s-cis* and *s-trans*. The energy difference between them is dependent of the steric size of the X group. For acrolein (where X = H), the sterically less crowded *s-trans* conformation is favored over the *c-cis* conformation by 1.7 kcal/mol and the interconversion energy barrier is about 5.0–6.4 kcal/mol [17]. In the cases of acrylic acid (X = OH) and methyl acrylate (X = OCH<sub>3</sub>), the energy differences between the *s-cis* and *s-trans* conformations are only 0.17 kcal/mol and 0.31 kcal/mol, respectively [17]. Even though the energy difference between the *s-cis* and *s-trans* conformation of azoxystrobin has not been determined




Scheme 2.

experimentally, it is very reasonable to assume that the energy difference is rather small and both conformations are equally plausible.

The X-ray co-crystal structure of bc1 with azoxystrobin was determined at 2.7 Å resolution [8c]. Since the cytochrome b structure has been solved several times under different conditions (with and without inhibitors), the structure of the protein itself is accurate. However, at such resolution, there is uncertainty regarding the bound conformation of the small molecular weight inhibitor. In fact, in the bc1/azoxystrobin structure, the


methoxyacrylate is refined as *s-cis* while in the analogous bc1/myxothiazol, an *s-trans* conformation was assumed. To address this issue, the bound azoxystrobin was taken out of the active and re-docked into the binding site. During the docking process, the protein is kept at the X-ray structure and the ligand is flexibly docked. The top 25 scoring structures were examined in detail and Fig. 1 showed the overlap of top 25 scoring docked poses of azoxystrobin in the Q<sub>o</sub> site.

Overall, the docking poses are quite consistent. The pyrimidine and the phenyl rings are essentially at the same locations in the binding pocket. The position of cyanophenoxy shows considerable variations, which are consistent with the active site topology. The methoxyacrylate moiety is *s-trans* in all of the 25 poses as opposed to the *s-cis*; the docked poses also indicate that the ester function could be in either Z or E configuration. This is likely an artifact of the simulation since it is known experimentally that the Z configuration (CH<sub>3</sub> is trans to the acrylate C=O bond) is generally favored in carboxylic esters [18]. Fig. 2 displays the overlap of the docked poses with the ester in the Z configuration and the refined structure (white) by Esser et al. [8c]. All of the top scoring docked poses agree with the refined structure by Esser et al. [8c] with respect to the



Scheme 3.





Scheme 5.

that of azoxystrobin, again consistent with the notion that the triazolone pharmacophore of **12** is a tied-back version of the normal methoxyacrylate pharmacophore of strobilurins.

The binding mode of **12** in the binding site also provides a straightforward explanation as to why only the (+)-**12** enantiomer is biologically active. Binding of the (–)-**12** enantiomer would require the flip of the triazolone ring along the inter-ring C–N bond in the binding pocket, switching positions of C=O with –OCH<sub>3</sub>, and the N–CH<sub>3</sub> with the sp<sup>2</sup> ring N atom. Since in the binding site (Fig. 3), the sp<sup>2</sup> ring N is already on the surface mesh, switching the sp<sup>2</sup> ring N by N–CH<sub>3</sub> will result in significant clash between the methyl group

and protein atoms. Thus, the (–)-**12** enantiomer cannot fit well in the binding pocket.


Xia and co-workers proposed previously that aromatic–aromatic (Ar–Ar) interactions play a critical role in the binding of famoxadone and azoxystrobin to the bc1 complex [8]. Several aromatic amino acids such as Tyr-131, Tyr-278, Tyr-273, and Phe-274 were found to be in close contact with the bound famoxadone [8a]. Interestingly, Tyr-131, Tyr-278, and Phe-274 are also in close contact with (+)-**12** in the binding pocket as revealed by our docking calculations. Thus, the proposed Ar–Ar interactions may also contribute significantly to the binding affinity of *N*-phenyl triazolones.

#### 4. Conclusions

The binding mode of azoxystrobin in the O<sub>o</sub> site of mitochondrial bc1 complex was examined using a molecular docking procedure. Based on previous extensive SAR, X-ray crystallographic studies, and the present molecular docking study, a consistent binding mode for azoxystrobin is presented. In this binding mode, azoxystrobin forms a hydrogen bond with the amide N–H of Pro270 through the ester carbonyl oxygen atom. The methoxyacrylate is in the *s-trans* form instead of the *s-cis* as proposed by Esser et al. previously. Subsequently, the binding mode of *N*-phenyl triazolone inhibitors with a conformationally constrained pharmacophore is investigated. According to our docking study, the binding site topology is consistent with only (+)-**12**, the biologically active enantiomer; the other atropisomer does not fit well. Our study also demonstrated the usefulness of molecular docking in elucidating the mode of binding of bc1 inhibitors. Similar approaches could be used to examine the binding modes of other novel inhibitors.

#### References

- [1] (a) P. Leroux, Recent developments in the mode of action of fungicides, *Pestic. Sci.* 47 (1996) 191–197;  
 (b) P.R. Rich, Quinone binding sites of membrane proteins as targets for inhibitors, *Pestic. Sci.* 47 (1996) 287–296.
- [2] (a) J.A. Sternberg, D. Geffken, J.B. Adams Jr., R. Postages, C.S. Sternberg, C.L. Campbell, W.K. Moberg, Famoxadone: the discovery and optimisation of a new agricultural fungicide, *Pest Manag. Sci.* 57 (2001) 143–152;  
 (b) Y.-J. Zheng, R. Shapiro, W.J. Marshal, D.B. Jordan, Synthesis and structural analysis of the active enantiomer of famoxadone, a potent inhibitor of cytochrome bc1, *Bioorg. Med. Chem. Lett.* 10 (2000) 1059–1062;  
 (c) D.B. Jordan, K.T. Kranis, M.A. Piccollelli, R.S. Schwartz, J.A. Sternberg,

Fig. 3. Binding of compound (+)-**12** in the Q<sub>o</sub> site in two (a and b) perspectives.



- K.M. Sun, Famoxadone and oxazolidinones: potent inhibitors of cytochrome bc1, *Biochem. Soc. Trans.* 27 (1999) 577–580;
- (d) S.O. Pember, L.C. Fleck, W.K. Moberg, M.P. Walker, Mechanistic differences in inhibition of ubiquinol cytochrome *c* reductase by the proximal Q<sub>o</sub>-site inhibitors famoxadone and methoxyacrylate stilbene, *Arch. Biochem. Biophys.* 435 (2005) 280–290.
- [3] H. Sauter, W. Steglich, T. Anke, Strobilurins: evolution of a new class of active substances, *Angew. Chem. Int. Ed.* 38 (1999) 1328–1349.
- [4] T. Ohshima, T. Komyoji, S. Mitani, N. Matsuo, T. Nakajima, Development of a novel fungicide, cyazofamid, *J. Pestic. Sci.* 29 (2004) 136–138.
- [5] A.R. Crofts, The cytochrome bc1 complex: function in the context of structure, *Annu. Rev. Physiol.* 66 (2004) 689–733.
- [6] T. Anke, W. Steglich,  $\beta$ -Methoxyacrylate antibiotics: from biological activities to synthetic analogues, in: U.P. Schlunegger (Ed.), *Biologically Active Molecules*, Springer, Berlin, 1989, pp. 9–25.
- [7] D. Xia, C.A. Yu, H. Kim, J.Z. Xia, A.M. Kachurin, L. Zhang, L. Yu, J. Deisenhofer, Crystal structure of the cytochrome bc1 complex from bovine heart mitochondria, *Science* 277 (1997) 60–66.
- [8] (a) X. Gao, X. Wen, C. Yu, L. Esser, S. Tsao, B. Quinn, L. Zhang, L. Yu, D. Xia, The crystal structure of mitochondrial cytochrome bc1 in complex with famoxadone: the role of aromatic-aromatic interaction in inhibition, *Biochemistry* 41 (2002) 11692–11702;
- (b) X. Gao, X. Wen, L. Esser, B. Quinn, L. Yu, D. Xia, Structural basis for the quinone reduction in the bc1 complex: a comparative analysis of crystal structures of mitochondrial cytochrome bc1 with bound substrate and inhibitors at the Q<sub>i</sub> site, *Biochemistry* 42 (2003) 9067–9080;
- (c) L. Esser, B. Quinn, Y. Li, M. Zhang, M. Elberry, L. Yu, C. Yu, D. Xia, Crystallographic studies of quinol oxidation site inhibitors: a modified classification of inhibitors for the cytochrome bc1 complex, *J. Mol. Biol.* 341 (2004) 281–302.
- [9] (a) Z. Zhang, L. Huang, V.M. Shulmeister, Y.I. Chi, K.K. Kim, L.W. Hung, A.R. Crofts, E.A. Berry, S.H. Kim, Electron transfer by domain movement in cytochrome bc1, *Nature* 392 (1998) 677–684;
- (b) L. Huang, D. Cobessi, E.Y. Tung, E.A. Berry, Binding of the respiratory chain inhibitor antimycin to the mitochondrial bc1 complex: a new crystal structure reveals an altered intramolecular hydrogen-bonding pattern, *J. Mol. Biol.* 351 (2005) 573–597.
- [10] S. Iwata, J.W. Lee, K. Okada, J.K. Lee, M. Iwata, B. Rasmussen, T.A. Link, S. Ramaswamy, B.K. Jap, Complete structure of the 11-subunit bovine mitochondrial cytochrome bc1 complex, *Science* 281 (1998) 64–71.
- [11] (a) C. Hunte, J. Koepke, C. Lange, T. Rossmann, H. Michel, Structure at 2.3 Å resolution of cytochrome bc(1) complex from yeast *Saccharomyces cerevisiae* co-crystallized with an antibody Fv fragment, *Structure* 15 (2000) 669–684;
- (b) H. Palsdottir, C.G. Lojero, B.L. Trumpower, C. Hunte, Structure of the yeast cytochrome bc1 complex with a hydroxyquinone anion Q<sub>o</sub> site inhibitor bound, *J. Biol. Chem.* 278 (2003) 31303–31311.
- [12] (a) R.J. Brown, G. Annis, A. Casalnuovo, D. Chan, R. Shapiro, W.J. Marshall, Synthesis and properties of axially-chiral *N*-(2,6-disubstituted)-phenyl triazolones, *Tetrahedron* 60 (2004) 4361–4375;
- (b) R.J. Brown, B. Ashworth, J.E. Drumm, D.A. Frazier, M.A. Hanagan, C. Happersett, G.E. Koether, D.J. Robinson, K.-M. Sun, P. Wojtkowski, Synthesis of fungicidal triazolones, in: D.R. Baker, J.G. Fenyes, G.P. Lahm, T.P. Selby, T.M. Stevenson (Eds.), *Synthesis and Chemistry of Agrochemicals VI*, ACS Symposium Series 800, American Chemical Society, 2002, pp. 327–339.
- [13] Y.-J. Zheng, D.A. Kleier, Conformational flexibility of antifungal atropisomeric strobilurin analogues: a quantum mechanical investigation, *J. Mol. Struct. THEOCHEM* 719 (2005) 69–74.
- [14] C. McMartin, R.S. Bohacek, QXP: powerful, rapid computer algorithms for structure-based drug design, *J. Comput. Aided Mol. Des.* 11 (1997) 333–344.
- [15] Flo+, ThistleSoft, 605 Colebrook Road, Colebrook, CT, 2004.
- [16] R.S. Bohacek, C. McMartin, Definition and display of steric, hydrophobic, and hydrogen-bonding properties of ligand binding sites in proteins using Lee and Richards accessible surface: validation of a high-resolution graphical tool for drug design, *J. Med. Chem.* 35 (1992) 1671–1684.
- [17] E.L. Eliel, S.H. Wilen, *Stereochemistry of Organic Compounds*, Wiley Interscience, John Wiley & Sons, Inc., 1993, p. 662 and the reference cited therein.
- [18] S. Ruschin, S.H. Bauer, Rotational isomers of methyl formate, located by the temperature-drift technique, *J. Phys. Chem.* 84 (1980) 3061.