

Conformational analysis using the HSEA-program¹, (basically Van de Waals energies) indicates that the conformations of B-C and A-C are very similar to those found in the trisaccharide. Comparison of the rotational freedom of B-C and A-C shows that the latter is more flexible. NMR-spectra, on the other hand, show large changes in chemical shifts for unit A. Thus from these results it can be concluded that several factors including conformational changes are responsible for the changes of the chemical shifts.

Reference

- 1 Lemiux, R U et al. *Can. J. Chem.* Vol 58 (1980) p 631

42


Three-dimensional and electronic analysis of Na⁺-dependent antidopaminergic agents: tropapride and zetidoline

S Collin*, D P Vercauteren*, G Evrard*, F Durant*, H van de Waterbeemd†, P A Carrupt† and B Testa†

*Laboratoire de Chimie Moléculaire Structurale, Facultés Universitaires Notre-Dame de la Paix, rue de Bruxelles, 61, B-5000 Namur, Belgium

†Ecole de Pharmacie, Université de Lausanne, CH-1005 Lausanne, Switzerland

'Tropapride' belongs to a well-known neuroleptic class: the benzamide nortropanes whose most significant feature is a selective binding to D₂ receptors in a highly Na⁺-dependent manner¹. A first 3D and electronic analysis of several analogues of tropapride has lead us to suggest a model of three pharmacophoric elements, a basic nitrogen lone pair, a phenyl ring, and a carbonyl function, for ligands at the D₂ receptor². In order to confirm this model, a comparison will be made of tropapride to zetidoline, the lead compound of a new neuroleptic family with the same biochemical properties as other atypical neuroleptics³.



Using a user-interactive flexible molecular fitting program, IFMFIT, zetidoline has been shown to possess the three pharmacophoric elements observed for benzamides. In addition, both show a similar steric hindrance. Molecular electrostatic potential maps obtained at the *ab initio* SCF-LCAO-MO level for zetidoline show that it has two well-defined regions in common with tropa-

pride: the negative and positive regions lie on opposite sides of the carbonyl and methoxy groups respectively.

References

- 1 Jalfre, M et al. *Arch. Int. Pharmacodyn. Ther.* Vol 264 (1983) p 232
- 2 Collin, S et al. *J. Cryst. Spec. Res.* Vol 16 (1986) p 255
- 3 Barone, D et al. *J. Pharmacol.* Vol 37 (1984) p 180

43

Computer assisted investigation of structure activity relationship in α_2 adrenoceptor ligands

P Bovy*, A De Prijck* and F Durant*

*Continental Pharma, 11 rue Granbonpre, B-1348 Mont-Saint-Guibert, Belgium

**FUNDP, 61 rue de Bruxelles, B-5000 Namur, Belgium

The aim of this work was to investigate within a set of molecules the common minimal structural features required to elicit α_2 adrenoceptor affinity in radioligand displacement experiments. *In vitro* radioligand binding assays were carried out using standard published procedures using *p*-aminoclonidine as the radioligand¹. The set of compounds was composed of reference drugs (rauwolscine², mianserine², idazoxan³, 1-methyl-4-(3-fluoro-2-pyridinyl)-piperazine²) and original compounds from Continental Pharma's collection (CP2953⁴, CP3848) displaying, to different extents, *in vitro* α_2 adrenoceptor affinity. Modelling studies were performed using a software package developed in FUNDP⁵. The workstation consists of a Megatek 7210 graphic processor connected to a host CPU processor PDP 11/60. Rauwolscine has been used as a reference template. We have been able to find good superpositions for a phenyl ring and a nitrogen atom (*sp*² or *sp*³ hybridized) common to all compounds. The 3D structures used for modelling studies were obtained either from X-ray data of the molecule itself or fragment assembly from the Cambridge Crystallographic Data File. For the flexible molecules, we have found different conformations with acceptable internal energy that allow superpositions. Thus all the molecules with affinity to the α_2 adrenoceptors have at least one conformation in which a N-atom is at a fixed position in space relative to an aromatic ring: the mean distance is 5.6 ± 0.15 Å with the nitrogen atom roughly lying in the same plane as the phenyl ring.

References

- 1 Rouot, B R and Snyder, S H *Life Sci.* Vol 25 (1979) pp 769-774
- 2 Pinder, R M *Drugs Fut.* Vol 10 (1985) pp 841-857
- 3 Chapleo, C B et al. *J. Med. Chem.* Vol 26 (1983) p 823
- 4 Lu Patent 85747 (1985)
- 5 Lejeune, J and Michel, A in *Chromatography and mass spectroscopy in nutrition science and food safety* (1984) pp 249-255