

STRUCTURE NOTE

Cytosolic Domain of the Human Mitochondrial Fission Protein Fis1 Adopts a TPR Fold

Julie A. Dohm,¹ Sarah J. Lee,² J. Marie Hardwick,² R. Blake Hill,^{1,3*} and Apostolos G. Gittis^{4*}¹Department of Biology, Johns Hopkins University, Baltimore, Maryland²Department of Molecular Microbiology and Immunology, Johns Hopkins University, School of Public Health, Baltimore, Maryland³Department of Chemistry, Johns Hopkins University, Baltimore, Maryland⁴Department of Biophysics, Johns Hopkins University, Baltimore, Maryland

Introduction. Mitochondria are polymorphic structures with fundamental roles in cellular differentiation, aging, and apoptosis. Mitochondria accommodate such diverse functions by establishing varying shapes and distributions in the cell via fusion and fission.¹ Thus, the components that mediate fission are likely to be targets for intracellular and extracellular signals that modulate mitochondrial activity in a variety of essential cellular processes.²

Fis1, Mdv1, and Dnm1 are three recently identified proteins essential for the fission of mitochondria in *Saccharomyces cerevisiae*. Mdv1 and Dnm1 assemble in a Fis1-dependent manner on the mitochondrial outer membrane.² Dnm1 is a dynamin-related GTPase that presumably oligomerizes to form rings around the outer surface of the mitochondrial tubules at sites of constriction, similar to structures formed during endocytosis by its homologue dynamin.³ Mdv1 associates with Fis1 and Dnm1 and is thought to act late in the fission process.⁴

The amino acid sequence of Fis1 is conserved from yeast to humans [Supplementary Figure A2], implicating a common mechanism for the control of mitochondrial fission.⁵ Fis1 is an integral membrane protein anchored in the mitochondrial outer membrane by its C-terminal domain. The cytosolic domain of yeast Fis1 has been shown to bind Mdv1 and has been speculated to bind Dnm1 and to self-associate.^{4,5} To help identify the mechanism by which Fis1 mediates interactions with itself and other components of the fission machinery, we determined the crystal structure of the cytosolic domain of human Fis1. Here, we present insights into proposed binding interactions of Fis1 from a 2.0 Å resolution structure of an indispensable component of the mitochondrial fission machinery.

Results. Fis1 is an α -helical protein with two divergent TPR motifs. The crystal structure of Fis1 without its C-terminal domain was determined by multiwavelength anomalous dispersion (MAD) phasing (Table I) and reveals an open array of six α -helices [Fig. 1(A) and Supplementary Figure A1]. The 26-residue helix α 1 is longer than the other helices and extends away from two

TABLE I. Summary of Crystallographic Data and Refinement Statistics

Data collection and phasing			
	λ_1	λ_2	λ_3
Wavelength (Å)	0.9767	0.9797	0.9645
Resolution (Å)	20–2.5	20–2.5	20–2.5
Unique reflections ^a	7860	7890	7912
Completeness (%) ^b	99.6 (99.6)	99.7 (99.9)	99.7 (99.9)
Redundancy	7	7	7
R _{merge} (%) ^c	4.2	4.3	4.4
$\langle I \rangle / \langle \sigma_i \rangle$ ^b	50.6 (26.8)	50.6 (26.8)	50.6 (26.8)
Figure of merit	SOLVE 0.65 RESOLVE 0.79		
Refinement statistics			
Resolution limits (Å)	20–2.0		
R _{cryst} /R _{free} (%) ^e	23.0/25.5		
No. of protein atoms	1010		
No. of nonprotein atoms	75		
RMSD bonds (Å)	0.006		
RMSD angles (°)	1.085		
Average B (Å ²)	34.1		

^aBijvoet pairs were scaled separately for the MAD data sets.^bValues in parentheses are for the highest resolution shell.^c $\sum_i |I(i, h) - \langle I(h) \rangle| / \sum_i I(i, h)$ ^d $\sum_i ||F_o| - |F_c|| / \sum_i |F_o|$ ^eR_{cryst} for 8% of reflections not used during the refinement.

three-helix bundles (α 2– α 4 and α 4– α 6) that share a common helix (α 4). Structural homology searches indicate that Fis1 resembles proteins that possess an α - α super-helix fold, most notably the tetratricopeptide repeat (TPR) motifs. The TPR motif is a well-known protein–protein interaction motif comprised of helix–turn–helix repeats of 34 residues arranged in an antiparallel fashion.⁶ The Fis1 structure superimposes onto the structures of other TPR proteins reasonably well^{7–9} (C α RMSD of \sim 1.5 Å), includ-

The Supplementary Materials Referred to in this article can be found at <http://www.interscience.wiley.com/jpages/0887-3585/suppmat/index.html>

*Correspondence to: Blake Hill, Department of Biology, Johns Hopkins University, 20A Mudd Hall, 3400 N. Charles St., Baltimore, MD 21218. E-mail: hill@jhu.edu and A.G. Gittis, Department of Biophysics, Johns Hopkins University, 007 Jenkins Hall, 3400 N. Charles St., Baltimore, MD 21218. E-mail: apo@groucho.med.jhmi.edu

Received 5 June 2003; Accepted 6 June 2003




Fig. 1. TPR fold of Fis1. **A:** Two ribbon drawings of Fis1 with labeled helices ($\alpha 1$ – $\alpha 6$) and TPR motifs color coded as follows: helix $\alpha 1$ blue, TPR1 ($\alpha 2$ – $\alpha 3$) yellow, TPR2 ($\alpha 4$ – $\alpha 5$) red and helix $\alpha 6$ green. Electron density was not observed for the first two and last two residues. Figures were prepared with MOLSCRIPT²³ and rendered in Raster3d.²⁴ **B:** Structure-based alignment of TPR motifs. The TPR motifs of p67^{phox}, PP5, and PEX5, structural homologs of Fis1 as found by DALI, VAST, and CE, are aligned with the TPR motifs of Fis1. Residues that conform to the TPR consensus sequence below the alignment are highlighted in orange.²⁵ Fis1 α -helical regions are indicated above the alignment by rectangles color-coded as in Figure 1(A). Helix $\alpha 6$ lacks the 34-residue TPR motif and is, therefore, excluded from the alignment. **C:** Stereoview of the final σ_A weighted $2F_o - F_c$ electron density map displaying interactions between labeled residues on helices $\alpha 2$ and $\alpha 4$ calculated to 2.0 Å resolution and contoured at 1.1 σ . Hydrogen bonds are indicated by dashes, and distances are labeled in red. The figure was generated using BOBSCRIPT²⁶ and rendered in Raster3d.²⁴

ing the membrane fusion protein Sec17.¹⁰ However, the amino acid sequence of Fis1 is only distantly related to TPR proteins; little sequence identity was found from sequence database queries using multiple rounds of PSI-BLAST. Structure-based sequence alignments were required to uncover two divergent TPR motifs in Fis1 [Fig. 1(B)]. TPR1 (residues 35–72) differs from the TPR consensus sequence by a three-residue insertion in the loop between helices $\alpha 2$ and $\alpha 3$. It is interesting that this loop is structurally constrained by hydrogen bonds formed between the backbone carbonyl oxygens of two conserved loop residues, Val46 and Thr48, and the guanidinium group of conserved Arg86 in helix $\alpha 4$ [Fig. 1(C)]. Consequently, the orientation of helices $\alpha 2$ and $\alpha 4$ is fixed, potentially defining a binding site. Loops of other TPR proteins were previously shown to mediate protein–protein interactions.¹¹




Fig. 2. Fis1 crystallographic dimer. Ribbon drawing of the Fis1 dimer found in the crystal lattice oriented down the twofold crystallographic symmetry axis. The figure was prepared as in Figure 1(A).

Fis1 forms a dimer in the crystal structure. In Fis1, helices $\alpha 2$, $\alpha 4$, and $\alpha 6$ are arranged to form an amphiphilic, concave surface [Fig. 1(A), right view]. The crystal packing of Fis1 may reveal one manner by which this concave surface mediates binding. In the crystal, a dimer of Fis1 is observed in which the concave surface of a Fis1 molecule makes contacts with helix $\alpha 1$ of a symmetry mate (Fig. 2). The symmetry-related dimer buries ~ 3000 Å² of otherwise exposed surface area on monomeric Fis1 (~ 1500 Å² from helix $\alpha 1$ alone), which supports two-hybrid data that detects yeast Fis1 self-association in vivo.⁶ However, Fis1 also exists as a monomer in solution as determined for the cytosolic domain of yeast Fis1 by size exclusion chromatography (data not shown). The monomeric Fis1 as seen in the crystal structure suggests that Fis1 may exhibit multiple modes of binding that involve the concave surface and helix $\alpha 1$. Indeed, molecular modeling studies suggest that only a few bond angle rotations about backbone ϕ , ψ would be required for an intramolecular association of helix $\alpha 1$. Thus, helix $\alpha 1$ may serve as a “switch” for modulating protein–protein interactions by regulating access to the concave binding surface.

Discussion. Fis1 adopts a TPR fold well-suited for binding interactions necessary for mitochondrial fission. The open structure of Fis1 creates a concave binding surface that can accommodate a helix to potentially promote intermolecular associations (as seen in the crystal packing of the structure) or intramolecular associations (as revealed by modeling). Alternatively, a protein with extended conformation, such as thought for the N-terminal domain of Mdv1, may bind the Fis1 surface in a manner similar to that observed for the extended peptides that bind to HOP, another TPR domain protein that modulates Hsp70/Hsp90 interactions.⁹

Besides the concave binding surface, Fis1 possesses additional structural features for mediating protein–protein interactions. We propose that helix $\alpha 1$ acts as a molecular switch by promoting either monomeric or dimeric conformations of Fis1 to modulate accessibility of different binding surfaces. Furthermore, the loops of Fis1, especially the loop in TPR1, may be important for interactions

with the GTPase Dnm1, as observed in the structure of the TPR domain of p67^{phox} in complex with the GTPase Rac.¹¹ Therefore, Fis1 is structurally equipped to be both free in solution and to associate with itself and with other members of the fission machinery. In addition, the structure of Fis1 is suitable as a scaffold for interactions with proteins involved in other cellular processes such as apoptosis. Several lines of evidence support changes in mitochondrial morphology during caspase-mediated cell death that are consistent with an increase in mitochondrial fission.^{12–17} Thus, Fis1 might serve as a target for signals that mediate cell death in addition to fission.

Materials and Methods. Protein expression, purification, and crystallization. The *Homo sapiens* Fis1 gene (CGI-135) without the C-terminal domain (1–123 aa) was subcloned into the plasmid pET15b and overexpressed in *E. coli* strain pLysS BL21(DE3)codon+ as a His₆-tagged fusion protein. The His₆-tagged protein was purified by Ni²⁺ affinity chromatography, cleaved with biotinylated thrombin, incubated with streptavidin agarose beads and further purified by Ni²⁺ affinity chromatography. The recombinant protein contains three extra vector derived amino acids (Gly-Ser-His) at the N-terminus. Crystals were grown by using hanging drop vapor diffusion at room temperature. Hanging drops consisted of equal volumes of the protein solution (Fis1 at 32 mg mL⁻¹ in 25 mM sodium acetate, 2 mM DTT, pH 5.0) and a reservoir solution (100 mM sodium acetate, 2.0 M ammonium sulfate, 5% glycerol, pH 4.6).

Data collection, structure determination, and refinement. For data collection, crystals were flash-frozen by rapid transfer into mother liquor with 20% (v/v) glycerol followed by direct placement in liquid nitrogen. Multiple three-wavelength MAD data sets were collected at Advanced Photon Source (Argonne, IL) beamline 19-ID on a CCD detector with crystals grown from SeMet-derived protein. All data were processed and scaled with the HKL2000 package.¹⁸ Crystals belong to space group P6₅22 with one molecule in the asymmetric unit ($a = b = 48.9$ Å, $c = 169.1$ Å, $\gamma = 120^\circ$). SOLVE/RESOLVE was used to locate the positions of the selenium atoms and for initial tracing.¹⁹ Reflection intensities were converted to structure factor amplitudes using the CCP4 suite.²⁰ Model building was performed in O²¹ and followed by refinement with CNS.²²

Coordinates. Atomic coordinates have been deposited in the Protein Data Bank (1NZN).

Summary. Fis1 is an integral membrane protein that acts in the fission of mitochondria by controlling the assembly, membrane distribution, and function of the mitochondrial fission machinery. Here we report the 2.0 Å resolution crystal structure of the cytosolic domain of human Fis1. The structure reveals an antiparallel array of α -helices homologous to tetratricopeptide repeat (TPR) proteins. Structure-based sequence alignments of Fis1 uncovered two divergent TPR motifs; the first TPR motif differs from the TPR consensus sequence by a three-

residue insertion in a loop that may be important for function. These TPR helices create an amphiphilic, concave surface that can accommodate a helix or, possibly, an extended segment. Indeed, this putative binding surface mediates homodimer formation of Fis1 in the crystal. The structure of Fis1 provides insight into the architecture of the proposed binding interactions that mediate mitochondrial fission.

ACKNOWLEDGEMENTS

We thank Steve Ginnel for help with data collection, Jeremy Berg, Eaton Lattman, Neil Clarke, Dan Leahy, Doug Barrick, and Scott Walsh for reading the manuscript and helpful discussions. RBH thanks Maria Bewley, John Flanagan, and David Jeruzalmi for helpful discussions. Use of the Argonne National Laboratory Structural Biology Center beamlines at the Advanced Photon Source was supported by the U.S. Department of Energy, Office of Biological and Environmental Research, under Contract No. W-31-109-ENG-38. AGG was partially supported by NSF grant MCB9982967. This work was supported by funds to RBH generously provided by the Zanvyl Krieger School of Arts and Sciences at the Johns Hopkins University.

REFERENCES

1. Bereiter-Hahn J, Voth M. Dynamics of mitochondria in living cells: shape changes, dislocations, fusion, and fission of mitochondria. *Microsc Res Tech* 1994;27(3):198–219.
2. Shaw JM, Nunnari J. Mitochondrial dynamics and division in budding yeast. *Trends Cell Biol* 2002;12(4):178–184.
3. Zhang P, Hinshaw J. Three-dimensional reconstruction of dynamin in the constricted state. *Nature Cell Biol* 2001;3:922–926.
4. Tieu Q, Okreglak V, Naylor K, Nunnari J. The WD repeat protein, Mdv1p, functions as a molecular adaptor by interacting with Dnm1p and Fis1p during mitochondrial fission. *J Cell Biol* 2002;158(August):445–452.
5. Mozdy AD, McCaffery JM, Shaw JM. Dnm1p GTPase-mediated mitochondrial fission is a multi-step process requiring the novel integral membrane component Fis1p. *J Cell Biol* 2000;151(2):367–379.
6. Blatch GL, Lassle M. The tetratricopeptide repeat: a structural motif mediating protein-protein interactions. *Bioessays* 1999;21(11):932–999.
7. Gatto GJ, Geisbrecht BV, Gould SJ, Berg JM. Peroxisomal targeting signal-1 recognition by the TPR domains of human PEX5. *Nat Struct Biol* 2000;7:1091–1095.
8. Das AK, Cohen PTW, Barford D. The structure of the tetratricopeptide repeats of protein phosphatase 5: implications for TPR-mediated protein-protein interactions. *EMBO J* 1998;17(5):1192–1199.
9. Scheufler C, Brinker A, Bourenkov G, Pegoraro S, Moroder L, Bartunik H, Hartl F, Moarefi I. Structure of TPR domain-peptide complexes: critical elements in the assembly of the Hsp70-Hsp90 multichaperone machine. *Cell* 2000;101:199–210.
10. Rice LM, Brunger AT. Crystal structure of the vesicular transport protein Sec17: implications for SNAP function in SNARE complex disassembly. *Mol Cell* 1999;4(July):85–95.
11. Lapouge K, Smith SJM, Walker PA, Gamblin SJ, Rittinger K. Structure of the TPR domain of p67^{phox} in complex with RacGTP. *Mol Cell* 2000;6:899–907.
12. Chen W, Calvo PA, Malide D, Gibbs J, Schubert U, Bacik I, Basta S, O'Neill R, Schickli J, Palese P, Henklein P, Bennink JR, Yewdell JW. A novel influenza A virus mitochondrial protein that induces cell death. *Nat Med* 2001;7(12):1306–1312.
13. Frank S, Gaume B, Bergmann-Leitner E, Leitner W, Robert E, Catez F, Smith C, Youle R. The Role of Dynamin-Related Protein 1, a Mediator of Mitochondrial Fission, in Apoptosis. *Dev Cell* 2001;1(October):515–525.

14. Desagher S, Martinou JC. Mitochondria as the central control point of apoptosis. *Trends Cell Biol* 2000;10(9):369–377.
15. Mancini M, Anderson BO, Caldwell E, Sedghinasab M, Paty PB, Hockenbery DM. Mitochondrial proliferation and paradoxical membrane depolarization during terminal differentiation and apoptosis in a human colon carcinoma cell line. *J Cell Biol* 1997;138(2):449–469.
16. Pinton P, Ferrari D, Rapizzi E, Di Virgilio FD, Pozzan T, Rizzuto R. The Ca^{2+} concentration of the endoplasmic reticulum is a key determinant of ceramide-induced apoptosis: significance for the molecular mechanism of Bcl-2 action. *EMBO J* 2001;20(11):2690–2701.
17. Karbowski M, Lee YJ, Gaume B, Jeong SY, Frank S, Nechushtan A, Santel A, Fuller M, Smith CL, Youle RJ. Spatial and temporal association of Bax with mitochondrial fission sites, Drp1, and Mfn2 during apoptosis. *J Cell Biol* 2002;159(6):931–938.
18. Otwinowski Z, Minor W. Processing of X-ray data collected in oscillation mode. *Methods Enzymol* 1997;276:307–326.
19. Terwilliger TC, Berendzen J. Automated MAD and MIR structure solution. *Acta Crystallogr D* 1999;55:849–861.
20. CCP4. The CCP4 Suite: programs for protein crystallography. *Acta Crystallogr D* 1994;50:760–763.
21. Jones TA, Zou JY, Cowan SW, Kjeldgaard M. Improved methods for building protein models in electron density maps and the location of errors in these models. *Acta Crystallogr A* 1991;47:110–119.
22. Brunger A, Adams P, Clore G, DeLano W, Gros P, Grosse-Kunstleve R, Jiang J, Kuszewski J, Nilges M, Pannu N, Read R, Rice L, Simonson T, Warren G. Crystallography & NMR system: a new software suite for macromolecular structure determination. *Acta Crystallogr D* 1998;54:905–921.
23. Kraulis PJ. MOLSCRIPT: a program to produce both detailed and schematic plots of protein structures. *J Appl Crystallogr* 1991;24:946–950.
24. Merritt EA, Murphy ME. RASTER3D version 2.0. A program for photorealistic molecular graphics. *Acta Crystallogr D* 1994;24:946–950.
25. Sikorski RS, Boguski MS, Goebel M, Hieter P. A repeating amino acid motif in CDC23 defines a family of proteins and a new relationship among genes required for mitosis and RNA synthesis. *Cell* 1990;60(January):307–317.
26. Esnouf RM. An extensively modified version of MolScript that includes greatly enhanced coloring capabilities. *J Mol Graph* 1997;15:132–134.
27. DeLano WL. The PyMOL Molecular Graphics System. San Carlos: DeLano Scientific; 2002.