

# Identification of New Repeating Motifs in Titin

Marion Greaser\*

University of Wisconsin, Muscle Biology Laboratory, Madison, Wisconsin

**ABSTRACT** Repeating motifs of 26–28 amino acids have been identified in the PEVK region of the giant elastic protein titin. These motifs, termed PPAK for the four amino acids that often constitute the beginning of the motif, occur 60 times in human soleus titin. PPAK motifs occur in groups of 2–12 that are separated by regions rich in glutamic acid (approximately 45%) and termed polyE segments. The fluctuation of the net charge between the PPAK and polyE regions suggests ionic interactions between these segments and their involvement in the elastic function of titin. *Proteins* 2001;43:145–149.

© 2001 Wiley-Liss, Inc.

**Key words:** protein domains; elasticity; PEVK; ionic interactions; secondary structure; muscle

## INTRODUCTION

Titin, also called connectin, is a >3-million dalton protein found in heart and skeletal muscle.<sup>1,2</sup> The protein is involved in a number of functions: connecting the thick filaments to the Z-lines,<sup>3</sup> serving as a template for assembly of the thick filaments,<sup>4</sup> preventing the sarcomere from being overstretched,<sup>5,6</sup> acting as a serine/threonine kinase,<sup>7,8</sup> and playing a role in sarcomere assembly.<sup>9–13</sup> It is believed to be the major component responsible for passive tension.<sup>14,15</sup> Additional information on this unusual protein can be found in several recent reviews.<sup>16–19</sup>

Since the earliest titin cDNA sequence became available, it was clear that a number of repeating motifs were involved in its amino acid sequence.<sup>20</sup> Two different types of 100-amino acid motifs were found with similarities to immunoglobulin and the fibronectin III domains. Part of the A-band region of titin also consisted of an 11-motif super-repeat structure.<sup>7</sup> Additional super-repeats have been found in the tandem Ig domains in the I-band section.<sup>21</sup> A 45-amino acid repeat has been identified near the Z line end with varying numbers ( $\geq 7$ ) expressed in different muscles.<sup>22</sup> Shorter serine–proline repeats considered as potential phosphorylation sites have also been found in regions of titin near the M-line<sup>23</sup> and Z-line.<sup>24</sup>

A unique feature of the titin sequence is the PEVK segment in the I-band region.<sup>25</sup> The PEVK is so named because approximately 75% of the amino acid residues are proline (P), glutamic acid (E), valine (V), and lysine (K). The length of the PEVK segment varies between 163 and 2174 residues, with the N2B isoform of the heart having the shorter length and the soleus muscle having the much longer segment.<sup>25</sup> Studies using antibodies to label muscle at various degrees of extension have demonstrated that

PEVK lengthens with stretch.<sup>26,27</sup> Thus, the PEVK region is believed to serve an elastic function in muscle. Although several reports have mentioned sequence repetitions in the PEVK, the nature of such repeats has not been described. The current paper describes two new types of titin repeating sequence that constitute the bulk of the PEVK.

## MATERIALS AND METHODS

A variety of gene and protein analysis software was used in the current study. These include BLAST,<sup>28,29</sup> MEME,<sup>30</sup> MotifSearch,<sup>31</sup> PeptideStructure,<sup>32</sup> PeptideSort, and Pileup. Most of these methods were used through SeqWeb Version 1.1 of the Genetics Computer Group Wisconsin Package Version 10.

## RESULTS AND DISCUSSION

A search of the PEVK sequence for repeating structure was prompted by the observation that a titin monoclonal antibody (9D10)<sup>33</sup> labeled two 0.55- $\mu$ m-wide zones per sarcomere that corresponded to the PEVK region of human soleus titin.<sup>34</sup> Since (1) there are only two sets of titin molecules per sarcomere,<sup>3,35,36</sup> (2) all the molecules are aligned in parallel in each set, and (3) most monoclonals label a zone no wider than 10 nm, the broad staining zone implied that multiple sequence regions were being recognized by the antibody. Numerous examples of the amino acid sequence PPAK were first found in the human soleus titin sequence (GenBank accession number X90569.1<sup>25</sup>) by visual inspection, and these occurred at 26–28 amino acid intervals.<sup>37</sup> A training set consisting of the best 23 sequences was used with the MEME program and a motif width of 28. The MEME output was then used in MotifSearch with the database. A total of 320 matches were recognized in the human soleus titin with this program, but there were numerous instances of overlap, presumably because of the amino acid redundancy and short within-motif sequence repeats. Motif borders were selected and overlap sequences eliminated with the help of the position *P*-values.<sup>30</sup> Motif positions 3–6 (typically KVP) were also useful in constructing a final alignment since this area of the motif showed the most limited variability.

Grant sponsor: College of Agricultural and Life Sciences, University of Wisconsin–Madison; Grant sponsor: National Institutes of Health; Grant number: HL62466.

\*Correspondence to: Marion Greaser, University of Wisconsin, Muscle Biology Laboratory, Madison, Wisconsin. E-mail: mgreaser@facstaff.wisc.edu

Received 17 July 2000; Accepted 8 December 2000



also found in other regions, but more typical structures are 2–3 consecutive glutamic acids alternating with one or two hydrophobic and/or proline residues. The following example begins at residue 6398: (**EEEAVSVQREEEY-EEYEEYDYKEFEEYEPTTEEYDQYEEY**). Often a single aromatic residue is sandwiched between two or three glutamic acids on both sides. There are also several regions where the glutamic acid residues alternate with other amino acids.

BLAST<sup>28,29</sup> and MotifSearch<sup>31</sup> searches were conducted to determine whether motifs similar to those found in the human soleus titin PEVK occurred in other proteins. The only similarity of sequence between the PEVK region and sequence patterns in the PROSITE database was with the proline-rich region found in keratinocyte proteins. The sequence KVPE occurs in several types of short motifs in these proteins (see <http://interpro.ebi.ac.uk> entry IPR000694), but this protein group is typically high in cysteine and glutamine, amino acids that are virtually absent from the titin PEVK. A few PPAK-like domains can be identified in the partial chicken skeletal titin (connectin) sequence,<sup>38</sup> but the latter sequence is partial, extending only a short distance into the PEVK region. There appears to be some similarity of the PPAK sequence to the first 27 residues of a myosin alkali light chain<sup>39</sup> (1MPP-KKKEPKKAPEPKKE EPK PAKPAEP28). A PolyE-like sequence is also found at the N-terminus of troponin T<sup>40</sup> (1MSDTEEQYEEEQPEEEAAEEEEEAPEEPEPVAE-PEEERP KP41). In addition, neurofilament H<sup>41</sup> has weak sequence similarities to both the PPAK and polyE segments. The 9D10 titin monoclonal antibody cross-reacts with this latter protein on Western blots<sup>37</sup>; other workers have identified titin monoclonals that react with neurofilament H as well.<sup>42–44</sup> These observations suggest that the two proteins may assume some common folding arrangement. The greatest similarity between the titin PEVK and neurofilament H is the repetition of lysines (either single or pairs) spaced every 6–8 residues in both proteins.

Secondary structure predictions using PeptideStructure<sup>32</sup> for the PPAK repeat regions provided limited insight on the protein folding in the PEVK region. The high proline content precludes significant  $\alpha$ -helix or typical  $\beta$ -sheet structure. However, predictions with the polyE segments consistently indicated considerable  $\alpha$ -helical content. The significant number of prolines present would prevent the polyE segments from assuming a completely straight helical orientation. Regions rich in prolines often contain the polyproline II structure,<sup>45</sup> but the PPAK repeats lack the glutamine residues often found in such structures.<sup>46</sup> Further experimental work using circular dichroism (CD) or nuclear magnetic resonance (NMR), or both, on synthetic or expressed peptides will be necessary to better define the protein secondary structure in the PEVK region.

The transitions back and forth between the tandem PPAK-containing segments and the polyE regions result in a cycling of the isoelectric points between these regions. Figure 2 shows that the pI values fluctuate between near 10 for the PPAK segments to ~3–4 for the polyE regions.




Fig. 2. Fluctuations in isoelectric point along the human soleus PEVK. Values for the different segments were obtained using PeptideSort.

The isoelectric points of the individual PPAK repeats are mostly in the 9–10 region, but a few have pI values of <7. The more negatively charged PPAKs tend to occur in the middle of the wider PPAK clusters, so there are more charge fluctuations than depicted in Figure 2. The isoelectric point of the PEVK region has previously been shown to be more negative than most of the rest of the titin molecule<sup>47</sup>; the calculated value for the human soleus PEVK is 5.06. The PPAK repeats constitute 75% of the amino acid number in the PEVK sequence while the polyE makes up 25%. Thus, the much smaller proportion polyE sequence dominates the charge for the PEVK region.

It is postulated that there may be charge–charge interaction between the positively charged PPAKs and the negatively charged polyE segments. Previous work has shown that there is an ionic strength effect on the stiffness of rat psoas myofibrils.<sup>27</sup> These workers suggested that this effect occurred in the PEVK region. Further work will be needed to validate or refute this hypothesis.

Elastic changes in other proteins upon stretch have been ascribed to several types of repeating amino acid patterns. The extensibility of the protein elastin is believed to involve a  $\beta$  spiral to extended chain transition of numerous repeats with the sequence VPGVG.<sup>48–50</sup> A similar  $\beta$  spiral produced from repetitive  $\beta$  turns is believed to occur in wheat gluten with GYYTSPQQ and PGQGQQ repeat sequences.<sup>51</sup> Although glycines appear to be important in the flexibility of these structures, this amino acid is rare among the PPAK domains (Table I). A recent study of heptads with a sequence of YSPTSPS from RNA polymerase II<sup>52</sup> showed that polymers with this repeat contain both  $\beta$  turn and polyproline II helix structure. Proline, glycine, glutamine, serine, and threonine, are amino acids commonly found in  $\beta$  turn regions,<sup>53</sup> none of the latter four occurs in the PPAK consensus sequence (Table I). Thus,



it is unclear whether similar mechanisms of reversible folding–unfolding occur in titin.

Many of the PPAK repeats may arise from exon duplications. A family of 27 amino acid segments derived from single exons found in titin genomic DNA (many of which are not expressed in human soleus titin) was recently reported.<sup>54</sup> The start site for these exons does not align with the PPAK repeats; rather, it corresponds to the PPAK consensus valine at position 5. The 27-amino acid repeat<sup>54</sup> really should have 28 amino acids in the protein, but the highly conserved valine is missing because of incomplete codons at both ends of each exon. Although the PPAK repeats described above could be realigned to the exon start sites, beginning the motif with the position 1 proline appears to be more appropriate from a protein structure standpoint since it better separates the PEVK sequence into regions of different charge.

A number of different titin isoforms are expressed by alternative splicing.<sup>25,54</sup> Splicing removes all the polyE segments in the human cardiac N2B isoform (Fig. 1) plus all but 5 of the 60 PPAK repeats found in the soleus titin isoform. The cardiac N2A still has two polyE domains and 20 PPAK motifs. Other titin isoforms with varying PEVK length also occur in skeletal and cardiac muscle.<sup>25,54</sup> The lack of a polyE segment in the PEVK region of the N2B isoform suggests that interaction of the PPAK and polyE is not essential for the folding and unfolding of the PEVK region in this isoform. Further structural studies will be necessary to determine if the PPAK and polyE repeats interact under physiological conditions and to define their role in the elastic function of titin.

## ACKNOWLEDGMENTS

The author thanks Dr. Jean-Yves Sgro for assistance with the GCG programs and Dr. John Trinick for comments and suggestions on an earlier version of this manuscript.

## REFERENCES

- Wang K, McClure J, Tu A. Titin: major myofibrillar components of striated muscle. *Proc Natl Acad Sci USA* 1979;76:3698–3702.
- Maruyama K. Connectin, an elastic protein from myofibrils. *J Biochem (Tokyo)* 1976;80:405–407.
- Furst DO, Osborn M, Nave R, Weber K. The organization of titin filaments in the half-sarcomere revealed by monoclonal antibodies in immunoelectron microscopy: a map of ten nonrepetitive epitopes starting at the Z line extends close to the M line. *J Cell Biol* 1988;106:1563–1572.
- Trinick J. Titin and nebulin: protein rulers in muscle? *Trends Biochem Sci* 1994;19:405–409.
- Horowitz R, Kempner ES, Bisher ME, Podolsky RJ. A physiological role for titin and nebulin in skeletal muscle. *Nature* 1986;323:160–164.
- Wang K, McCarter R, Wright J, Beverly J, Ramirez-Mitchell R. Regulation of skeletal muscle stiffness and elasticity by titin isoforms: a test of the segmental extension model of resting tension. *Proc Natl Acad Sci USA* 1991;88:7101–7105.
- Labeit S, Gautel M, Lakey A, Trinick J. Towards a molecular understanding of titin. *EMBO J* 1992;11:1711–1716.
- Mayans O, van der Ven PF, Wilm M, Mues A, Young P, Furst DO, Wilmanns M, Gautel M. Structural basis for activation of the titin kinase domain during myofibrillogenesis. *Nature* 1998;395:863–869.
- Wang SM, Greaser ML, Schultz E, Bulinski JC, Lin JJ, Lessard JL. Studies on cardiac myofibrillogenesis with antibodies to titin, actin, tropomyosin, and myosin. *J Cell Biol* 1988;107:1075–1083.
- Schultheiss T, Lin ZX, Lu MH, Murray J, Fischman DA, Weber K, Masaki T, Imamura M. Differential distribution of subsets of myofibrillar proteins in cardiac nonstriated and striated myofibrils. *J Cell Biol* 1990;110:1159–1172.
- Handel SE, Greaser ML, Schultz E, Wang SM, Bulinski JC, Lin JJ, Lessard JL. Chicken cardiac myofibrillogenesis studied with antibodies specific for titin and the muscle and nonmuscle isoforms of actin and tropomyosin. *Cell Tissue Res* 1991;263:419–430.
- Rhee D, Sanger JM, Sanger JW. The premyofibril: evidence for its role in myofibrillogenesis. *Cell Motil Cytoskeleton* 1994;28:1–24.
- Gregorio CC, Trombitas K, Centner T, Kolmerer B, Stier G, Kunke K, Suzuki K, Obermayr F, Herrmann B, Granzier H, Sorimachi H, Labeit S. The NH2 terminus of titin spans the Z-disc: its interaction with a novel 19-kD ligand (T-cap) is required for sarcomeric integrity. *J Cell Biol* 1998;143:1013–1027.
- Granzier HL, Wang K. Passive tension and stiffness of vertebrate skeletal and insect flight muscles: the contribution of weak cross-bridges and elastic filaments. *Biophys J* 1993;65:2141–2159.
- Granzier HL, Irving TC. Passive tension in cardiac muscle: contribution of collagen, titin, microtubules, and intermediate filaments. *Biophys J* 1995;68:1027–1044.
- Labeit S, Kolmerer B, Linke WA. The giant protein titin. Emerging roles in physiology and pathophysiology. *Circ Res* 1997;80:290–294.
- Maruyama K. Connectin/titin, giant elastic protein of muscle. *FASEB J* 1997;11:341–345.
- Trinick J, Tskhovrebova L. Titin: a molecular control freak. *Trends Cell Biol* 1999;9:377–380.
- Gregorio CC, Granzier H, Sorimachi H, Labeit S. Muscle assembly: a titanic achievement? *Curr Opin Cell Biol* 1999;11:18–25.
- Labeit S, Barlow DP, Gautel M, Gibson T, Holt J, Hsieh CL, Francke U, Leonard K, Wardale J, Whiting A. A regular pattern of two types of 100-residue motif in the sequence of titin. *Nature* 1990;345:273–276.
- Gautel M. The super-repeats of titin/connectin and their interactions: glimpses at sarcomeric assembly. *Adv Biophys* 1996;33:27–37.
- Gautel M, Goulding D, Bullard B, Weber K, Furst DO. The central Z-disk region of titin is assembled from a novel repeat in variable copy numbers. *J Cell Sci* 1996;109:2747–2754.
- Gautel M, Leonard K, Labeit S. Phosphorylation of KSP motifs in the C-terminal region of titin in differentiating myoblasts. *EMBO J* 1993;12:3827–3834.
- Besteyen MG, Wolff JA, Greaser ML. Characterization of a 5.4 kb cDNA fragment from the Z-line region of rabbit cardiac titin reveals phosphorylation sites for proline-directed kinases. *J Cell Sci* 1995;108:3029–3037.
- Labeit S, Kolmerer B. Titins: giant proteins in charge of muscle ultrastructure and elasticity. *Science* 1995;270:293–296.
- Trombitas K, Greaser M, Labeit S, Jin JP, Kellermayer M, Helmes M, Granzier H. Titin extensibility in situ: entropic elasticity of permanently folded and permanently unfolded molecular segments. *J Cell Biol* 1998;140:853–859.
- Linke WA, Ivemeyer M, Mundel P, Stockmeier MR, Kolmerer B. Nature of PEVK-titin elasticity in skeletal muscle. *Proc Natl Acad Sci USA* 1998;95:8052–8057.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol* 1990;215:403–410.
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 1997;25:3389–3402.
- Bailey TL, Gribskov M. Score distributions for simultaneous matching to multiple motifs. *J Comput Biol* 1997;4:45–59.
- Gribskov M, Luthy R, Eisenberg D. Profile analysis. *Methods Enzymol* 1990;183:149–159.
- Jameson B, Wolf H. The antigenic index: a novel algorithm for predicting antigenic determinants. *CABIOS* 1988;4:181–186.
- Wang SM, Greaser ML. Immunocytochemical studies using a monoclonal antibody to bovine cardiac titin on intact and extracted myofibrils. *J Muscle Res Cell Motil* 1985;6:293–312.
- Trombitas K, Greaser M, French G, Granzier H. PEVK extension of human soleus muscle titin revealed by immunolabeling with the anti-titin antibody 9D10. *J Struct Biol* 1998;122:188–196.
- Itoh Y, Suzuki T, Kimura S, Ohashi K, Higuchi H, Sawada H, Shimizu T, Shibata M, Maruyama K. Extensible and less-

- extensible domains of connectin filaments in stretched vertebrate skeletal muscle sarcomeres as detected by immunofluorescence and immunoelectron microscopy using monoclonal antibodies. *J Biochem (Tokyo)* 1988;104:504–508.
36. Whiting A, Wardale J, Trinick J. Does titin regulate the length of muscle thick filaments? *J Mol Biol* 1989;205:263–268.
  37. Greaser M, Wang S-M, Berri M, Mozdziak PE, Kumazawa Y. Sequence and mechanical implications of cardiac PEVK. *Adv. Exp. Med. Biol.* 2000;481:53–66.
  38. Maruyama K, Endo T, Kume H, Kawamura Y, Kanzawa N, Nakauchi Y, Kimura S, Kawashima S. A novel domain sequence of connectin localized at the I band of skeletal muscle sarcomeres: homology to neurofilament subunits. *Biochem Biophys Res Commun* 1993;194:1288–1291.
  39. Nakamura S, Nabeshima YI, Kobayashi H, Nabeshima Y, Nonomura Y, Fujii-Kuriyama Y. Single chicken cardiac myosin alkali light chain gene generates two different mRNAs by alternative splicing of a complex exon. *J Mol Biol* 1988;203:895–904.
  40. Gahlmann R, Troutt AB, Wade RP, Gunning P, Kedes L. Alternative splicing generates variants in important functional domains of human slow skeletal troponin T. *J Biol Chem* 1987;262:16122–16126.
  41. Robinson PA, Wion D, Anderton BH. Isolation of a cDNA for the rat heavy neurofilament polypeptide (NF-H). *FEBS Lett* 1986;209:203–205.
  42. Shimizu T, Matsumura K, Itoh Y, Mannen T, Maruyama K. An immunological homology between neurofilament and muscle elastic filament: a monoclonal antibody cross-reacts with neurofilament subunits and connectin. *Biomed Res* 1988;9:227–233.
  43. Matsumura K, Shimizu T, Mannen T, Maruyama K. The immunological homology between two filamentous cross-linker phosphoproteins, connectin and cross-bridge region of neurofilament-H, is not affected by the phosphorylation state. *J Biochem (Tokyo)* 1989;105:226–230.
  44. Mencarelli C, Magi B, Marzocchi B, Armellini D, Pallini V. Evolution of the “titin epitope” in neurofilament proteins. *Comp Biochem Physiol [B]* 1991;100:741–744.
  45. Williamson MP. The structure and function of proline-rich regions of proteins. *Biochem J* 1994;297:249–260.
  46. Stapley BJ, Creamer TP. A survey of left-handed polyproline II helices. *Protein Sci* 1999;8:587–595.
  47. Tskhovrebova L, Trinick J. Direct visualization of extensibility in isolated titin molecules. *J Mol Biol* 1997;265:100–106.
  48. Urry DW, Haynes B, Zhang H, Harris RD, Prasad KU. Mechanochemical coupling in synthetic polypeptides by modulation of an inverse temperature transition. *Proc Natl Acad Sci USA* 1988;85:3407–3411.
  49. Urry DW. Entropic elastic processes in protein mechanisms. I. Elastic structure due to an inverse temperature transition and elasticity due to internal chain dynamics. *J Protein Chem* 1988;7:1–34.
  50. Reiersen H, Clarke AR, Rees AR. Short elastin-like peptides exhibit the same temperature-induced structural transitions as elastin polymers: implications for protein engineering. *J Mol Biol* 1998;283:255–264.
  51. Field JM, Tatham AS, Shewry PR. The structure of a high- $M_r$  subunit of durum-wheat (*Triticum durum*) gluten. *Biochem J* 1987;247:215–221.
  52. Bienkiewicz EA, Woody A-YM, Woody RW. Conformation of the RNA polymerase II C-terminal domain: circular dichroism of long and short fragments. *J Mol Biol* 2000;297:119–133.
  53. Hutchinson EG, Thornton JM. A revised set of potentials for beta-turn formation in proteins. *Protein Sci* 1994;3:2207–2216.
  54. Freiburg A, Trombitas K, Hell W, Cazorla O, Fougereousse F, Centner T, Kolmerer B, Witt C, Beckmann JS, Gregorio CC, Granzier H, Labeit S. Series of exon-skipping events in the elastic spring region of titin as the structural basis for myofibrillar elastic diversity. *Circ Res* 2000;86:1114–1121.