Two-row recurrence for Koornwinder polynomials

Houcine Ben Dali
Harvard University

joint work with Lauren Williams

Conférence en l'honneur de Philippe Biane

Paris, Sorbonne Université

October 19, 2026

Koornwinder polynomials: overview


Koornwinder polynomials are Laurent polynomials in n variables $z_1, \ldots, z_n$, depending on six parameters q, t, a, b, c, d.

- Koornwinder '92: symmetric polynomials, indexed by partitions, introduced as multivariable orthogonal polynomials.
- Cherednik '95: introduced double affine Hecke algebras of reduced types.
- Sahi '99, '00: introduced the double affine Hecke algebra of type $(C^\vee, C)$ and used it to define the nonsymmetric Koornwinder polynomials.
- Stokman '00: orthogonality of nonsymmetric Koornwinder polynomials.

Koornwinder polynomials: overview

Koornwinder polynomials are Laurent polynomials in n variables $z_1, \ldots, z_n$, depending on six parameters q, t, a, b, c, d.

- Koornwinder '92: symmetric polynomials, indexed by partitions, introduced as multivariable orthogonal polynomials.
- Cherednik '95: introduced double affine Hecke algebras of reduced types.
- Sahi '99, '00: introduced the double affine Hecke algebra of type $(C^\vee, C)$ and used it to define the nonsymmetric Koornwinder polynomials.
- Stokman '00: orthogonality of nonsymmetric Koornwinder polynomials.




Macdonald '03

Koornwinder polynomials: overview

Koornwinder polynomials are Laurent polynomials in n variables $z_1, \ldots, z_n$, depending on six parameters q, t, a, b, c, d.


- Koornwinder '92: symmetric polynomials, indexed by partitions, introduced as multivariable orthogonal polynomials.
- Cherednik '95: introduced double affine Hecke algebras of reduced types.
- Sahi '99, '00: introduced the double affine Hecke algebra of type $(C^\vee, C)$ and used it to define the nonsymmetric Koornwinder polynomials.
- Stokman '00: orthogonality of nonsymmetric Koornwinder polynomials.



Macdonald '03

$\Longrightarrow$ They generalize several families from symmetric function theory.

Koornwinder polynomials: overview



♦ Koornwinder '92, Macdonald '03; detailed treatment: Yamaguchi–Yanagida '23.

* Symmetric case: van Diejen '95. Nonsymmetric case: BD–Williams '26.

Motivations and previous results

- **Combinatorics.** Type A Macdonald polynomials have several combinatorial formulas: fillings (Haglund–Haiman–Loehr '05) and multiline queues (Corteel–Mandelshtam–Williams '22).
  - ▶ Generalize these formulas to Koornwinder polynomials?

Motivations and previous results

- **Combinatorics.** Type A Macdonald polynomials have several combinatorial formulas: fillings (Haglund–Haiman–Loehr '05) and multiline queues (Corteel–Mandelshtam–Williams '22).
  ▶ Generalize these formulas to Koornwinder polynomials?
  Alcove walk formulas (Ram–Yip '11, Colmenarejo–Gagnon–Ram '26); the case $\lambda = (1, \ldots, 1, 0, \ldots, 0)$: rhombic staircase tableaux (Corteel–Mandelshtam–Williams '24).

Motivations and previous results

- **Combinatorics.** Type A Macdonald polynomials have several combinatorial formulas: fillings (Haglund–Haiman–Loehr '05) and multiline queues (Corteel–Mandelshtam–Williams '22).
  ▶ Generalize these formulas to Koornwinder polynomials?
  Alcove walk formulas (Ram–Yip '11, Colmenarejo–Gagnon–Ram '26); the case $\lambda = (1, \ldots, 1, 0, \ldots, 0)$: rhombic staircase tableaux (Corteel–Mandelshtam–Williams '24).

- **Integrability.** Vertex models for nonsymmetric Macdonald polynomials (Borodin–Wheeler '22); only special cases for Koornwinder (Wheeler–Zinn-Justin '16).
  ▶ A vertex model for Koornwinder polynomials?

Motivations and previous results

- **Combinatorics.** Type A Macdonald polynomials have several combinatorial formulas: fillings (Haglund–Haiman–Loehr '05) and multiline queues (Corteel–Mandelshtam–Williams '22).
  ▶ Generalize these formulas to Koornwinder polynomials?
  Alcove walk formulas (Ram–Yip '11, Colmenarejo–Gagnon–Ram '26); the case $\lambda = (1, \ldots, 1, 0, \ldots, 0)$: rhombic staircase tableaux (Corteel–Mandelshtam–Williams '24).

- **Integrability.** Vertex models for nonsymmetric Macdonald polynomials (Borodin–Wheeler '22); only special cases for Koornwinder (Wheeler–Zinn-Justin '16).
  ▶ A vertex model for Koornwinder polynomials?

- **Interacting particle systems.** At $q = 1$ and $z_i = 1$: connection to the multispecies open ASEP (Cantini–Garbali–de Gier–Wheeler '16).

Main results (BD–Williams '26)

Theorem A (two-row recurrence)

Koornwinder polynomials satisfy a recurrence relation.

Theorem B (local rules)

The coefficients of this recurrence are characterized by a finite family of local rules.

Theorem C (top homogeneous part)

The top homogeneous parts of nonsymmetric Koornwinder polynomials are nonsymmetric type A Macdonald polynomials.

Notation


Fix $n > 0$.

- A partition is $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n \geq 0)$, and a signed composition is $\kappa = (\kappa_1, \ldots, \kappa_n) \in \mathbb{Z}^n$.

Notation

Fix $n > 0$.

- A partition is $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n \geq 0)$, and a signed composition is $\kappa = (\kappa_1, \ldots, \kappa_n) \in \mathbb{Z}^n$.
- The finite Weyl group $W_0 \simeq S_n \ltimes (\mathbb{Z}/2\mathbb{Z})^n$ of type C_n is the group of signed permutations. It acts on $\mathbb{Z}^n$ by permuting the entries and changing their signs.



Notation

Fix $n > 0$.

- A partition is $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n \geq 0)$, and a signed composition is $\kappa = (\kappa_1, \ldots, \kappa_n) \in \mathbb{Z}^n$.
- The finite Weyl group $W_0 \simeq S_n \ltimes (\mathbb{Z}/2\mathbb{Z})^n$ of type C_n is the group of signed permutations. It acts on $\mathbb{Z}^n$ by permuting the entries and changing their signs.
- $W_0 \cdot \lambda$ is the set of signed compositions obtained by permuting the parts of λ and changing their signs. For example,

$$W_0 \cdot (2, 0) = \{(2, 0), (-2, 0), (0, 2), (0, -2)\}.$$

Notation

Fix $n > 0$.

- A partition is $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n \geq 0)$, and a signed composition is $\kappa = (\kappa_1, \ldots, \kappa_n) \in \mathbb{Z}^n$.
- The finite Weyl group $W_0 \simeq S_n \ltimes (\mathbb{Z}/2\mathbb{Z})^n$ of type C_n is the group of signed permutations. It acts on $\mathbb{Z}^n$ by permuting the entries and changing their signs.
- $W_0 \cdot \lambda$ is the set of signed compositions obtained by permuting the parts of λ and changing their signs. For example,

$$W_0 \cdot (2, 0) = \{(2, 0), (-2, 0), (0, 2), (0, -2)\}.$$

- We work with Laurent polynomials in $\mathbb{Q}(q, t, a, b, c, d)[z_1^{\pm 1}, \ldots, z_n^{\pm 1}]$. The group W_0 acts by permuting the variables and inverting them, $z_i \mapsto z_i^{-1}$.

Notation

Fix $n > 0$.

- A partition is $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n \geq 0)$, and a signed composition is $\kappa = (\kappa_1, \ldots, \kappa_n) \in \mathbb{Z}^n$.
- The finite Weyl group $W_0 \simeq S_n \ltimes (\mathbb{Z}/2\mathbb{Z})^n$ of type C_n is the group of signed permutations. It acts on $\mathbb{Z}^n$ by permuting the entries and changing their signs.
- $W_0 \cdot \lambda$ is the set of signed compositions obtained by permuting the parts of λ and changing their signs. For example,

$$W_0 \cdot (2, 0) = \{(2, 0), (-2, 0), (0, 2), (0, -2)\}.$$

- We work with Laurent polynomials in $\mathbb{Q}(q, t, a, b, c, d)[z_1^{\pm 1}, \ldots, z_n^{\pm 1}]$. The group W_0 acts by permuting the variables and inverting them, $z_i \mapsto z_i^{-1}$.
- A Laurent polynomial is symmetric if it is invariant under W_0.

Notation

Fix $n > 0$.

- A partition is $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n \geq 0)$, and a signed composition is $\kappa = (\kappa_1, \ldots, \kappa_n) \in \mathbb{Z}^n$.
- The finite Weyl group $W_0 \simeq S_n \ltimes (\mathbb{Z}/2\mathbb{Z})^n$ of type C_n is the group of signed permutations. It acts on $\mathbb{Z}^n$ by permuting the entries and changing their signs.
- $W_0 \cdot \lambda$ is the set of signed compositions obtained by permuting the parts of λ and changing their signs. For example,

$$W_0 \cdot (2, 0) = \{(2, 0), (-2, 0), (0, 2), (0, -2)\}.$$

- We work with Laurent polynomials in $\mathbb{Q}(q, t, a, b, c, d)[z_1^{\pm 1}, \ldots, z_n^{\pm 1}]$. The group W_0 acts by permuting the variables and inverting them, $z_i \mapsto z_i^{-1}$.
- A Laurent polynomial is symmetric if it is invariant under W_0.

Example

$z_1^2 + z_1^{-2} + z_2^2 + z_2^{-2} - bd\left(z_1 z_2 + z_1 z_2^{-1} + z_1^{-1} z_2 + z_1^{-1} z_2^{-1}\right)$ is symmetric.

Notation

Fix $n > 0$.

- A partition is $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n \geq 0)$, and a signed composition is $\kappa = (\kappa_1, \ldots, \kappa_n) \in \mathbb{Z}^n$.
- The finite Weyl group $W_0 \simeq S_n \ltimes (\mathbb{Z}/2\mathbb{Z})^n$ of type C_n is the group of signed permutations. It acts on $\mathbb{Z}^n$ by permuting the entries and changing their signs.
- $W_0 \cdot \lambda$ is the set of signed compositions obtained by permuting the parts of λ and changing their signs. For example,


$$W_0 \cdot (2, 0) = \{(2, 0), (-2, 0), (0, 2), (0, -2)\}.$$

- We work with Laurent polynomials in $\mathbb{Q}(q, t, a, b, c, d)[z_1^{\pm 1}, \ldots, z_n^{\pm 1}]$. The group W_0 acts by permuting the variables and inverting them, $z_i \mapsto z_i^{-1}$.
- A Laurent polynomial is symmetric if it is invariant under W_0.


Indexing:

- ▷ symmetric polynomials: partitions $\lambda = (\lambda_1, \ldots, \lambda_n)$,
- ▷ nonsymmetric Laurent polynomials: signed compositions $\kappa \in \mathbb{Z}^n$.

Two nonsymmetric refinements of the symmetric Koornwinder polynomials



Two nonsymmetric refinements of the symmetric Koornwinder polynomials



Proposition (Cantini–Garbali–de Gier–Wheeler '16)

$$K_\lambda = \sum_{\kappa \in W_0 \cdot \lambda} F_\kappa.$$

Example

$K_{(2,0)} = F_{(2,0)} + F_{(-2,0)} + F_{(0,2)} + F_{(0,-2)}$.

Definitions of E_κ and F_κ

The affine Hecke algebra of type $\widetilde{C}_n$ is generated by $T_0, \ldots, T_n$, with braid relations given by the diagram and $(T_i - t_i)(T_i + 1) = 0$.



Definitions of E_κ and F_κ

The affine Hecke algebra of type $\widetilde{C}_n$ is generated by $T_0, \ldots, T_n$, with braid relations given by the diagram and $(T_i - t_i)(T_i + 1) = 0$.



It acts on Laurent polynomials via the polynomial representation.

Definitions of E_κ and F_κ

The affine Hecke algebra of type $\widetilde{C}_n$ is generated by $T_0, \ldots, T_n$, with braid relations given by the diagram and $(T_i - t_i)(T_i + 1) = 0$.



It acts on Laurent polynomials via the polynomial representation.

E_κ: eigenfunctions of the Cherednik operators (Sahi '99, '00)

$$Y_i = T_i \cdots T_{n-1} T_n T_{n-1} \cdots T_1 \\ T_0 T_1^{-1} \cdots T_{i-1}^{-1},$$

$$Y_i E_\kappa = y_i(\kappa)\, E_\kappa,$$

for $1 \le i \le n$, normalized by $[z^\kappa] E_\kappa = 1$.

Definitions of E_κ and F_κ

The affine Hecke algebra of type $\widetilde{C}_n$ is generated by $T_0, \ldots, T_n$, with braid relations given by the diagram and $(T_i - t_i)(T_i + 1) = 0$.



It acts on Laurent polynomials via the polynomial representation.

E_κ: eigenfunctions of the Cherednik operators (Sahi '99, '00)

$$Y_i = T_i \cdots T_{n-1} T_n T_{n-1} \cdots T_1 \, T_0 T_1^{-1} \cdots T_{i-1}^{-1},$$

$$Y_i E_\kappa = y_i(\kappa)\, E_\kappa,$$

for $1 \le i \le n$, normalized by $[z^\kappa] E_\kappa = 1$.


F_κ: the unique family satisfying the qKZ relations (Kasatani '11), normalized by $[z^\kappa] F_\kappa = 1$:

$$\begin{aligned}
T_0 F_{\kappa_1,\ldots} &= q^{\kappa_1} F_{-\kappa_1,\ldots}, && \kappa_1 < 0,\\
T_0 F_{\kappa_1,\ldots} &= t_0\, F_{\kappa_1,\ldots}, && \kappa_1 = 0,\\
T_i F_{\ldots,\kappa_i,\kappa_{i+1},\ldots} &= t\, F_{\ldots,\kappa_i,\kappa_{i+1},\ldots}, && \kappa_i = \kappa_{i+1},\\
T_i F_{\ldots,\kappa_i,\kappa_{i+1},\ldots} &= F_{\ldots,\kappa_{i+1},\kappa_i,\ldots}, && \kappa_i > \kappa_{i+1},\\
T_n F_{\ldots,\kappa_n} &= t_n\, F_{\ldots,\kappa_n}, && \kappa_n = 0,\\
T_n F_{\ldots,\kappa_n} &= F_{\ldots,-\kappa_n}, && \kappa_n > 0.
\end{aligned}$$

Koornwinder polynomials and the multispecies ASEP

Theorem (Cantini–Garbali–de Gier–Wheeler '16)

The stationary probability of a state μ of the multispecies ASEP with open boundaries on n sites is proportional to
$F_\mu(z_1 = \cdots = z_n = 1; q = 1, t, a, b, c, d)$.



Dynamics of the multispecies ASEP with open boundaries

Here a, b, c, d are related to the boundary rates by

$$\alpha = \frac{-ac(1-t)}{(1+a)(1+c)}, \qquad \gamma = \frac{1-t}{(1+a)(1+c)},$$

$$\beta = \frac{-bd(1-t)}{(1+b)(1+d)}, \qquad \delta = \frac{1-t}{(1+b)(1+d)}.$$

Some examples of Koornwinder ASEP polynomials for $n = 3$

The top degree part is represented in red; we will see that it corresponds to type A ASEP polynomials.

$$F_{(0,1,0)} = z_2 + \frac{t\big(t^2ac(b+d) - a - c\big)}{1 - t^4abcd}$$

$$\begin{aligned}
F_{(0,2,0)} = {} & z_2^2 + \frac{1-t}{1-qt}\,(z_1 + qz_3)\,z_2 + \frac{q(1-t)}{1-qt}\,(z_1^{-1} + z_3^{-1})\,z_2 + \frac{qt\big(qt^2ac(b+d) - a - c\big)}{1 - q^2t^4abcd}\,z_2^{-1} \\
& + \frac{qt(1-t)\big(qt^2ac(b+d) - a - c\big)}{(1-qt)(1-q^2t^4abcd)}\,\big(z_1 + q\,z_1^{-1} + z_3 + z_3^{-1}\big) \\
& - \frac{1}{(1-qt)(1-q^2t^4abcd)}\Big(q^3t^5\big(ac(b+d) + bd(a+c)\big) + q^2t^4ac(b+d) - q^2t^3ac(b+d) \\
& \quad - q^2t^2bd(a+c) - q^2t^3(a+b+c+d) - qt^2ac(b+d) - qt^2(a+c) \\
& \quad + qt(a+c) + q(b+d) + a + c\Big)\,z_2 \\
& - \frac{1}{(1-qt)(1-qt^4abcd)(1-q^2t^4abcd)}\Big(q^4t^6abcd\big(t^3abcd + t^2(a+c)(b+d) \\
& \quad - t^2abcd + t^2ac + tabcd - 1\big) \\
& \quad + q^3t^4\big(t^4a^2c^2(b+d)^2 - t^4a^2bc^2d - t^3a^2b^2c^2d^2 - t^2bd(a+c)^2 - t^2ac(a+c)(b+d) \\
& \quad - t^2ac(b+d)^2 - tabcd(a+c)(b+d) + t^2abcd - ta^2bc^2d - tabcd + abcd\big) \\
& \quad - q^2t^2\big(t^4ac(a+c)(b+d) + t^3a^2c^2(b+d)^2 - t^3a^2bc^2d + t^3abcd \\
& \quad - t^2(a+c)^2 - t^2(a+c)(b+d) - t^2abcd - tbd(a+c)^2 - tac(a+c)(b+d) \\
& \quad - tac(b+d)^2 + t^2ac + tabcd - 1\big) \\
& \quad + q\big(t^3ac(a+c)(b+d) + t^4ac + t^3abcd - t(a+c)^2 - t(a+c)(b+d) \\
& \quad - t^2 + tac + t - 1\big) - tac\Big)
\end{aligned}$$

Some examples of Koornwinder ASEP polynomials for $n = 3$


$$F_{(0,2,1)} =$$

$$\frac{z_2^2 z_3 + \frac{(1-t)}{(1-qt^2)}}{(1-qt^2)(1-t^2abcd)(1-q^2t^4abcd)} z_1 z_2 z_3 + \frac{1}{(1-qt^2)(1-t^2abcd)(1-q^2t^4abcd)} \left(q^3t^8a^2b^2c^2d + q^3t^8a^2b^2c^2d^2 + q^3t^7a^2b^2cd^2 + q^3t^7ab^2c^2d^2 + q^2t^7a^2b^2c^2d + q^2t^7a^2bc^2d^2 - q^3t^7a^2bcd - q^3t^7abc^2d - q^2t^6a^2b^2c^2d - q^2t^6a^2bc^2d^2 + q^3t^6a^2bcd + q^3t^6abc^2d - q^2t^6a^2bcd - q^2t^6abc^2d - q^2t^4a^2b^2cd^2 - q^2t^4ab^2c^2d^2 + q^2t^5a^2bcd - q^2t^5ab^2cd + q^2t^5abc^2d - qt^4a^2bc^2d^2 - q^3t^5abc - q^3t^5abd - q^3t^5acd - q^3t^5bcd - q^2t^4a^2bcd - qt^5a^2bcd - q^2t^4abc^2d - qt^5abc^2d + qt^4a^2bcd + qt^4abc^2d - q^2t^4abc - q^2t^4acd + q^2t^3abc + q^2t^3acd + qt^2ab^2cd + qt^2abcd^2 - qt^3abc + q^2t^2abd - qt^3acd + q^2t^2bcd + t^2a^2bcd + t^2abc^2d + q^2t^3a + q^2t^3b + q^2t^3c + 2qt^2abc + q^2t^3d + 2qt^2acd + qt^3a + qt^3c - qta - qtc - qb - qd - a - c\right) z_2 z_3 + \frac{t(tabc+tacd-a-c)}{(1-t^2abcd)} z_2^2 + \frac{(1-t)qt^2(qt^2abc+qt^2acd-a-c)}{(1-qt^2)(1-q^2t^4abcd)} z_1 z_3 - \frac{t(1-t)}{(1-qt^2)(1-t^2abcd)(1-q^2t^4abcd)} \left(q^2t^5a^2b^2c^2d + q^2t^5a^2bc^2d^2 - q^2t^5a^2bcd - q^2t^5abc^2d + q^2t^4abc + q^2t^4acd - q^2t^3abc - q^2t^3acd + qt^3abc + qt^3acd - qt^2abc - qt^2acd - qt^2a - qt^2c - tabc - tacd + qta + qtc + a + c\right) z_1 z_2 + \frac{(1-t)q}{(1-qt^2)} z_1^{-1} z_2 z_3 + \frac{1}{(1-qt^2)(1-t^2abcd)(1-qt^3abcd)(1-q^2t^4abcd)} \left(q^4t^{10}a^3b^3c^3d^3 + q^4t^{10}a^3b^3c^2d^2 + q^4t^{10}a^3b^2c^3d^2 + q^4t^{10}a^2b^3c^3d^2 + q^4t^{10}a^3b^2c^2d^3 + q^4t^{10}a^2b^2c^3d^3 + q^3t^{10}a^3b^3c^3d + q^3t^{10}a^3b^2c^3d^2 + q^3t^{10}a^3bc^3d^3 - q^3t^8a^3b^3c^3d^3 + q^3t^9a^3b^2c^3d^2 - q^4t^9a^2b^2c^2d^2 - 2q^3t^8a^3b^2c^3d^2 - q^3t^8a^3b^2c^2d - q^3t^9a^2b^2c^3d - q^3t^9a^3bc^2d^2 - q^4t^8a^2b^2c^2d^2 - q^3t^7a^3b^3c^2d^2 - q^3t^9a^2bc^3d^2 - q^3t^7a^2b^3c^3d^2 - q^3t^7a^3b^2c^2d^3 - q^3t^7a^2b^2c^3d^3 - q^3t^8a^2b^3c^2d - q^2t^7a^3b^3c^3d - q^3t^8a^3b^2c^2d - 3q^3t^8a^2b^2c^2d^2 - q^3t^8ab^2c^3d^2 - q^2t^7a^3b^2c^3d^2 - q^3t^8a^2bc^2d^3 - q^2t^7a^3bc^3d^3 - q^4t^8a^2b^2c^2d - q^4t^8ab^2c^2d - q^2t^8a^3b^2c^2d - q^2t^8a^2b^2c^3d - q^4t^8a^2bcd^2 - q^4t^8abc^2d^2 - q^2t^8a^3bc^2d^2 - q^3t^7a^2b^2c^2d^2 - q^2t^8a^2bc^3d^2 + q^3t^8a^2bc^2d + 2q^3t^6a^2b^2c^2d^2 - q^3t^7a^2b^2c^2 - 3q^3t^7a^2bc^2d + q^2t^6a^2b^2c^3d - q^3t^7a^2c^2d^2 + q^2t^6a^3bc^2d^2 - q^2t^6a^2b^2c^2d^2 + q^2t^6a^2bc^3d^2 + q^4t^7abcd + q^2t^7a^3bcd + 2q^3t^6a^2bc^2d + 2q^2t^7a^2bc^2d + q^2t^5a^2b^3c^2d + q^2t^7abc^3d + q^2t^5a^3b^2cd^2 + 4q^2t^5a^2b^2c^2d^2 + q^2t^5ab^2c^3d^2 + q^2t^5a^2bc^3d^2 - q^2t^6a^2b^2c^2 + q^3t^5a^2b^2cd + q^2t^6a^2b^2cd - 3q^2t^6a^2bc^2d + q^3t^5ab^2c^2d + q^2t^6ab^2c^2d + qt^5a^3b^2c^2d + qt^5a^2b^2c^3d + q^3t^5a^2bcd^2 + q^2t^6a^2c^2d^2 + q^3t^5abc^2d^2 + q^2t^6abc^2d^2 + qt^5a^3bc^2d^2 + qt^5a^2bc^3d^2 + q^3t^6a^2bc + q^3t^6ab^2c + q^3t^6abc^2 + 2q^2t^5a^2b^2c^2 + q^3t^6a^2bd + q^3t^6a^2cd + 3q^3t^6abcd + q^3t^6ac^2d + q^3t^6bc^2d + 3q^2t^5a^2bc^2d + qt^6a^2bc^2d + q^3t^6acd^2 + 2q^2t^5a^2c^2d^2 + qt^4a^2b^2c^2d^2 + q^3t^5abcd - qt^5a^2bc^2d + 2q^2t^5a^2bc + 2q^2t^5a^2cd - q^3t^4abcd + q^2t^5abcd - qt^4a^3bcd + 2q^2t^5ac^2d - qt^4abc^3d - 2q^2t^4a^2bc - 2q^2t^4abc^2 - 2q^2t^4a^2cd + q^2t^4abcd - qt^3a^2b^2cd - 2q^2t^4ac^2d - qt^3ab^2c^2d - qt^3a^2bcd^2 - qt^3abc^2d^2 + qt^4a^2bc - q^2t^3abc^2 + qt^4abc^2 - q^2t^3a^2bd + qt^4a^2cd - 4q^2t^3abcd - qt^4abcd + qt^4ac^2d - q^2t^3bc^2d - t^3a^2bc^2d - q^2t^3acd^2 - q^2t^4a^2 - q^2t^4ab - q^2t^4ac - q^2t^4bc - 2qt^3a^2bc - q^2t^4c^2 - 2qt^3abc^2 - q^2t^4ad - q^2t^4cd - 2qt^3a^2cd + qt^3abcd - 2qt^3ac^2d - qt^4ac - 2qt^2abcd - qt^3a^2 - qt^3ac - qt^3c^2 - q^2t^3 + 2qt^2a^2 + 2qt^2ac + 2qt^2c^2 + qtab + qtbc + qtad + qtcd + qt^2 + tac - qt + q\right) z_3 + \frac{1}{(1-qt^2)(1-t^2abcd)(1-qt^3abcd)(1-q^2t^4abcd)} \left(q^4t^{11}a^3b^3c^3d + q^4t^{11}a^3b^2c^3d^2 + q^4t^{11}a^3bc^3d^3 + q^4t^{10}a^3b^3c^2d^2 + q^4t^{10}a^3b^2c^3d^2 + q^4t^{10}a^2b^3c^3d^2 + q^4t^{10}a^3b^2c^2d^3 + q^4t^{10}a^2b^2c^3d^3 + q^3t^{10}a^3b^3c^3d + q^3t^{10}a^3b^2c^3d^2 + q^3t^{10}a^3bc^3d^3 - q^4t^{10}a^3b^3c^2d - q^4t^{10}a^3b^2c^2d^2 - q^4t^{10}a^2b^3c^2d^2 - q^4t^{10}a^3bc^2d^3 - q^4t^{10}a^2bc^3d^3 - q^3t^{10}a^3b^2c^2d - q^3t^{10}a^2b^3c^2d - q^3t^{10}a^3bc^2d^2 - q^4t^{10}a^2b^2c^2d^2 - q^4t^{10}a^2bc^3d^2 - q^4t^{10}ab^2c^3d^2 + q^3t^9a^3b^3c^2d^2 - q^3t^9a^3b^2c^3d^2 - q^3t^9a^3bc^3d^3 - q^4t^9a^2b^2c^2d^3 - q^3t^8a^3b^3c^3d^3 + q^3t^9a^3b^3c^3d - q^4t^9a^2b^3c^3d^2 - q^3t^9a^3b^2c^2d^2 - q^3t^9a^2b^3c^2d^2 - q^3t^9a^2b^2c^3d^2 - q^4t^9a^2b^2c^2d^3 + q^3t^8a^3b^2c^2d^3 + q^3t^8a^2b^3c^2d^3 + q^3t^8a^2b^2c^3d^3 + q^3t^9a^2b^2c^2d^2 + q^3t^8a^3b^2c^2d^2 + q^3t^8a^2b^3c^2d^2 + q^3t^8a^2b^2c^3d^2 + q^3t^8a^2b^2c^2d^3 - q^2t^7a^3b^3c^3d^3 + q^3t^8a^3b^3c^3d^2 + q^3t^8a^3b^2c^3d^3 + q^3t^8a^2b^3c^3d^3 + q^3t^8a^3b^3c^2d^3 + q^3t^8a^3b^2c^3d^2 + q^3t^8a^2b^3c^3d^2 + q^3t^8a^2b^2c^3d^3 - q^3t^8ab^2c^3d^3 - q^2t^7a^3b^2c^3d^3 - q^3t^8a^2bc^2d^3 - q^2t^7a^3bc^3d^3 - q^4t^8a^2b^2c^2d^2 - q^4t^8ab^2c^2d^2 - q^2t^8a^3b^2c^2d^2 - q^2t^8a^2b^2c^3d^2 - q^4t^8a^2bcd^2 - q^4t^8abc^2d^2 - q^2t^8a^3bc^2d^2 - q^3t^7a^2b^2c^2d^2 - q^2t^8a^2bc^3d^2 + q^3t^8a^2bcd + 2q^3t^6a^2b^2c^2d^2 - q^3t^7a^2b^2c^2 - 3q^3t^7a^2bc^2d + q^2t^6a^2b^2c^3d - q^3t^7a^2c^2d^2 + q^2t^6a^3bc^2d^2 - q^2t^6a^2b^2c^2d^2 + q^2t^6a^2bc^3d^2 + q^4t^7abcd + q^2t^7a^3bcd + 2q^3t^6a^2bc^2d + 2q^2t^7a^2bc^2d + q^2t^5a^2b^3c^2d + q^2t^7abc^3d + q^2t^5a^3b^2cd^2 + 4q^2t^5a^2b^2c^2d^2 + q^2t^5ab^2c^3d^2 + q^2t^5a^2bc^3d^2 - q^2t^6a^2b^2c^2 + q^3t^5a^2b^2cd + q^2t^6a^2b^2cd - 3q^2t^6a^2bc^2d + q^3t^5ab^2c^2d + q^2t^6ab^2c^2d + qt^5a^3b^2c^2d + qt^5a^2b^2c^3d + q^3t^5a^2bcd^2 + q^2t^6a^2c^2d^2 + q^3t^5abc^2d^2 + q^2t^6abc^2d^2 + qt^5a^3bc^2d^2 + qt^5a^2bc^3d^2 + q^3t^6a^2bc + q^3t^6ab^2c + q^3t^6abc^2 + 2q^2t^5a^2b^2c^2 + q^3t^6a^2bd + q^3t^6a^2cd + 3q^3t^6abcd + q^3t^6ac^2d + q^3t^6bc^2d + 3q^2t^5a^2bc^2d + qt^6a^2bc^2d + q^3t^6acd^2 + 2q^2t^5a^2c^2d^2 + qt^4a^2b^2c^2d^2 + q^3t^5abcd - qt^5a^2bc^2d + 2q^2t^5a^2bc + 2q^2t^5a^2cd - q^3t^4abcd + q^2t^5abcd - qt^4a^3bcd + 2q^2t^5ac^2d - qt^4abc^3d - 2q^2t^4a^2bc - 2q^2t^4abc^2 - 2q^2t^4a^2cd + q^2t^4abcd - qt^3a^2b^2cd - 2q^2t^4ac^2d - qt^3ab^2c^2d - qt^3a^2bcd^2 - qt^3abc^2d^2 + qt^4a^2bc - q^2t^3abc^2 + qt^4abc^2 - q^2t^3a^2bd + qt^4a^2cd - 4q^2t^3abcd - qt^4abcd + qt^4ac^2d - q^2t^3bc^2d - t^3a^2bc^2d - q^2t^3acd^2 - q^2t^4a^2 - q^2t^4ab - q^2t^4ac - q^2t^4bc - 2qt^3a^2bc - q^2t^4c^2 - 2qt^3abc^2 - q^2t^4ad - q^2t^4cd - 2qt^3a^2cd + qt^3abcd - 2qt^3ac^2d - qt^4ac - 2qt^2abcd - qt^3a^2 - qt^3ac - qt^3c^2 - q^2t^3 + 2qt^2a^2 + 2qt^2ac + 2qt^2c^2 + qtab + qtbc + qtad + qtcd + ta^2 + 2tac + tc^2 - qt + q\right) z_2 - \frac{(1-t)qt^2}{(1-qt^2)(1-t^2abcd)(1-qt^3abcd)(1-q^2t^4abcd)} \left(q^2t^7a^3b^3c^3d + q^2t^7a^3b^2c^3d^2 + q^2t^7a^3bc^3d^3 + q^2t^6a^3b^2c^3d^2 - q^2t^6a^3b^2c^2d - q^2t^6a^2b^2c^3d - q^2t^6a^3b^2c^2d^2 - qt^5a^3b^2c^2d - qt^5a^2b^2c^3d - q^2t^6a^2b^2c^2d^2 - q^2t^6a^2bc^3d^2 - qt^5a^3bc^3d^2 - qt^5a^2bc^4d^2 - q^2t^5a^2b^2c^2d^2 - q^2t^5a^2bc^3d^2 - qt^5a^2b^2c^3d^2 - qt^5a^2bc^4d^2 - q^2t^5a^2bc^2d^2 - q^2t^5a^2b^2c^2d^2 - q^2t^5a^2bc^3d^2 - q^2t^5a^2b^2c^3d^2 - q^2t^5a^2bc^2d^3 - q^2t^5a^2b^2c^2d^3 - q^2t^5a^2bc^3d^3 - q^2t^5a^2b^2c^3d^3

Branching rules for symmetric polynomials

Symmetric Schur polynomials:

$$s_\lambda(z_1, \ldots, z_n) = \sum_{\substack{\mu \subseteq \lambda \\ \lambda/\mu \text{ horizontal strip}}} z_n^{|\lambda|-|\mu|}\, s_\mu(z_1, \ldots, z_{n-1}).$$



$\lambda = (5, 2, 2)$, $\mu = (3, 2, 1)$

λ/μ is a horizontal strip: at most one box in each column.

Branching rules for symmetric polynomials

Symmetric Macdonald polynomials:

$$P_\lambda(z_1,\dots,z_n;q,t) = \sum_{\substack{\mu\subseteq\lambda \\ \lambda/\mu \text{ horizontal strip}}} \psi_{\lambda/\mu}(q,t)\, z_n^{|\lambda|-|\mu|} P_\mu(z_1,\dots,z_{n-1};q,t),$$

with branching coefficients (Macdonald '95)

$$\psi_{\lambda/\mu}(q,t) = \prod_{s\in R_{\lambda/\mu}\setminus C_{\lambda/\mu}} \frac{b_\mu(s)}{b_\lambda(s)}, \qquad b_\lambda(s) = \frac{1-q^{a_\lambda(s)}t^{l_\lambda(s)+1}}{1-q^{a_\lambda(s)+1}t^{l_\lambda(s)}},$$

where $R_{\lambda/\mu}$ (resp. $C_{\lambda/\mu}$) is the union of rows (resp. columns) meeting λ/μ.

Branching rules for symmetric polynomials

Symmetric Macdonald polynomials:

$$P_\lambda(z_1,\dots,z_n;q,t) = \sum_{\substack{\mu\subseteq\lambda \\ \lambda/\mu \text{ horizontal strip}}} \psi_{\lambda/\mu}(q,t)\, z_n^{|\lambda|-|\mu|} P_\mu(z_1,\dots,z_{n-1};q,t),$$

with branching coefficients (Macdonald '95)

$$\psi_{\lambda/\mu}(q,t) = \prod_{s\in R_{\lambda/\mu}\setminus C_{\lambda/\mu}} \frac{b_\mu(s)}{b_\lambda(s)}, \qquad b_\lambda(s) = \frac{1-q^{a_\lambda(s)}t^{l_\lambda(s)+1}}{1-q^{a_\lambda(s)+1}t^{l_\lambda(s)}},$$

where $R_{\lambda/\mu}$ (resp. $C_{\lambda/\mu}$) is the union of rows (resp. columns) meeting λ/μ.

Branching rules also exist for

- symmetric interpolation Macdonald polynomials (Okounkov '98),
- symmetric Koornwinder polynomials (van Diejen–Emsiz '15).

Branching rules for symmetric polynomials

Symmetric Macdonald polynomials:

$$P_\lambda(z_1,\dots,z_n;q,t) = \sum_{\substack{\mu\subseteq\lambda \\ \lambda/\mu \text{ horizontal strip}}} \psi_{\lambda/\mu}(q,t)\, z_n^{|\lambda|-|\mu|} P_\mu(z_1,\dots,z_{n-1};q,t),$$

with branching coefficients (Macdonald '95)

$$\psi_{\lambda/\mu}(q,t) = \prod_{s\in R_{\lambda/\mu}\setminus C_{\lambda/\mu}} \frac{b_\mu(s)}{b_\lambda(s)}, \qquad b_\lambda(s) = \frac{1-q^{a_\lambda(s)}t^{l_\lambda(s)+1}}{1-q^{a_\lambda(s)+1}t^{l_\lambda(s)}},$$

where $R_{\lambda/\mu}$ (resp. $C_{\lambda/\mu}$) is the union of rows (resp. columns) meeting λ/μ.

Branching rules also exist for

- symmetric interpolation Macdonald polynomials (Okounkov '98),
- symmetric Koornwinder polynomials (van Diejen–Emsiz '15).

These branching rules have two limitations:

- they do not have a direct analogue for nonsymmetric polynomials;
- except for the Schur case, the branching coefficients are complicated rational expressions in q and t, and the resulting formulas do not make integrality and positivity manifest.

Type A two-row recurrence

We denote by $F_\kappa^A(z_1, \ldots, z_n; q, t)$ the type A ASEP polynomials.

- They are homogeneous polynomials.
- They are indexed by compositions $\kappa \in \mathbb{N}^n$.

Type A two-row recurrence

We denote by $F_\kappa^A(z_1, \ldots, z_n; q, t)$ the type A ASEP polynomials.

- They are homogeneous polynomials.
- They are indexed by compositions $\kappa \in \mathbb{N}^n$.

$$F^A_{(0,1,0)} = z_2,$$

$$F^A_{(0,2,0)} = z_2^2 + \frac{1-t}{1-qt}\big(z_1 + qz_3\big)z_2,$$

$$F^A_{(0,2,1)} = z_2^2 z_3 + \frac{1-t}{1-qt^2}\, z_1 z_2 z_3.$$

Type A two-row recurrence

We denote by $F_\kappa^A(z_1, \ldots, z_n; q, t)$ the type A ASEP polynomials.

- They are homogeneous polynomials.
- They are indexed by compositions $\kappa \in \mathbb{N}^n$.

$$F^A_{(0,1,0)} = z_2,$$
$$F^A_{(0,2,0)} = z_2^2 + \frac{1-t}{1-qt}\big(z_1 + qz_3\big)z_2,$$
$$F^A_{(0,2,1)} = z_2^2 z_3 + \frac{1-t}{1-qt^2}\, z_1 z_2 z_3.$$

Theorem (Corteel–Mandelshtam–Williams '22)


For $\kappa \in \mathbb{N}^n$,

$$F_\kappa^A(z_1, \ldots, z_n; q, t) = \prod_{i:\, \kappa_i > 0} z_i \sum_{\substack{\mu \text{ permutation} \\ \text{of } \kappa^-}} c_\kappa^\mu(q,t)\, F_\mu^A(z_1, \ldots, z_n; q, t),$$

for some coefficients $c_\kappa^\mu(q,t)$, where κ^- is obtained from κ by decreasing each nonzero part by 1. Moreover, the coefficients $c_\kappa^\mu(q,t)$ are combinatorial.

Type A two-row recurrence


$$F_\kappa^A(z_1,\ldots,z_n;q,t) = \prod_{i:\,\kappa_i>0} z_i \sum_{\substack{\mu \text{ permutation} \\ \text{of } \kappa^-}} c_\kappa^\mu(q,t)\, F_\mu^A(z_1,\ldots,z_n;q,t)$$



$\kappa = (2,2,0,0,0,4,2,1)$

Type A two-row recurrence

$$F_\kappa^A(z_1,\ldots,z_n;q,t) = \prod_{i:\,\kappa_i>0} z_i \sum_{\substack{\mu \text{ permutation} \\ \text{of } \kappa^-}} c_\kappa^\mu(q,t)\, F_\mu^A(z_1,\ldots,z_n;q,t)$$




$\kappa = (2,2,0,0,0,4,2,1)$

$\prod_{i:\,\kappa_i>0} z_i = z_1 z_2 z_6 z_7 z_8$

Type A two-row recurrence

$$F_\kappa^A(z_1,\ldots,z_n;q,t) = \prod_{i:\,\kappa_i>0} z_i \sum_{\substack{\mu \text{ permutation} \\ \text{of } \kappa^-}} c_\kappa^\mu(q,t)\, F_\mu^A(z_1,\ldots,z_n;q,t)$$




$\kappa = (2,2,0,0,0,4,2,1)$

$\prod_{i:\,\kappa_i>0} z_i = z_1 z_2 z_6 z_7 z_8$

$\kappa^- = (1,1,0,0,0,3,1,0)$


Type A two-row recurrence

$$F_\kappa^A(z_1,\ldots,z_n;q,t) = \prod_{i:\,\kappa_i>0} z_i \sum_{\substack{\mu \text{ permutation} \\ \text{of } \kappa^-}} c_\kappa^\mu(q,t)\, F_\mu^A(z_1,\ldots,z_n;q,t)$$



$\kappa = (2,2,0,0,0,4,2,1)$


$\prod_{i:\,\kappa_i>0} z_i = z_1 z_2 z_6 z_7 z_8$



$\kappa^- = (1,1,0,0,0,3,1,0)$


permute in all possible ways

Example:




$\mu = (0,1,0,1,3,1,0,0)$

Type A two-row recurrence



A multiline queue (MLQ) with 4 rows.

Type A two-row recurrence




A multiline queue (MLQ) with 4 rows.

The weight of an MLQ is $\mathrm{wt} = \mathrm{wt}_{\mathrm{ball}}\,\mathrm{wt}_{\mathrm{pairing}}$, where:

- Ball weight: each ball in column i contributes z_i; here $z_1 z_2^2 z_3^2 z_4 z_5 z_6^2 z_7 z_8$.
- Pairing weight: a rational function in q, t, multiplicative over the layers: here $\mathrm{wt}_{\mathrm{pairing}} = \mathrm{wt}_1\,\mathrm{wt}_2\,\mathrm{wt}_3$.

Type A two-row recurrence




A multiline queue (MLQ) with 4 rows, of type $(2, 2, 0, 0, 0, 4, 2, 1)$.

The weight of an MLQ is $\mathrm{wt} = \mathrm{wt}_{\mathrm{ball}}\, \mathrm{wt}_{\mathrm{pairing}}$, where:

- Ball weight: each ball in column i contributes z_i; here $z_1 z_2^2 z_3^2 z_4 z_5 z_6^2 z_7 z_8$.
- Pairing weight: a rational function in q, t, multiplicative over the layers: here $\mathrm{wt}_{\mathrm{pairing}} = \mathrm{wt}_1\, \mathrm{wt}_2\, \mathrm{wt}_3$.

The type of an MLQ is given by the composition of its bottom row (0 for an empty position).

Type A two-row recurrence



A multiline queue (MLQ) with 4 rows, of type $(2, 2, 0, 0, 0, 4, 2, 1)$.

The weight of an MLQ is $\mathrm{wt} = \mathrm{wt}_{\mathrm{ball}}\, \mathrm{wt}_{\mathrm{pairing}}$, where:

- Ball weight: each ball in column i contributes z_i; here $z_1 z_2^2 z_3^2 z_4 z_5 z_6^2 z_7 z_8$.
- Pairing weight: a rational function in q, t, multiplicative over the layers: here $\mathrm{wt}_{\mathrm{pairing}} = \mathrm{wt}_1\, \mathrm{wt}_2\, \mathrm{wt}_3$.

The type of an MLQ is given by the composition of its bottom row (0 for an empty position).


Theorem (Corteel–Mandelshtam–Williams '22)

For $\kappa \in \mathbb{N}^n$, the type A ASEP polynomial F_κ^A is the generating function of MLQs of type κ.

Type A two-row recurrence

$$F_\kappa^A(z_1,\dots,z_n;q,t) = \prod_{i:\,\kappa_i>0} z_i \sum_{\substack{\mu \text{ permutation} \\ \text{of } \kappa^-}} c_\kappa^\mu(q,t)\, F_\mu^A(z_1,\dots,z_n;q,t)$$

Proof:




Since the pairing weight is multiplicative over layers, $c_\kappa^\mu(q,t)$ is the generating function of generalized two-line queues.


Type A two-row recurrence

$$F_\kappa^A(z_1,\dots,z_n;q,t) = \prod_{i:\,\kappa_i>0} z_i \sum_{\substack{\mu \text{ permutation} \\ \text{of } \kappa^-}} c_\kappa^\mu(q,t)\, F_\mu^A(z_1,\dots,z_n;q,t)$$

Proof:



$\equiv$




$\kappa = (2,2,0,0,0,4,2,1)$ $\qquad$ $\mu = (0,1,0,1,3,1,0,0)$

Since the pairing weight is multiplicative over layers, $c_\kappa^\mu(q,t)$ is the generating function of generalized two-line queues.

Remark: One may think of this as a *transposed* branching rule: instead of removing one column, we remove one row.

Type A two-row recurrence

Remark: A variant of the two-row recurrence technique was used to give a combinatorial formula for interpolation Macdonald polynomials (BD–Williams '25).



Two-row recurrence for Koornwinder polynomials

Theorem A (BD–Williams '26)

For $\kappa \in \mathbb{Z}^n$,


$$F_\kappa = \sum_{\substack{\nu \in \mathbb{Z}^n \\ \nu_i \in \{0, \mathrm{Sign}(\kappa_i)\}}} z^\nu \sum_{\mu \in W_0 \cdot \kappa^-} a_\kappa^\mu(\nu)\, F_\mu,$$

for some coefficients $a_\kappa^\mu(\nu) \in \mathbb{Q}(q,t,a,b,c,d)$.

- $\mathrm{Sign}(k) \in \{-1, 0, 1\}$ is the sign of $k \in \mathbb{Z}$.
- κ^- is obtained from κ by decreasing the absolute value of each nonzero part by 1.

Two-row recurrence for Koornwinder polynomials


$$F_\kappa = \sum_{\substack{\nu \in \mathbb{Z}^n \\ \nu_i \in \{0, \mathrm{Sign}(\kappa_i)\}}} z^\nu \sum_{\mu \in W_0 \cdot \kappa^-} a_\kappa^\mu(\nu)\, F_\mu$$



$\kappa = (2, -1, 0, 3, -2, 1)$

Two-row recurrence for Koornwinder polynomials

$$F_\kappa = \sum_{\substack{\nu \in \mathbb{Z}^n \\ \nu_i \in \{0, \mathrm{Sign}(\kappa_i)\}}} z^\nu \sum_{\mu \in W_0 \cdot \kappa^-} a_\kappa^\mu(\nu)\, F_\mu$$




$\kappa = (2, -1, 0, 3, -2, 1)$

for each basement cell, choose $z_i^{\mathrm{Sign}(\kappa_i)}$ or 1 according to ν

$\nu = (1, -1, 0, 0, -1, 0)$


Example:



$z^\nu = z_1 z_2^{-1} z_5^{-1}$

Two-row recurrence for Koornwinder polynomials

$$F_\kappa = \sum_{\substack{\nu \in \mathbb{Z}^n \\ \nu_i \in \{0, \mathrm{Sign}(\kappa_i)\}}} z^\nu \sum_{\mu \in W_0 \cdot \kappa^-} a_\kappa^\mu(\nu)\, F_\mu$$




$\kappa = (2, -1, 0, 3, -2, 1)$


for each basement cell, choose $z_i^{\mathrm{Sign}(\kappa_i)}$ or 1 according to ν

$\nu = (1, -1, 0, 0, -1, 0)$

Example:




$z^\nu = z_1 z_2^{-1} z_5^{-1}$



$\kappa^- = (1, 0, 0, 2, -1, 0)$

Two-row recurrence for Koornwinder polynomials

$$F_\kappa = \sum_{\substack{\nu\in\mathbb{Z}^n \\ \nu_i\in\{0,\mathrm{Sign}(\kappa_i)\}}} z^\nu \sum_{\mu\in W_0\cdot\kappa^-} a_\kappa^\mu(\nu)\, F_\mu$$




$\kappa = (2,-1,0,3,-2,1)$


for each basement cell, choose $z_i^{\mathrm{Sign}(\kappa_i)}$ or 1 according to ν

$\nu = (1,-1,0,0,-1,0)$

Example:




$z^\nu = z_1 z_2^{-1} z_5^{-1}$



$\kappa^- = (1,0,0,2,-1,0)$

permute and flip signs in all possible ways

Example:



$\mu = (0,-2,1,0,0,-1)$

Main proof idea

- Use the double affine Hecke algebra of type $(C^\vee, C)$ (Sahi '99) to define a refinement of the polynomial representation, in which we have a formal copy $\mathsf{Z}_1, \ldots, \mathsf{Z}_n$ of the variables $z_1, \ldots, z_n$.
- We construct polynomials $\widehat{E}_\kappa$ and $\widehat{F}_\kappa$ in the refined representation that satisfy

$$\widehat{F}_\kappa = \sum_\nu \mathsf{Z}^\nu \sum_\mu a_\kappa^\mu(\nu)\, F_\mu(z_1, \ldots, z_n; q, t, a, b, c, d)$$

$$\Big\downarrow \mathsf{Z}_i \mapsto z_i$$

$$F_\kappa(z_1, \ldots, z_n; q, t, a, b, c, d) = \sum_\nu z^\nu \sum_\mu a_\kappa^\mu(\nu)\, F_\mu(z_1, \ldots, z_n; q, t, a, b, c, d).$$

Two-line configurations

$$F_\kappa = \sum_{\substack{\nu \in \mathbb{Z}^n \\ \nu_i \in \{0, \mathrm{Sign}(\kappa_i)\}}} z^\nu \sum_{\mu \in W_0 \cdot \kappa^-} a_\kappa^\mu(\nu)\, F_\mu$$

Each coefficient $a_\kappa^\mu(\nu)$ is encoded by a two-line configuration $\begin{bmatrix} \sigma \\ \tau \end{bmatrix}$:


- the top row $\sigma \in \mathbb{Z}^n$ is obtained by augmenting the nonzero parts of μ;
- the bottom row $\tau \in (\mathbb{Z} \cup \overline{\mathbb{Z}})^n$, determined by κ and ν, has entries that are integers or barred integers.

Example:

$\mu = (0, -2, 1, 0, 0, -1)$

$\kappa = (2, -1, 0, 3, -2, 1)$

$\nu = (1, -1, 0, 0, -1, 0)$



$$\longrightarrow \quad a_\kappa^\mu(\nu) = \begin{bmatrix} 0 & -3 & 2 & 0 & 0 & -2 \\ 2 & -1 & 0 & \overline{3} & -2 & \overline{1} \end{bmatrix}$$

Local rules

Theorem B (BD–Williams '26)

The two-line configurations are characterized by a family of local rules.

Only the columns involved are represented; the other columns are the same on both sides. Here $m > \ell$ and $k > 0$.

In the bulk (two consecutive columns):

$$\begin{bmatrix} m & \ell \\ -k & 0 \end{bmatrix} = t \begin{bmatrix} \ell & m \\ 0 & -k \end{bmatrix} + (1-t) \begin{bmatrix} m & \ell \\ 0 & -k \end{bmatrix}.$$

On the boundary (first or last column):

$$\begin{bmatrix} 0 & \cdots \\ \overline{k} & \cdots \end{bmatrix} = -q^{k-1}ac \begin{bmatrix} 0 & \cdots \\ \overline{-k} & \cdots \end{bmatrix} - q^{k-1}(a+c) \begin{bmatrix} 0 & \cdots \\ -k & \cdots \end{bmatrix}.$$

In total, there are 73 local rules.

A cyclic algorithm

Each local rule "transports" a column to the next position, plus residual terms.

$$\begin{bmatrix} m & \ell \\ -k & 0 \end{bmatrix} = \underbrace{t \begin{bmatrix} \ell & m \\ 0 & -k \end{bmatrix}}_{\text{transported term}} + \underbrace{(1-t) \begin{bmatrix} m & \ell \\ 0 & -k \end{bmatrix}}_{\text{residual terms}},$$

$$\begin{bmatrix} 0 & \cdots \\ \overline{k} & \cdots \end{bmatrix} = \underbrace{-q^{k-1}ac \begin{bmatrix} 0 & \cdots \\ \overline{-k} & \cdots \end{bmatrix}}_{\text{transported term}} \underbrace{-q^{k-1}(a+c) \begin{bmatrix} 0 & \cdots \\ -k & \cdots \end{bmatrix}}_{\text{residual terms}}.$$

A cyclic algorithm

Each local rule "transports" a column to the next position, plus residual terms.

$$\begin{bmatrix} m & \ell \\ -k & 0 \end{bmatrix} = \underbrace{t \begin{bmatrix} \ell & m \\ 0 & -k \end{bmatrix}}_{\text{transported term}} + \underbrace{(1-t) \begin{bmatrix} m & \ell \\ 0 & -k \end{bmatrix}}_{\text{residual terms}},$$

$$\begin{bmatrix} 0 & \cdots \\ \overline{k} & \cdots \end{bmatrix} = \underbrace{-q^{k-1}ac \begin{bmatrix} 0 & \cdots \\ \overline{-k} & \cdots \end{bmatrix}}_{\text{transported term}} \underbrace{-q^{k-1}(a+c) \begin{bmatrix} 0 & \cdots \\ -k & \cdots \end{bmatrix}}_{\text{residual terms}}.$$

Fact: There exists a partial order on two-line configurations for which the residual terms are smaller.

A cyclic algorithm

Each local rule "transports" a column to the next position, plus residual terms. We solve the system by induction, moving the active column cyclically:

$$\mathcal{C} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix}$$

A cyclic algorithm

Each local rule "transports" a column to the next position, plus residual terms. We solve the system by induction, moving the active column cyclically:

$$\mathcal{C} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix}$$

$$= t \begin{bmatrix} 0 & 2 & 0 \\ 1 & 0 & -2 \end{bmatrix} + \text{smaller terms}$$

A cyclic algorithm

Each local rule "transports" a column to the next position, plus residual terms. We solve the system by induction, moving the active column cyclically:

$$\mathcal{C} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix}$$

$$= t \begin{bmatrix} 0 & 2 & 0 \\ 1 & 0 & -2 \end{bmatrix} + \text{smaller terms}$$

$$= t \begin{bmatrix} 0 & 0 & 2 \\ 1 & -2 & 0 \end{bmatrix} + \text{smaller terms}$$

A cyclic algorithm

Each local rule "transports" a column to the next position, plus residual terms. We solve the system by induction, moving the active column cyclically:

$$\mathcal{C} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix}$$

$$= t \begin{bmatrix} 0 & 2 & 0 \\ 1 & 0 & -2 \end{bmatrix} + \text{smaller terms}$$

$$= t \begin{bmatrix} 0 & 0 & 2 \\ 1 & -2 & 0 \end{bmatrix} + \text{smaller terms}$$

$$= t \begin{bmatrix} 0 & 0 & -2 \\ 1 & -2 & 0 \end{bmatrix} + \text{smaller terms}$$

A cyclic algorithm

Each local rule "transports" a column to the next position, plus residual terms. We solve the system by induction, moving the active column cyclically:

$$\mathcal{C} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix}$$

$$= t \begin{bmatrix} 0 & 2 & 0 \\ 1 & 0 & -2 \end{bmatrix} + \text{smaller terms}$$

$$= t \begin{bmatrix} 0 & 0 & 2 \\ 1 & -2 & 0 \end{bmatrix} + \text{smaller terms}$$

$$= t \begin{bmatrix} 0 & 0 & -2 \\ 1 & -2 & 0 \end{bmatrix} + \text{smaller terms}$$

$$= t^2 \begin{bmatrix} 0 & -2 & 0 \\ 1 & 0 & -2 \end{bmatrix} + \text{smaller terms}$$

A cyclic algorithm

Each local rule "transports" a column to the next position, plus residual terms. We solve the system by induction, moving the active column cyclically:

$$\mathcal{C} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix}$$

$$= t \begin{bmatrix} 0 & 2 & 0 \\ 1 & 0 & -2 \end{bmatrix} + \text{smaller terms}$$

$$= t \begin{bmatrix} 0 & 0 & 2 \\ 1 & -2 & 0 \end{bmatrix} + \text{smaller terms}$$

$$= t \begin{bmatrix} 0 & 0 & -2 \\ 1 & -2 & 0 \end{bmatrix} + \text{smaller terms}$$

$$= t^2 \begin{bmatrix} 0 & -2 & 0 \\ 1 & 0 & -2 \end{bmatrix} + \text{smaller terms}$$

$$= t^2 \begin{bmatrix} -2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix} + \text{smaller terms}$$

A cyclic algorithm

Each local rule "transports" a column to the next position, plus residual terms. We solve the system by induction, moving the active column cyclically:

$$\mathcal{C} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix}$$

$$= t \begin{bmatrix} 0 & 2 & 0 \\ 1 & 0 & -2 \end{bmatrix} + \text{smaller terms}$$

$$= t \begin{bmatrix} 0 & 0 & 2 \\ 1 & -2 & 0 \end{bmatrix} + \text{smaller terms}$$

$$= t \begin{bmatrix} 0 & 0 & -2 \\ 1 & -2 & 0 \end{bmatrix} + \text{smaller terms}$$

$$= t^2 \begin{bmatrix} 0 & -2 & 0 \\ 1 & 0 & -2 \end{bmatrix} + \text{smaller terms}$$

$$= t^2 \begin{bmatrix} -2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix} + \text{smaller terms}$$

$$= qt^2 \underbrace{\begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix}}_{=\mathcal{C}} + \text{smaller terms}$$

A cyclic algorithm

Each local rule "transports" a column to the next position, plus residual terms. We solve the system by induction, moving the active column cyclically:

$$\mathcal{C} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix}$$

$$= t \begin{bmatrix} 0 & 2 & 0 \\ 1 & 0 & -2 \end{bmatrix} + \text{smaller terms}$$

$$= t \begin{bmatrix} 0 & 0 & 2 \\ 1 & -2 & 0 \end{bmatrix} + \text{smaller terms}$$

$$= t \begin{bmatrix} 0 & 0 & -2 \\ 1 & -2 & 0 \end{bmatrix} + \text{smaller terms}$$

$$= t^2 \begin{bmatrix} 0 & -2 & 0 \\ 1 & 0 & -2 \end{bmatrix} + \text{smaller terms}$$

$$= t^2 \begin{bmatrix} -2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix} + \text{smaller terms}$$

$$= qt^2 \underbrace{\begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & -2 \end{bmatrix}}_{=\mathcal{C}} + \text{smaller terms}$$

$$(1 - qt^2)\,\mathcal{C} = \underbrace{(1-t) \begin{bmatrix} 2 & 0 & 0 \\ 1 & 0 & -2 \end{bmatrix} + t(1-t) \begin{bmatrix} 0 & 0 & -2 \\ 1 & 0 & -2 \end{bmatrix}}_{\text{smaller terms}}.$$

Application 1: nonsymmetric Askey–Wilson polynomials

Theorems A and B give explicit recurrences in the one-variable case ($n = 1$):

Theorem

For $k \geq 1$,

$$F_{(k+1)} = \big(z + u_k\big)F_{(k)} + \big(v_k z + w_k\big)F_{(-k)},$$
$$F_{(-k-1)} = \big(\widetilde{v}_k z^{-1} + \widetilde{w}_k\big)F_{(k)} + \big(z^{-1} + \widetilde{u}_k\big)F_{(-k)},$$

with $u_k = w_k\widetilde{v}_k$, $\widetilde{u}_k = v_k\widetilde{w}_k$, and

$$v_k = q^{k-1}\frac{ac + q - acq^k - abcdq^k}{1 - abcdq^{2k-1}}, \qquad w_k = q^k\frac{-(a+c) + ac(b+d)q^k}{1 - abcdq^{2k}},$$
$$\widetilde{v}_k = \frac{1 + bd - bdq^k - abcdq^{k-1}}{1 - abcdq^{2k-1}}, \qquad \widetilde{w}_k = \frac{-(b+d) + bd(a+c)q^k}{1 - abcdq^{2k}}.$$

This result refines the three-term relation for symmetric Askey–Wilson polynomials $K_{(k)} = F_{(k)} + F_{(-k)}$.

Application 1: nonsymmetric Askey–Wilson polynomials

Theorems A and B give explicit recurrences in the one-variable case ($n = 1$):

Theorem

For $k \geq 1$,

$$F_{(k+1)} = \big(z + u_k\big)F_{(k)} + \big(v_k z + w_k\big)F_{(-k)},$$
$$F_{(-k-1)} = \big(\widetilde{v}_k z^{-1} + \widetilde{w}_k\big)F_{(k)} + \big(z^{-1} + \widetilde{u}_k\big)F_{(-k)},$$

with $u_k = w_k\widetilde{v}_k$, $\widetilde{u}_k = v_k\widetilde{w}_k$, and

$$v_k = q^{k-1}\frac{ac + q - acq^k - abcdq^k}{1 - abcdq^{2k-1}}, \qquad w_k = q^k\frac{-(a+c) + ac(b+d)q^k}{1 - abcdq^{2k}},$$
$$\widetilde{v}_k = \frac{1 + bd - bdq^k - abcdq^{k-1}}{1 - abcdq^{2k-1}}, \qquad \widetilde{w}_k = \frac{-(b+d) + bd(a+c)q^k}{1 - abcdq^{2k}}.$$

This result refines the three-term relation for symmetric Askey–Wilson polynomials $K_{(k)} = F_{(k)} + F_{(-k)}$.

All these coefficients are rational functions of q^k. In particular, for $n = 1$, all two-line configurations are given by 8 rational functions of q^k. For any n, there are only finitely many rational functions to compute.

Application 2: top homogeneous parts and type A

Definition

A two-line configuration is simple if all its labels are regular (no bars), and two nonzero labels in the same column have the same sign.

Example

$$\begin{bmatrix} 2 & -2 & 0 & 0 & 3 & 0 \\ 3 & -2 & 1 & -1 & 0 & 2 \end{bmatrix}$$ is simple,

$$\begin{bmatrix} 0 & -3 & 2 & 0 & 0 & -2 \\ 2 & -1 & 0 & \overline{3} & -2 & \overline{1} \end{bmatrix}$$ is not.


Application 2: top homogeneous parts and type A

Definition

A two-line configuration is simple if all its labels are regular (no bars), and two nonzero labels in the same column have the same sign.

Proposition

A simple two-line configuration $\begin{bmatrix} \sigma \\ \tau \end{bmatrix}$ *is the generating function of reflected two-line queues with top row* σ *and bottom row* τ*.*




$$\begin{bmatrix} 3 & 0 & 3 & 0 & 0 & -2 \\ 3 & -1 & 0 & 0 & -2 & -3 \end{bmatrix}$$

Application 2: top homogeneous parts and type A

Proposition

A simple two-line configuration $\begin{bmatrix} \sigma \\ \tau \end{bmatrix}$ *is the generating function of reflected two-line queues with top row* σ *and bottom row* τ*.*



Together with the two-row recurrence (Theorem A), this gives:


Theorem C (BD–Williams '26)

For $\mu \in \mathbb{N}^n$*,*


$$\mathrm{Top}(F_\mu) = F^A_\mu, \qquad \mathrm{Top}\left(E_\mu(z_1,\dots,z_n;q,t)\right) = E^A_{\mu^{\mathrm{rev}}}(z_n,\dots,z_1;q^{-1},t^{-1}),$$

where $\mu^{\mathrm{rev}} = (\mu_n,\dots,\mu_1)$*.*


Summary


Type A Macdonald
Multiline queue formula (CMW '22)
Two-row recurrence

Summary


Type A Macdonald
Multiline queue formula (CMW '22)
Two-row recurrence
Theorem C
Top part
Koornwinder
Local rules
qKZ
Theorem B
Two-row recurrence
Theorem A
DAHA


Summary


Type A Macdonald
Multiline queue formula (CMW '22)
Two-row recurrence
??
Theorem C
Top part
Koornwinder
Local rules
qKZ
Theorem B
Two-row recurrence
Theorem A
DAHA

The affine Weyl group of type $\widetilde{C}_n$


The affine Weyl group W of type $\widetilde{C}_n$ is generated by $s_0, s_1, \ldots, s_n$ with relations

$$\begin{aligned} s_i s_j &= s_j s_i, && |i-j| \geq 2, \\ s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}, && 1 \leq i < n-1, \\ s_0 s_1 s_0 s_1 &= s_1 s_0 s_1 s_0, \\ s_{n-1} s_n s_{n-1} s_n &= s_n s_{n-1} s_n s_{n-1}, \\ s_i^2 &= 1, && 0 \leq i \leq n. \end{aligned}$$



The finite Weyl group $W_0 = \langle s_1, \ldots, s_n \rangle \simeq S_n \ltimes (\mathbb{Z}/2\mathbb{Z})^n$ is the group of signed permutations.

The affine Weyl group of type $\widetilde{C}_n$



W acts on Laurent polynomials in $z_1, \ldots, z_n$ by

$$s_i(z_1, \ldots, z_i, z_{i+1}, \ldots, z_n) = (z_1, \ldots, z_{i+1}, z_i, \ldots, z_n), \qquad 1 \leq i < n,$$
$$s_n(z_1, \ldots, z_{n-1}, z_n) = (z_1, \ldots, z_{n-1}, z_n^{-1}),$$
$$s_0(z_1, z_2, \ldots, z_n) = (qz_1^{-1}, z_2, \ldots, z_n).$$

The affine Hecke algebra $\mathcal{H}$ of type $\widetilde{C}_n$

The affine Hecke algebra $\mathcal{H}$ is generated by $T_0, T_1, \ldots, T_n$, subject to the braid relations

$$\begin{aligned} T_i T_j &= T_j T_i, && |i-j| \geq 2, \\ T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1}, && 1 \leq i < n-1, \\ T_0 T_1 T_0 T_1 &= T_1 T_0 T_1 T_0, \\ T_{n-1} T_n T_{n-1} T_n &= T_n T_{n-1} T_n T_{n-1}, \end{aligned}$$

and the quadratic relations

$$(T_i - t_i)(T_i + 1) = 0, \qquad 0 \leq i \leq n,$$

where $t_0 = -ac/q$, $t_i = t$ for $1 \leq i < n$, and $t_n = -bd$.



When $t_i = 1$ for all $0 \leq i \leq n$, we get back the affine Weyl group W, where T_i corresponds to s_i.

Polynomial representation of $\mathcal{H}$

The affine Hecke algebra $\mathcal{H}$ acts on Laurent polynomials in $z_1, \ldots, z_n$ by the Demazure–Lusztig operators

$$T_i = t + \frac{tz_i - z_{i+1}}{z_i - z_{i+1}}\,(s_i - 1), \qquad 1 \le i < n,$$

$$T_n = -bd + \frac{(1 - bz_n)(1 - dz_n)}{1 - z_n^2}\,(s_n - 1),$$

$$T_0 = -\frac{ac}{q} + \frac{(1 - a/z_1)(1 - c/z_1)}{1 - q/z_1^2}\,(s_0 - 1).$$

Theorem (Sahi '99)

This representation is faithful.

Koornwinder ASEP polynomials


Fix a partition λ. The Koornwinder ASEP polynomials $\{F_\eta\}_{\eta \in W_0 \cdot \lambda}$ are the unique family of Laurent polynomials, normalized by $[z^\eta]F_\eta = 1$, satisfying the qKZ relations (Kasatani '11)

$$
\begin{aligned}
T_0 F_{\eta_1,\eta_2,\ldots,\eta_n} &= q^{\eta_1} F_{-\eta_1,\eta_2,\ldots,\eta_n}, && \eta_1 < 0, \\
T_0 F_{\eta_1,\eta_2,\ldots,\eta_n} &= t_0\, F_{\eta_1,\eta_2,\ldots,\eta_n}, && \eta_1 = 0, \\
T_i F_{\ldots,\eta_i,\eta_{i+1},\ldots} &= t\, F_{\ldots,\eta_i,\eta_{i+1},\ldots}, && \eta_i = \eta_{i+1},\ 1 \le i < n, \\
T_i F_{\ldots,\eta_i,\eta_{i+1},\ldots} &= F_{\ldots,\eta_{i+1},\eta_i,\ldots}, && \eta_i > \eta_{i+1},\ 1 \le i < n, \\
T_n F_{\ldots,\eta_n} &= t_n\, F_{\ldots,\eta_n}, && \eta_n = 0, \\
T_n F_{\ldots,\eta_n} &= F_{\ldots,-\eta_n}, && \eta_n > 0.
\end{aligned}
$$

Recall that $t_0 = -ac/q$ and $t_n = -bd$.

The order on two-line configurations associated to $\lambda = (2, 0)$

On the six W_0-orbits associated to $\lambda = (2, 0)$, we show the poset induced by the relations $[C] > [C']$ if C' appears as a smaller term in a cycle of an element of $[C]$.



Each node represents a W_0-orbit.