

PSYCHIATRIC NURSE PRACTITIONER SERVICES

DSM-5-TR Evaluation Checklist

Monica Preder, MSN, APRN, PMHNP-BC • Board Certified Psychiatric Nurse Practitioner

Date: September 15, 2026

Practice: Therapeutic Health Services (THS) • Washington State Telehealth

|               |      |                    |                |
|---------------|------|--------------------|----------------|
| Patient Name: | DOB: | Evaluation Date:   | MRN / Chart #: |
|               |      | September 15, 2026 |                |

**Alcohol Use Disorder (AUD)**

F10.10 (Mild) / F10.20 (Moderate/Severe)

A problematic pattern of alcohol use leading to clinically significant impairment or distress, manifested by at least 2 of 11 criteria occurring within a 12-month period. Mild: 2-3 symptoms, Moderate: 4-5 symptoms, Severe: 6+ symptoms.

DSM-5 11 Criteria Checklist (Within 12-Month Period)

Mild: 2–3 • Moderate: 4–5 • Severe: 6+

- ☐ 1: Alcohol is often taken in larger amounts or over a longer period than was intended.
- ☐ 2: Persistent desire or unsuccessful efforts to cut down or control alcohol use.
- ☐ 3: Great deal of time spent in activities necessary to obtain alcohol, use alcohol, or recover from its effects.
- ☐ 4: Craving, or a strong desire or urge to use alcohol.
- ☐ 5: Recurrent alcohol use resulting in a failure to fulfill major role obligations at work, school, or home.
- ☐ 6: Continued alcohol use despite having persistent or recurrent social or interpersonal problems caused or exacerbated by alcohol.
- ☐ 7: Important social, occupational, or recreational activities given up or reduced because of alcohol use.
- ☐ 8: Recurrent alcohol use in situations in which it is physically hazardous (e.g., driving).
- ☐ 9: Alcohol use continued despite knowledge of having a persistent physical or psychological problem likely caused/exacerbated by alcohol.
- ☐ 10: Tolerance (need for markedly increased amounts to achieve intoxication, or markedly diminished effect with same amount).
- ☐ 11: Withdrawal (characteristic autonomic hyperactivity, tremor, insomnia, nausea, agitation; or alcohol/benzodiazepine taken to relieve withdrawal).

CLINICAL SPECIFIERS & SUBTYPES

Current Severity:

- ☐ Mild (2–3 symptoms: F10.10)
- ☐ Moderate (4–5 symptoms: F10.20)
- ☐ Severe (6+ symptoms: F10.20)

Remission Status:

- ☐ Active use
- ☐ In early remission (no criteria met for 3 to 12 months, except craving)
- ☐ In sustained remission (no criteria met for >= 12 months, except craving)

Diagnostic Conclusion & Medical Decision Making (MDM):

☐ Full Diagnostic Criteria Met    ☐ Sub-threshold Symptoms    ☐ Ruled Out (Alternative Differential Documented Below)

|                                                                                                                    |                                                             |
|--------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------|
| Clinician Signature:                                                                                               | Date & Clinical Time:                                       |
| Monica Preder, MSN, APRN, PMHNP-BC<br>Board Certified Psychiatric Nurse Practitioner • DEA / WA State APRN on file | Time Spent in Diagnostic Assessment & Counseling: _____ min |