

PSYCHIATRIC NURSE PRACTITIONER SERVICES

DSM-5-TR Evaluation Checklist

Monica Preder, MSN, APRN, PMHNP-BC • Board Certified Psychiatric Nurse Practitioner

Date: September 15, 2026

Practice: Therapeutic Health Services (THS) • Washington State Telehealth

Patient Name:	DOB:	Evaluation Date:	MRN / Chart #:
		September 15, 2026	

Bipolar I Disorder (Mania, Hypomania & Depression)

F31.x (Bipolar I) / F31.81 (Bipolar II)

Bipolar I requires at least one lifetime Manic Episode (lasting >= 1 week or requiring hospitalization). Hypomania requires >= 4 consecutive days without marked impairment. Bipolar depression involves a Major Depressive Episode.

Criterion A: Distinct Mood Episode

Distinct period of persistently elevated, expansive, or irritable mood + increased activity/energy

- ☐ **A_Mania:** MANIA: Lasting at least 1 week and present most of the day, nearly every day (or any duration if hospitalization is required).
- ☐ **A_Hypomania:** HYPOMANIA: Lasting at least 4 consecutive days, present most of the day, nearly every day.

Criterion B: Symptom Checklist (Requires >= 3 symptoms, or >= 4 if mood is only irritable)

Represents a noticeable change from baseline

- ☐ **B1:** Inflated self-esteem or grandiosity (grandiosity may include delusions of invincibility, superiority, special abilities, fortune, or messianic beliefs).
- ☐ **B2:** Decreased need for sleep (e.g., feels fully rested after only 3 hours of sleep).
- ☐ **B3:** More talkative than usual or pressure to keep talking (rapid, loud, interruptive speech).
- ☐ **B4:** Flight of ideas or subjective experience that thoughts are racing.
- ☐ **B5:** Distractibility (attention too easily drawn to unimportant or irrelevant external stimuli).
- ☐ **B6:** Increase in goal-directed activity (socially, at work/school, or sexually) or psychomotor agitation (purposeless, non-goal-directed activity).
- ☐ **B7:** Excessive involvement in activities that have a high potential for painful consequences (e.g., unrestrained buying sprees, sexual indiscretions, foolish business investments).

Criterion C & D: Severity, Hospitalization & Psychosis

Distinguishes Mania from Hypomania

- ☐ **Mania_Severity:** MANIA (Criterion C): Severe enough to cause marked impairment in social/occupational functioning, or necessitates hospitalization, or features psychotic symptoms (grandiose/persecutory delusions).
- ☐ **Hypomania_Severity:** HYPOMANIA (Criterion E): Not severe enough to cause marked impairment or hospitalization, and ABSOLUTELY NO psychotic features (presence of psychosis = automatic Mania).

CLINICAL SPECIFIERS & SUBTYPES

Current / Most Recent Episode:

- ☐ Manic episode (F31.1x / F31.2x)
- ☐ Hypomanic episode (F31.0)
- ☐ Depressed episode (F31.3x / F31.4x / F31.5)
- ☐ Unspecified (F31.9)

Diagnostic Features:

- ☐ With psychotic features (mood-congruent)
- ☐ With psychotic features (mood-incongruent)
- ☐ With anxious distress
- ☐ With rapid cycling (>= 4 mood episodes in 12 months)
- ☐ With mixed features

Diagnostic Conclusion & Medical Decision Making (MDM):

☐ Full Diagnostic Criteria Met ☐ Sub-threshold Symptoms ☐ Ruled Out (Alternative Differential Documented Below)

Clinician Signature:

Monica Preder, MSN, APRN, PMHNP-BC
Board Certified Psychiatric Nurse Practitioner • DEA / WA State APRN on file

Date & Clinical Time:

Time Spent in Diagnostic Assessment & Counseling: _____
min