

|               |      |                    |                |
|---------------|------|--------------------|----------------|
| Patient Name: | DOB: | Evaluation Date:   | MRN / Chart #: |
|               |      | September 15, 2026 |                |

Sedative, Hypnotic, or Anxiolytic Use Disorder

F13.10 (Mild) / F13.20 (Moderate/Severe)

Problematic pattern of benzodiazepine, sedative, or hypnotic drug use leading to clinically significant impairment or distress within a 12-month period.

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |                                        |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------|
| DSM-5 11 Criteria Checklist (Within 12-Month Period)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | Mild: 2–3 • Moderate: 4–5 • Severe: 6+ |
| <div><input type="checkbox"/> 1: Sedatives/anxiolytics often taken in larger amounts or over a longer period than intended.</div> <div><input type="checkbox"/> 2: Persistent desire or unsuccessful efforts to reduce or control sedative/anxiolytic use.</div> <div><input type="checkbox"/> 3: Great deal of time spent trying to obtain, use, or recover from sedatives/anxiolytics.</div> <div><input type="checkbox"/> 4: Craving, or strong desire to use the sedative/anxiolytic.</div> <div><input type="checkbox"/> 5: Recurrent sedative use results in failure to fulfill major obligations at work, school, or home.</div> <div><input type="checkbox"/> 6: Continued use despite persistent social or interpersonal problems.</div> <div><input type="checkbox"/> 7: Important activities given up or reduced because of sedative/anxiolytic use.</div> <div><input type="checkbox"/> 8: Recurrent sedative use in situations that are physically hazardous.</div> <div><input type="checkbox"/> 9: Continued use despite knowledge of physical/psychological problem likely caused by drug.</div> <div><input type="checkbox"/> 10: Tolerance developed (not counted if taken solely under appropriate medical supervision).</div> <div><input type="checkbox"/> 11: Withdrawal syndrome (autonomic hyperactivity, hand tremor, insomnia, nausea/vomiting, transient hallucinations, psychomotor agitation, anxiety, grand mal seizures; or use to avoid symptoms).</div> |                                        |

CLINICAL SPECIFIERS & SUBTYPES

Current Severity:

☐ Mild (2–3 symptoms: F13.10)

☐ Moderate (4–5 symptoms: F13.20)

☐ Severe (6+ symptoms: F13.20)

Remission Status:

☐ Active use

☐ In early remission (3–12 months)

☐ In sustained remission (> 12 months)

Diagnostic Conclusion & Medical Decision Making (MDM):

☐ Full Diagnostic Criteria Met

☐ Sub-threshold Symptoms

☐ Ruled Out (Alternative Differential Documented Below)

|                                                                                                                                       |                                                                                       |
|---------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| Clinician Signature:                                                                                                                  | Date & Clinical Time:                                                                 |
| <div>Monica Preder, MSN, APRN, PMHNP-BC</div> <div>Board Certified Psychiatric Nurse Practitioner • DEA / WA State APRN on file</div> | <div>Time Spent in Diagnostic Assessment &amp; Counseling: _____</div> <div>min</div> |