

This Mathematica notebook contains the computations for the paper “Two connected populations with Allee effect” where the sequence of nonnegative steady state of the system (1) of the paper for fixed values of the Allee threshold parameter, but varying spatial dispersal rate is computed in the one way migration case.

Part 2: Right side of Figure 4.

This section contains the functions that are used in the next section.

Consider the parametric system of polynomial equations given at equation (2) at the paper with the inequality constraints on the variables and the parameters, and $\delta_{1,2}=1$, $\delta_{2,1}=0$.

We define systemEquations to be the list of the two polynomial equations resulted by letting $\dot{N}_1 = \dot{N}_2 = 0$.

```
In[1]:= systemEquations = {N1 * (1 - N1) * (N1 - b) - a * N1, N2 * (1 - N2) * (N2 - b) + a * N1}
Out[1]= {-a N1 + (1 - N1) N1 (-b + N1), a N1 + (1 - N2) N2 (-b + N2)}
```

The following function in addition to what solvor function in part 1 does, substitute each solution at the Jacobian matrix of the system and computed the eigenvalues to decide stability of these points as steady states.

```

In[2]:= solvorStability[aa_, bb_] := (
  equations = ReplaceAll[{a → aa, b → bb}][systemEquations];
  solutions =
    NSolve[equations[[1]] == 0 && equations[[2]] == 0 && N1 ≥ 0 && N2 ≥ 0, {N1, N2}, Reals];
  pointsWithStability = ConstantArray[{0, 0}, 1], Length[solutions]];
  For[i = 1, i ≤ Length[solutions], i++,
    pointsWithStability[[i]][[1]] = ReplaceAll[solutions[[i]]][{N1, N2}];
  ];
  JacobM = ConstantArray[0, {2, 2}];
  For[i = 1, i ≤ 2, i++,
    For[j = 1, j ≤ 2, j++,
      JacobM[[i, j]] = D[equations[[i]], Nj];
    ];
  ];
  For[i = 1, i ≤ Length[pointsWithStability], i++,
    JacobMSubsed = ReplaceAll[{N1 → pointsWithStability[[i]][[1]][[1]],
      N2 → pointsWithStability[[i]][[1]][[2]]}][JacobM];
    EigenvaluesJacobMSubsed = Eigenvalues[JacobMSubsed];
    For[j = 1, j ≤ Length[EigenvaluesJacobMSubsed], j++,
      If[Re[EigenvaluesJacobMSubsed[[j]]] ≥ 0, {pointsWithStability[[i]][[2]] = 0;
        Break}];
    ];
  ];
  Return[pointsWithStability];
)

```

The following function finds the critical values of a for a fixed given value of b . This is done by solving $f(a,b)=0$ after substituting the given choice of b . Where f is any of the two polynomials defining the boundary between the open regions in Figure 3 of the paper. They are accessibly by asking the RootFinding[Parametric] package of Maple to show the algebraic description of the cells.

```

In[3]:= criticalValuesA[bb_] := (
  equation1 = ReplaceAll[{b -> bb}] [
    16 * a^3 * b^6 + 16 * a^2 * b^7 - 4 * a * b^8 + 108 * a^4 * b^4 + 60 * a^3 * b^5 - 75 * a^2 * b^6 +
    10 * a * b^7 + b^8 - 54 * a^4 * b^3 - 12 * a^3 * b^4 + 42 * a^2 * b^5 + 4 * a * b^6 - 4 * b^7 +
    729 * a^6 + 1458 * a^5 * b + 405 * a^4 * b^2 - 328 * a^3 * b^3 + 50 * a^2 * b^4 - 20 * a * b^5 +
    6 * b^6 - 54 * a^4 * b - 12 * a^3 * b^2 + 42 * a^2 * b^3 + 4 * a * b^4 - 4 * b^5 + 108 * a^4 +
    60 * a^3 * b - 75 * a^2 * b^2 + 10 * a * b^3 + b^4 + 16 * a^3 + 16 * a^2 * b - 4 * a * b^2];
  equation2 = ReplaceAll[{b -> bb}] [4 * a - b^2 + 2 * b - 1];
  solutions1 = Solve[equation1 == 0 && a ≥ 0, {a}, Reals];
  points1 = ConstantArray[{0, 0}, Length[solutions1]];
  For[i = 1, i ≤ Length[solutions1], i++,
    points1[[i]] = ReplaceAll[solutions1[[i]]][a];
  ];
  solutions2 = Solve[equation2 == 0 && a ≥ 0, {a}, Reals];
  points2 = ConstantArray[{0, 0}, Length[solutions2]];
  For[i = 1, i ≤ Length[solutions2], i++,
    points2[[i]] = ReplaceAll[solutions2[[i]]][a];
  ];
  criticalValues = Union[points1, points2];
  Return[Sort[criticalValues, Less]];
)

```

The following function receives a choice of b and the list of values as critical values of a and a list specifying how many sample points between two critical values must be considered and then returns a list of solutions for the given value of b and different values of a .

```

In[4]:= solvorSequencePoints[bb_, alist_, NNL_] := (
  criticalN = Length[alist] - 1;
  pointsL = {};
  (* a=alist[[1]] *)
  AppendTo[pointsL, solvorStability[alist[[1]], bb]];
  (* alist[[i]] until alist[[i+1]] *)
  For[l1 = 1, l1 ≤ criticalN, l1++,
    NN = NNL[[l1]];
    For[l2 = 1, l2 ≤ NN, l2++,
      aa = ((NN - l2) * alist[[l1]] + l2 * alist[[l1 + 1]]) / NN;
      AppendTo[pointsL, solvorStability[aa, bb]];
    ];
  ];
  Return[pointsL];
)

```

The following function is a sub-function for the next function.

```

In[5]:= plotSubTask[pointos_] := (
  inputo1 = {};
  inputo2 = {};
  For[l1 = 1, l1 ≤ Length[pointos], l1++,
    For[l2 = 1, l2 ≤ Length[pointos[[l1]]], l2++,
      AppendTo[inputo1, {pointos[[l1]][[l2]][[1]]}];
      colorstability = If[pointos[[l1]][[l2]][[2]] == 1,
        {255/255, 36/255, 0/255}, {251/255, 206/255, 177/255}];
      AppendTo[inputo2, {Directive[RGBColor[colorstability[[1]],
        colorstability[[2]], colorstability[[3]]], PointSize[0.02]]}];
    ];
  ];
  Return[{inputo1, inputo2}]
)

```

The following function plots the steady states for a fixed value of b , but varying values of a and color them with respect their stability as in the right side of Figure 4 of the paper. Red means stable and pink means unstable.

```

In[6]:= plotTask[bb_, alist_, NNL_] := (
  SSlist = solvorSequencePoints[bb, alist, NNL];
  subTaskOutput = plotSubTask[SSlist];
  Return[ListPlot[subTaskOutput[[1]], PlotStyle → subTaskOutput[[2]], AxesLabel →
    {Style["N1", FontSize → 16, Bold, Black], Style["N2", FontSize → 16, Bold, Black]}]]
)

```

This section contains the plots used in Figure 4 of the paper.

Sequences for $0 < b < 0.4961661$.

$b = 0.1$

```

In[7]:= N[criticalValuesA[1/10]]

```

```


Out[7]= {0.00238297, 0.0194548, 0.2025}

```

```

In[8]:= plotTask[1/10,
  {0, 0.002382970637023988, 0.019454811603742996, 0.2025, 0.21}, {5, 5, 100, 1}]

```




$b = 0.3$

In[]:= **N[criticalValuesA[3/10]]**

Out[]:= {0.01986, 0.0504942, 0.1225}

In[]:= **plotTask[3/10,**
 $\{0, 0.01986001392249426^{\sim}, 0.050494173879918135^{\sim}, 0.1225^{\sim}, 0.13\}, \{20, 20, 90, 1\}$ **]**




Sequence for $b=0.4961661$. (Critical Value)

In[]:= **N[criticalValuesA[4961661 * 10⁻⁷]]**

Out[]:= {0.057545, 0.0634621, 0.0634621}

```
In[ ]:= plotTask[4961661 * 10^(-7),
  {0, 0.057544964217530525`, 0.06346214969730232`, 0.064}, {50, 20, 1}]
```




Sequence for $0.4961661 < b < 1/2$.

$b = 0.948$

```
In[ ]:= N[criticalValuesA[498 / 1000]]
```

```
Out[ ]:= {0.0586017, 0.0628843, 0.063001}
```

```
In[ ]:= plotTask[498 / 1000,
  {0, 0.05860173612241946`, 0.06288434240012102`, 0.063001`, 0.064}, {50, 10, 3, 1}]
```




Sequence for $b=1/2$. (Critical value)

```
In[ ]:= N[criticalValuesA[1 / 2]]
```

```
Out[ ]:= {0.0610042, 0.0625}
```

```
In[ ]:= plotTask[1/2, {0, 0.0610042339640731`, 0.0625`, 0.065}, {40, 10, 1}]
```




Sequences for $b > 1/2$.

$b = 0.6$

```
In[ ]:= N[criticalValuesA[6/10]]
```

```
Out[ ]:= {0.04}
```

```
In[ ]:= plotTask[6/10, {0, 0.04`, 0.05}, {80, 1}]
```




$b = 0.9$

```
In[ ]:= N[criticalValuesA[9/10]]
```

```
Out[ ]:= {0.0025}
```

```
In[ ]:= plotTask[9/10, {0, 0.0025`, 0.003}, {20, 1}]
```



End of the file.

Source paper: Gergely Rost and AmirHosein Sadeghimanesh, Two connected populations with Allee effect, 2021.