

# Linear Algebra Review

# Why do we need Linear Algebra?

- We will associate coordinates to
  - 3D points in the scene
  - 2D points in the CCD array
  - 2D points in the image
- Coordinates will be used to
  - Perform geometrical transformations
  - Associate 3D with 2D points
- Images are matrices of numbers
  - We will find properties of these numbers

# Matrices

$$A_{n \times m} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1m} \\ a_{21} & a_{22} & \cdots & a_{2m} \\ a_{31} & a_{32} & \cdots & a_{3m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nm} \end{bmatrix}$$

Sum:

$$C_{n \times m} = A_{n \times m} + B_{n \times m}$$

$$c_{ij} = a_{ij} + b_{ij}$$


A and B must have the same dimensions

Example:

$$\begin{bmatrix} 2 & 5 \\ 3 & 1 \end{bmatrix} + \begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix} = \begin{bmatrix} 8 & 7 \\ 4 & 6 \end{bmatrix}$$

# Matrices

Product:

$$C_{n \times p} = A_{n \times m} B_{m \times p}$$


A and B must have compatible dimensions

$$c_{ij} = \sum_{k=1}^m a_{ik} b_{kj}$$

$$A_{n \times n} B_{n \times n} \neq B_{n \times n} A_{n \times n}$$

Examples:

$$\begin{bmatrix} 2 & 5 \\ 3 & 1 \end{bmatrix} \cdot \begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix} = \begin{bmatrix} 17 & 29 \\ 19 & 11 \end{bmatrix}$$

$$\begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix} \cdot \begin{bmatrix} 2 & 5 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} 18 & 32 \\ 17 & 10 \end{bmatrix}$$

# Matrices

Transpose:

$$C_{m \times n} = A^T_{n \times m}$$

$$c_{ij} = a_{ji}$$

$$(A + B)^T = A^T + B^T$$

$$(AB)^T = B^T A^T$$

If  $A^T = A$   $A$  is symmetric

Examples:

$$\begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix}^T = \begin{bmatrix} 6 & 1 \\ 2 & 5 \end{bmatrix} \quad \begin{bmatrix} 6 & 2 \\ 1 & 5 \\ 3 & 8 \end{bmatrix}^T = \begin{bmatrix} 6 & 1 & 3 \\ 2 & 5 & 8 \end{bmatrix}$$

# Matrices

Determinant:      A must be square

$$\det \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = a_{11}a_{22} - a_{21}a_{12}$$

$$\det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

Example:  $\det \begin{bmatrix} 2 & 5 \\ 3 & 1 \end{bmatrix} = 2 - 15 = -13$

# Matrices

Inverse:

A must be square

$$A_{n \times n} A_{n \times n}^{-1} = A_{n \times n}^{-1} A_{n \times n} = I$$


$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}^{-1} = \frac{1}{a_{11}a_{22} - a_{21}a_{12}} \begin{bmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{bmatrix}$$

Example:  $\begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix}^{-1} = \frac{1}{28} \begin{bmatrix} 5 & -2 \\ -1 & 6 \end{bmatrix}$

$$\begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix}^{-1} \cdot \begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix} = \frac{1}{28} \begin{bmatrix} 5 & -2 \\ -1 & 6 \end{bmatrix} \cdot \begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix} = \frac{1}{28} \begin{bmatrix} 28 & 0 \\ 0 & 28 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

# 2D Vector

$$\mathbf{v} = (x_1, x_2)$$



Magnitude:  $\|\mathbf{v}\| = \sqrt{x_1^2 + x_2^2}$

If  $\|\mathbf{v}\| = 1$ ,  $\mathbf{v}$  Is a UNIT vector


$$\frac{\mathbf{v}}{\|\mathbf{v}\|} = \left( \frac{x_1}{\|\mathbf{v}\|}, \frac{x_2}{\|\mathbf{v}\|} \right) \text{ Is a unit vector}$$

Orientation:  $\theta = \tan^{-1} \left( \frac{x_2}{x_1} \right)$




# Vector Addition

$$\mathbf{v} + \mathbf{w} = (x_1, x_2) + (y_1, y_2) = (x_1 + y_1, x_2 + y_2)$$




# Vector Subtraction

$$\mathbf{v} - \mathbf{w} = (x_1, x_2) - (y_1, y_2) = (x_1 - y_1, x_2 - y_2)$$




# Scalar Product

$$a\mathbf{v} = a(x_1, x_2) = (ax_1, ax_2)$$



# Inner (dot) Product




$$v \cdot w = (x_1, x_2) \cdot (y_1, y_2) = x_1 y_1 + x_2 \cdot y_2$$

The inner product is a **SCALAR!**

$$v \cdot w = (x_1, x_2) \cdot (y_1, y_2) = \|v\| \cdot \|w\| \cos \alpha$$

$$v \cdot w = 0 \Leftrightarrow v \perp w$$

# Orthonormal Basis



$$\mathbf{i} = (1,0) \quad \|\mathbf{i}\| = 1$$

$$\mathbf{j} = (0,1) \quad \|\mathbf{j}\| = 1$$

$$\mathbf{i} \cdot \mathbf{j} = 0$$


$$\mathbf{v} = (x_1, x_2)$$

$$\mathbf{v} = x_1 \mathbf{i} + x_2 \mathbf{j}$$

$$\mathbf{v} \cdot \mathbf{i} = (x_1 \mathbf{i} + x_2 \mathbf{j}) \cdot \mathbf{i} = x_1 \cdot 1 + x_2 \cdot 0 = x_1$$

$$\mathbf{v} \cdot \mathbf{j} = (x_1 \mathbf{i} + x_2 \mathbf{j}) \cdot \mathbf{j} = x_1 \cdot 0 + x_2 \cdot 1 = x_2$$

# Vector (cross) Product



$$u = v \times w$$

The cross product is a **VECTOR!**

Magnitude:  $\|u\| = \|v \times w\| = \|v\| \|w\| \sin \alpha$

Orientation:

$$u \perp v \Rightarrow u \cdot v = (v \times w) \cdot v = 0$$
$$u \perp w \Rightarrow u \cdot w = (v \times w) \cdot w = 0$$

# Vector Product Computation


$$\mathbf{k} = (0, 0, 1) \quad \|\mathbf{k}\| = 1$$

$$\mathbf{j} = (0, 1, 0) \quad \|\mathbf{j}\| = 1 \quad \mathbf{i} \cdot \mathbf{j} = 0, \mathbf{i} \cdot \mathbf{k} = 0, \mathbf{j} \cdot \mathbf{k} = 0$$

$$\mathbf{i} = (1, 0, 0) \quad \|\mathbf{i}\| = 1$$

$$\mathbf{u} = \mathbf{v} \times \mathbf{w} = (x_1, x_2, x_3) \times (y_1, y_2, y_3)$$

$$\mathbf{u} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{vmatrix}$$




$$= (x_2 y_3 - x_3 y_2) \mathbf{i} + (x_3 y_1 - x_1 y_3) \mathbf{j} + (x_1 y_2 - x_2 y_1) \mathbf{k}$$


# 2D Geometrical Transformations



# 2D Translation



# 2D Translation Equation




$$\mathbf{P} = (x, y)$$

$$\mathbf{t} = (t_x, t_y)$$

$$\mathbf{P}' = (x + t_x, y + t_y) = \mathbf{P} + \mathbf{t}$$

# 2D Translation using Matrices



$$\mathbf{P} = (x, y)$$

$$\mathbf{t} = (t_x, t_y)$$

$$\mathbf{P}' \rightarrow \begin{bmatrix} x + t_x \\ y + t_y \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

The diagram illustrates the matrix representation of 2D translation. The translation vector  $\mathbf{t} = (t_x, t_y)$  is used to translate a point  $\mathbf{P} = (x, y)$  to a new point  $\mathbf{P}' = (x + t_x, y + t_y)$ . The transformation is represented by a 2x3 matrix multiplied by a 3x1 column vector. The matrix is  $\begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \end{bmatrix}$ , where the translation components  $t_x$  and  $t_y$  are highlighted with a pink box and a pink arrow. The column vector is  $\begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$ , where the original coordinates  $x$  and  $y$  are boxed, and the bottom element 1 is circled in pink with a pink arrow pointing to it. The result of the multiplication is the translated point  $\mathbf{P}'$ .

# Homogeneous Coordinates

- Multiply the coordinates by a non-zero scalar and add an extra coordinate equal to that scalar. For example,

$$(x, y) \rightarrow (x \cdot z, y \cdot z, z) \quad z \neq 0$$

$$(x, y, z) \rightarrow (x \cdot w, y \cdot w, z \cdot w, w) \quad w \neq 0$$

- **NOTE:** If the scalar is 1, there is no need for the multiplication!


# Back to Cartesian Coordinates:

- Divide by the last coordinate and eliminate it. For example,

$$(x, y, z) \quad z \neq 0 \rightarrow (x/z, y/z)$$

$$(x, y, z, w) \quad w \neq 0 \rightarrow (x/w, y/w, z/w)$$

# 2D Translation using Homogeneous Coordinates




$$\mathbf{P} = (x, y) \rightarrow (x, y, 1)$$

$$\mathbf{t} = (t_x, t_y) \rightarrow (t_x, t_y, 1)$$


$$\mathbf{P}' \rightarrow \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix} = \underbrace{\begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}}_{\mathbf{T}} \cdot \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$\mathbf{P}' = \mathbf{T} \cdot \mathbf{P}$$

# Scaling



# Scaling Equation



$$\mathbf{P} = (x, y) \rightarrow (x, y, 1)$$


$$\mathbf{P}' = (s_x x, s_y y) \rightarrow (s_x x, s_y y, 1)$$

$$\mathbf{P}' \rightarrow \begin{bmatrix} s_x x \\ s_y y \\ 1 \end{bmatrix} = \underbrace{\begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix}}_{\mathbf{S}} \cdot \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$\mathbf{P}' = \mathbf{S} \cdot \mathbf{P}$$



# Scaling & Translating



$$P'' = T.P' = T.(S.P) = (T.S).P$$

# Scaling & Translating

$$P'' = T.P' = T.(S.P) = (T.S).P$$


$$\begin{aligned} \mathbf{P}'' = \mathbf{T} \cdot \mathbf{S} \cdot \mathbf{P} &= \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \\ &= \begin{bmatrix} s_x & 0 & t_x \\ 0 & s_y & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x x + t_x \\ s_y y + t_y \\ 1 \end{bmatrix} \end{aligned}$$

# Translating & Scaling $\neq$ Scaling & Translating

$$P'' = S \cdot P' = S \cdot (T \cdot P) = (S \cdot T) \cdot P$$


$$\begin{aligned} \mathbf{P}'' = \mathbf{S} \cdot \mathbf{T} \cdot \mathbf{P} &= \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \\ &= \begin{bmatrix} s_x & 0 & s_x t_x \\ 0 & s_y & s_y t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x x + s_x t_x \\ s_y y + s_y t_y \\ 1 \end{bmatrix} \end{aligned}$$

# Rotation



# Rotation Equations

Counter-clockwise rotation by an angle  $\theta$



$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$\mathbf{P}' = \mathbf{R} \cdot \mathbf{P}$$

# Degrees of Freedom

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

R is 2x2  4 elements

BUT! There is only 1 degree of freedom:  $\theta$

The 4 elements must satisfy the following constraints:

$$\mathbf{R} \cdot \mathbf{R}^T = \mathbf{R}^T \cdot \mathbf{R} = \mathbf{I}$$

$$\det(\mathbf{R}) = 1$$

# Scaling, Translating & Rotating



Order matters!

$$P' = S.P$$

$$P'' = T.P' = (T.S).P$$


$$P''' = R.P'' = R.(T.S).P = (R.T.S).P$$



$$R.T.S \neq R.S.T \neq T.S.R \dots$$

# 3D Rotation of Points

Rotation around the coordinate axes, **counter-clockwise**:



$$R_x(\alpha) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \alpha & -\sin \alpha \\ 0 & \sin \alpha & \cos \alpha \end{bmatrix}$$

$$R_y(\beta) = \begin{bmatrix} \cos \beta & 0 & \sin \beta \\ 0 & 1 & 0 \\ -\sin \beta & 0 & \cos \beta \end{bmatrix}$$

$$R_z(\gamma) = \begin{bmatrix} \cos \gamma & -\sin \gamma & 0 \\ \sin \gamma & \cos \gamma & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



# 3D Rotation (axis & angle)

$$\mathbf{n} = [n_1 \quad n_2 \quad n_3]^T, \text{ angle } \theta$$

$$\mathbf{R} = \mathbf{I} \cos \theta + \mathbf{I}(1 - \cos \theta) \begin{bmatrix} n_1^2 & n_1 n_2 & n_1 n_3 \\ n_1 n_2 & n_2^2 & n_2 n_3 \\ n_1 n_3 & n_2 n_3 & n_3^2 \end{bmatrix} + \sin \theta \begin{bmatrix} 0 & -n_3 & n_2 \\ n_3 & 0 & -n_1 \\ -n_2 & n_1 & 0 \end{bmatrix}$$

# 3D Translation of Points

Translate by a vector  $\mathbf{t}=(t_x,t_y,t_z)^T$ :

$$T = \begin{bmatrix} 1 & 0 & 0 & t_x \\ 0 & 1 & 0 & t_y \\ 0 & 0 & 1 & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

