

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/305733074>

# Data and Doing: Using Wound Size to Evaluate Wound Care in Venous Ulcers

**Article** in *Advances in Skin & Wound Care* · August 2016

DOI: 10.1097/01.ASW.0000483250.24007.23

---

CITATIONS

6

---

READS

178

2 authors, including:



Gregory Bohn

38 PUBLICATIONS 478 CITATIONS

SEE PROFILE

## Data and Doing: Using Wound Size to Evaluate Wound Care in Venous Ulcers

Barbara Bates-Jensen, PhD, RN, FAAN; and Gregory A. Bohn, MD, FACS

### ABSTRACT

Measuring and monitoring wound progress by size are an important management tool. Wound progress and size can help assess the effectiveness of therapy and predict healing, while preparing for application of advanced wound products and treatments. The authors outline methods of tracking wound size and predictors of healing in venous leg ulcers.

**KEYWORDS:** wound measurement, predicting wound healing, advanced wound therapy

ADV SKIN WOUND CARE 2016;29:347–8.

The use of wound size as an essential component of wound assessment is included in all clinical practice guidelines for wounds.<sup>1–6</sup> Measuring wound size is an established standard practice in wound care clinics and other healthcare settings. Although it seems intuitive that assessing for changes in wound size is an essential component of evaluating wound treatment, clearly the wound has to get smaller to heal; interpreting the rate of and trend in wound size change is often overlooked as a predictor for wound healing. Furthermore, research studies on wound size and other wound and patient characteristics on initial wound presentation can be valuable to keep in mind when assessing a patient's prognostic chance to heal.<sup>7–9</sup>

Understanding whether a particular wound be difficult to heal is useful in clinical practice and allows the clinician to prepare for failure of the wound to progress.<sup>10</sup> Needed changes in treatment can be anticipated, thus preventing delay in initiation of advanced wound therapies that may favorably impact ability to heal. The timely readjustment of treatment may improve time to wound closure with resultant lower probability of complications.<sup>11</sup>

Although the demonstrated failure of wound treatment to improve the wound is inherent in certain reimbursement requirements, further delay in the implementation of advanced wound therapies could be avoided by preparing for the likely event that the wound, based on initial evaluation, is likely to fail to progress. Therefore, early wound screening based on

changes and trends in wound size as a predictor of wound healing likelihood is best practice for wound clinicians.

### Wound Measurement Methods

Multiple methods are available to measure wounds to determine their surface area. Simple linear measurement of length and width multiplied to estimate surface area can be obtained using standard reference points (eg, head to toe and side to side, clock face 12:00 to 6:00 and 9:00 to 3:00; or longest aspect by perpendicular widest aspect), acetate film tracing with area calculation, or laser scanner technology embedded in cameras or devices, which have all been used and described.<sup>10</sup>

The accuracy of various methods differs; however, routinely using a consistent reproducible wound measurement method that can be applied by varied caregivers with each wound assessment allows for comparison of weekly change to track healing progress.<sup>10,12</sup> Linear measurements of length and width to calculate surface area have been correlated with healing and have been commonly used to determine wound size and surface area<sup>9,13</sup>; however, newer technology is offering advanced measurement options. Use of laser technology embedded in a camera-type device has simplified obtaining and improved the accuracy of wound size measures. Deciding which method to use to obtain wound size is often a matter of clinician preference.

### Using Data to Predict Healing in Venous Ulcers

Treating patients with venous ulcers (VUs) can be somewhat predictable, based on known criteria both at initial presentation and during reevaluation during care. Venous leg ulcers comprise approximately 30% to 40% of chronic wounds and can be some of the most challenging to heal. The goal in treating VUs is to achieve healing in 12 weeks. If a wound fails to heal by the 12th week of treatment, the wound is considered to fall outside the expected healing parameters.<sup>2,4,7,8</sup>

There are 2 methods of using data and research evidence to predict which VUs are more likely to heal. The first method involves evaluating characteristics during the initial assessment.

Barbara Bates-Jensen, PhD, RN, FAAN, is Professor, Nursing and Medicine, University of California–Los Angeles School of Nursing and David Geffen School of Medicine; a Fellow in the American Academy of Nursing; and President of the Wound Reach Foundation. Gregory Bohn, MD, FACS, is a Surgeon and Medical Director, West Shore Medical Center for Wound Care and Hyperbaric Medicine, Manistee, Michigan; President of the American Board of Wound Healing; a Fellow of the American College of Hyperbaric Medicine and of the American Professional Wound Care Association; and Vice President of the Wound Reach Foundation. Dr Bates-Jensen has disclosed that she is a member of the Cardinal Health Advisory Board; her institution has received grants from the National Institutes of Health and the US Department of Veterans Affairs; she is recipient of the UCLA SEM Scanner patent; royalties from her book authorship are paid directly to the Wound Care Foundation; and she was reimbursed by Bruin Biometrics for travel expenses to attend the European Wound Management Association meeting to present a lecture (unpaid). Dr. Bohn has disclosed that he is a volunteer board member of the Wound Reach Foundation and the Association for the Advancement of Wound Care (AAWC) (current President); he is compensated by the AAWC for travel expenses related to association work; and he has previously been reimbursed for providing educational services to Wound Care Advantage. "Quality Measures" is a review column published in partnership with the Wound Reach Foundation, a 501(c)(3) nonprofit charity organization. Submitted January 20, 2016; accepted March 24, 2016.

- wound duration 6 months or greater = 1 point
- wound size 5 cm<sup>2</sup> or greater = 1 point

If the total score equals zero, the positive predictive value is approximately 93% that the VU will heal with compression. If the total score is 1 or 2, it is unlikely that compression alone will result in ulcer healing, so these VUs should be treated with advanced therapies.<sup>7,9,14,15</sup> A second approach includes evaluation of the patient's ankle-brachial index. Patients presenting with wounds less than 10 cm<sup>2</sup>, wounds of less than 12 months' duration, and with an ankle-brachial index of 0.8 or greater indicating probable absence of peripheral arterial disease have an 80% chance of healing with standard therapy within 24 weeks.<sup>7,14</sup> Clinicians must understand that when negative risk factors are present, the expectation for success may be much less, even at 24 weeks of care. Identifying this poorer potential to heal at the outset of care may help clinicians prepare to intervene quickly and attempt to change the course of care when benchmarks of healing are not present initially.

When a VU has not met goals of healing, it is recommended to reevaluate the treatment plan and possibly change therapy. In addition, Medicare, Medicare intermediaries, and commercial insurers use these data assessments as criteria to warrant application of advanced modalities.<sup>17</sup> Given the lack of response to standard therapy, advanced wound therapy can be

## Summary

The data on wound size, duration, and presence of peripheral arterial disease on initial presentation, along with monitoring for surface area reductions, are not new. However, use of these data in clinical practice remains uneven. Incorporating evidence-based therapy to treat patients with VUs would help to ensure that patients receive the best treatment available from the initial assessment visit. Having the ability to assess progress as it relates to treatment gives the clinician a means by which to evaluate the effectiveness of that treatment and how the patient responds to it. If the patient is not responding to treatment, the clinician has the option to change or modify the treatment regimen to effect improvement and healing. In wound care, change in wound surface area is best used to follow wound progress. ●

## References

1. Lazarus GS, Cooper DM, Knighton DR, et al. Definitions and guidelines for the assessment of wounds and evaluation of healing. *Arch Dermatol* 1994;130:489-93.
2. Robson MC, Cooper DM, Gould LJ, et al. Guidelines for the treatment of venous ulcers. *Wound Rep Reg* 2006;12:649-62.
3. Whitney J, Phillips L, Aslam R, et al. Guidelines for the treatment of pressure ulcers. *Wound Rep Reg* 2006;14:663-79.
4. Glowiczki P, Comerota AJ, Dalsing MC, et al. The care of patients with varicose veins and associated chronic venous diseases: Clinical practice guidelines of the Society for Vascular Surgery and the American Venous Forum journal of vascular surgery. *J Vasc Surg* 2011;53(5 Suppl):2S-48S.
5. Association for the Advancement of Wound Care Venous Ulcer Guideline. Malvern, Pennsylvania: Association for the Advancement of Wound Care. December 2010. <http://aawconline.org/professional-resources/resources/>. Last accessed May 16, 2016.
6. National Pressure Ulcer Advisory Panel, European Pressure Ulcer Advisory Panel and Pan Pacific Pressure Injury Alliance. Prevention and Treatment of Pressure Ulcers: Clinical Practice Guideline. Emily Haesler (Ed.). Cambridge Media: Osborne Park, Australia; 2014.
7. Kantor J, Margolis DJ. A multicentre study of percentage change in venous leg ulcer area as a prognostic index of healing at 24 weeks. *Br J Dermatol* 2000;142:960-4.
8. Kantor J, Margolis DJ. Expected healing rates for chronic wounds. *Wounds* 2000;12(6):155-8.
9. Kantor J, Margolis DJ. Efficacy and prognostic value of simple wound measurements. *Arch Dermatol* 1998;134:157-4.
10. Gethin G. The importance of continuous wound measuring. *Wounds UK* 2006;2:60-8.
11. Wu SC, Driver VR, Wrobel JS, Armstrong DG. Foot ulcers in the diabetic patient, prevention, and treatment. *Vasc Health and Risk Management* 2007;3(1):65-76.
12. Milic DJ, Zivic SS, Bogdanovic DC, Karanovic ND, Golubovic ZV. Risk factors related to the failure of venous leg ulcers to heal with compression treatment. *J Vasc Surg* 2009;49:1242-7.
13. Bryant JL, Brooks TL, Schmidt B, Mostow EN. Reliability of wound measuring techniques in an outpatient wound center. *Ostomy Wound Manage* 2001;47(4):44-51.
14. Margolis DJ, Berlin JA, Brian L. Risk factors associated with the failure of a venous leg ulcer to heal. *Arch Derm* 1999;135:920-6.
15. Gelfand JM, Hoffstad O, Margolis DJ. Surrogate Endpoints for the treatment of venous leg ulcers. *J Invest Dermatol* 2002;119:1420-25.
16. Warriner R, Synder R, Cardinal M. Subsequent delayed progress in diabetic wounds receiving standards care that meet initial positive area reduction predictive of healing at four weeks. Presented at Diabetic Global Conference, Los Angeles, California, March 18-20, 2010.
17. Novitas: Local Coverage Determination for Application of Bioengineered Skin Substitutes to Lower Extremity Chronic Non-Healing Wounds (L35041). [https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=35041&ContrId=339&ver=28&ContrVer=1&CntcrtrSelected=339\\*\\*1&Cntcrctr=339&s=All&DocType=Active%7cFuture&ac=AAAGAAIAAAAAAA%3d%3d&LstAccesedJune21,2016](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=35041&ContrId=339&ver=28&ContrVer=1&CntcrtrSelected=339**1&Cntcrctr=339&s=All&DocType=Active%7cFuture&ac=AAAGAAIAAAAAAA%3d%3d&LstAccesedJune21,2016).