

Point Set Topology

Hoyan Mok

April 10, 2020

Contents

1	Topological Spaces and Continuous Mappings	2
1.1	Metric Space	2
1.2	Topological Space	4
1.3	Neighbourhood	5
1.4	Continuous Mappings	7
1.5	Closure	7
1.6	Interior Points and Boundary	10
1.7	Basis	11
2	Subspace	12

1 Topological Spaces and Continuous Mappings

1.1 Metric Space

Definition 1.1. function

$$d: X^2 \rightarrow \mathbb{R} \quad (1-1)$$

$\forall x_1, x_2, x_3 \in X$ satisfied:

- a) $d(x_1, x_2) = 0 \Leftrightarrow x_1 = x_2$;
- b) $d(x_1, x_2) = d(x_2, x_1)$ (symmetry);
- c) $d(x_1, x_3) \leq d(x_1, x_2) + d(x_2, x_3)$ (Triangle inequality),

is called a **metric** or **distance** in X . Such X is said to be equipped with metric d , (X, d) is called a **metric space**.

Some examples:

- $(\mathbb{R}^n; d_p)$, where $d_p(x_1, x_2) = \left(\sum_{i=1}^n |x_1^i - x_2^i|^p \right)^{1/p}$, while $d_\infty(x_1, x_2) = \max_{1 \leq i \leq n} |x_1^i - x_2^i|$.
- Similarly we can define metric spaces as $(C[a, b]; d_p)$ or $C_p[a, b]$. $d_p(f, g) = \left(\int_a^b |f - g|^p dx \right)^{\frac{1}{p}}$. C_∞ is called a **Chebyshev metric**.
- On class $\tilde{\mathfrak{R}}[a, b]$ over $\mathfrak{R}[a, b]$ similar metric can be defined. Functions are considered of one same class if they are equivalent except on a set not larger than null set.

Hilbert space denoted by $(\mathbb{H}; d)$ is defined as:

$$\mathbb{H} = \left\{ x = (x_1, x_2, \dots) \mid \forall i \in \mathbb{Z}_+ \left(\forall x_i \in \mathbb{R} \wedge \sum_{i=1}^{\infty} x_i^2 < \infty \right) \right\} \quad (1-2)$$

equipped with a metric d :

$$d: \mathbb{H}^2 \rightarrow \mathbb{R}; x, y \mapsto \sqrt{\sum_{i=1}^{\infty} (x_i - y_i)^2}. \quad (1-3)$$

To justify this definition, we need to introduce a lemma:

Lemma 1.

$$\forall n \in \mathbb{Z} \forall u \in \mathbb{R}^n \forall v \in \mathbb{R}^n \left(\sum_{i=1}^n u_i v_i \leq \sqrt{\sum_{i=1}^n u_i^2} \sqrt{\sum_{i=1}^n v_i^2} \right) \quad (1-4)$$

This is called **Schwarz inequality**.

Proof. If $\forall i = 1, \dots, n (v_i = 0)$ the equivalence has already been satisfied, therefore the following considers the situation that $\exists i \in \{1, \dots, n\} (v_i \neq 0)$. $\forall \lambda \in \mathbb{R}$, the quadratic polynomial of λ

$$\sum_{i=1}^n (u_i + \lambda v_i)^2 = \sum_{i=1}^n u_i^2 + 2\lambda \sum_{i=1}^n u_i v_i + \lambda^2 \sum_{i=1}^n v_i^2 \geq 0$$

has at most one root. Hence $\Delta \leq 0$ will lead to the inequality 1-4. \square

Apply this inequality to $\sum_{i=1}^n (u_i + v_i)^2 = \sum_{i=1}^n u_i^2 + \sum_{i=1}^n v_i^2 + 2 \sum_{i=1}^n v_i \sum_{i=1}^n u_i$ we can get

$$\sum_{i=1}^n (u_i + v_i)^2 \leq \sum_{i=1}^n u_i^2 + \sum_{i=1}^n v_i^2 + 2 \sqrt{\sum_{i=1}^n u_i^2} \sqrt{\sum_{i=1}^n v_i^2} = \left(\sqrt{\sum_{i=1}^n u_i^2} + \sqrt{\sum_{i=1}^n v_i^2} \right)^2,$$

in which substitute u_i, v_i by $x_i - y_i, x_i + y_i$ will result in triangle inequality. The inequality holds as the n limits to $+\infty$.

Definition 1.2. Let (X, d) be a metric space, the **distant** between a non-empty set $\emptyset \neq A \in \mathcal{P}(X)$ and a point x is defined as:

$$d(A, x) = \inf\{d(x, y) \mid y \in A\},$$

and we let $d(x, A) = d(A, x)$. Also, the **distant** between two non-empty sets A, B is defined as:

$$d(A, B) = \inf\{d(x, y) \mid x \in A \wedge y \in B\}.$$

A metric space (X, d) is called **discrete** if

$$\forall x \in X \left(\exists \delta_x \in \mathbb{R}_+ (\forall y \in X (y \neq x \rightarrow d(x, y) > \delta_x)) \right).$$

Lemma 2. If (X, d) is a metric space, then $\forall a, b, u, v, |d(a, b) - d(u, v)| \leq d(a, u) + d(b, v)$.

Proof. Without loss of generality, we assume that $d(a, b) > d(u, v)$. According to the triangle inequality (see def. 1.1), $d(a, b) \leq d(a, u) + d(u, v) + d(v, b)$, which is to prove. \square

Definition 1.3. $\delta \in \mathbb{R}_+, a \in X$. Set

$$B(a; \delta) = \{x \in X \mid d(a, x) < \delta\}$$

is then called a **ball** with centre $a \in X$, and a radius of δ , or a **δ -ball** of point a .

Definition 1.4. An **open set** $G \subset X$ in metric space (X, d) satisfies: $\forall x \in G, \exists B(x; \delta)$, s.t. $B(x; \delta) \subset G$.

Definition 1.5. A set $F \subset X$ in metric space (X, d) is said to be a **closed set** if its complement $\mathbb{C}_X(F)$ is open.

It can be proved that \emptyset and X itself is both open and closed.

Proposition 1. a) An infinite union of open sets is an open set.

b) A finite intersection of open sets is an open set.

c) A finite union of closed sets is a closed set.

d) An infinite intersection of closed sets is a closed set.

Proof. a) If open sets $G_\alpha \subset X, \forall \alpha \in A, \forall a \in \bigcap_{\alpha \in A} G_\alpha, \exists \alpha_0 \in A, a \in G_{\alpha_0},$
 $\exists B(a; \delta) \subset G_{\alpha_0} \subset \bigcap_{\alpha \in A} G_\alpha.$

b) Open sets $G_1 \cup G_2 \subset X, a \in G_1 \cap G_2$, therefore $\exists \delta_1, \delta_2 \in \mathbb{R}_+, B(a; \delta_1) \subset G_1, B(a; \delta_2) \subset G_2$, without loss of generality, let $\delta_1 \geq \delta_2, a \in B(a; \delta_1) \cap B(a; \delta_2) = B(a; \delta_2) \subset G_1 \cap G_2.$

c) Just consider $\mathbb{C}_X(\bigcap_{\alpha \in A} F_\alpha) = \bigcup_{\alpha \in A} \mathbb{C}_X(F_\alpha)$ and a).

d) Similarly, $\mathbb{C}_X(F_1 \cup F_2) = \mathbb{C}_X(F_1) \cap \mathbb{C}_X(F_2).$

□

1.2 Topological Space

Definition 1.6. We say X is equipped with a **topological space** or equipped with **topology** if we assigned a $\mathcal{T} \subset 2^X$, which has got the following propoties:

a) $\emptyset \in \mathcal{T}; X \in \mathcal{T}.$

b) $\forall \alpha \in A (G_\alpha \in \mathcal{T}) \rightarrow \bigcup_{\alpha \in A} G_\alpha \in \mathcal{T}.$

c) $G_1 \in \mathcal{T} \wedge G_2 \in \mathcal{T} \rightarrow G_1 \cap G_2 \in \mathcal{T}.$

Then we call (X, \mathcal{T}) a **topological space**. Every $G \in \mathcal{T}$ is called an **open set**.

Definition 1.7. A topology \mathcal{T}_d insisting of the open sets in a metric space (X, d) is called a **topology induced by metric d**.

A trivial example of topological space is **trivial topology**, which consists only of empty set and the space itself, i.e. $\mathcal{T} = \{\emptyset, X\}$. Another trivial example of topological space is **discrete topology**, which consists of all the subsets of the space i.e. $\mathcal{T} = 2^X$.

A **cofinite space** is a base set X equipped with a topology \mathcal{T} defined as follows:

$$\mathcal{T} = \{U \in 2^X \mid U = \emptyset \vee \mathbb{C}_X U \text{ is finite}\} \quad (1-5)$$

Proposition 2. The set \mathcal{T} under definition 1-5 is a topology.

Proof. a) $\emptyset \in \mathcal{T}, X \in \mathcal{T}$.

b) $\forall i \in I (|\mathbb{C}_X A_i| \in \mathbb{N}) \rightarrow \forall i_0 \in I (|\bigcap_{i \in I} \mathbb{C}_X A_i| \leq |\mathbb{C}_X A_{i_0}|)$, therefore $\bigcup_{i \in I} A_i \in \mathcal{T}$.

c) $\forall A \in \mathcal{T} \forall B \in \mathcal{T} (A \cap B = \emptyset \in \mathcal{T} \vee \mathbb{C}_X(A \cap B) = \mathbb{C}_X A \cup \mathbb{C}_X B \text{ is finite})$, therefore $\forall A \in \mathcal{T} \forall B \in \mathcal{T} (A \cap B \in \mathcal{T})$. \square

Similarly, **countable complement space** can be defined.

Let X be equiped with two topology $\mathcal{T}_1, \mathcal{T}_2$. $\mathcal{T}_1 \cup \mathcal{T}_2$ is possibly not a topology of X . For example, $\mathcal{T}_1 = \{(x, +\infty) \mid x \in \mathbb{R}\} \cup \{\emptyset, \mathbb{R}\}$ and $\mathcal{T}_2 = \{(-\infty, y) \mid y \in \mathbb{R}\} \cup \{\emptyset, \mathbb{R}\}$ are both topologies of \mathbb{R} , but there union $\mathcal{T}_1 \cup \mathcal{T}_2$ is not.

Theorem 1.1. Let X be equiped with two topology $\mathcal{T}_1, \mathcal{T}_2$. Their intersection $\mathcal{T}_1 \cap \mathcal{T}_2$ is also a topology on X .

Proof. a) $\{\emptyset, X\} \subseteq \mathcal{T}_1 \wedge \{\emptyset, X\} \subseteq \mathcal{T}_2 \rightarrow \{\emptyset, X\} \subseteq \mathcal{T}_1 \cap \mathcal{T}_2$.

b) $\forall \alpha \in A (G_\alpha \in \mathcal{T}_1 \cap \mathcal{T}_2) \rightarrow \bigcup_{\alpha \in A} G_\alpha \in \mathcal{T}_1 \wedge \bigcup_{\alpha \in A} G_\alpha \in \mathcal{T}_2$.

c) $\forall G_1 \in \mathcal{T}_1 \cap \mathcal{T}_2 \forall G_2 \in \mathcal{T}_1 \cap \mathcal{T}_2 (G_1 \cap G_2 \in \mathcal{T}_1 \wedge G_1 \cap G_2 \in \mathcal{T}_2)$ \square

Definition 1.8. Let (X, \mathcal{T}) be a topological space. If there exists a metric $d: X^2 \rightarrow \mathbb{R}$ s.t. (X, \mathcal{T}) is induced by d then call (X, \mathcal{T}) a **metrizable space**, (X, d) is its **metrization**.

1.3 Neighbourhood

Definition 1.9. Let (X, \mathcal{T}) be a topological space. A set $U(x)$ is said to be a **neighbourhood** of a point $x \in X$ if $\exists G \in \mathcal{T} (G \subseteq U(x) \wedge x \in G)$. If $U(x) \in \mathcal{T}$, it is called an **open neighbourhood**. Subset class $\{U(x) \subseteq X \mid U(x) \text{ is a neighbourhood of } x\}$ is called the **neighbourhood system** of point x , denoted by \mathcal{U}_x .

Theorem 1.2. Let (X, \mathcal{T}) be a topological space, U is a subset of X . U is an open set iff $\forall x \in U, U$ is a neighbourhood of x .

Proof. The necessity is trivial. $\forall x \in U \exists V(x)$ s.t. $V(x)$, being a subset of U , is an open neighbourhood of x . By definition of topology, $\bigcup_{x \in U} V(x) \in \mathcal{T}$. $\forall x \in U (x \in \bigcup_{x \in U} V(x)) \rightarrow U \subseteq \bigcup_{x \in U} V(x)$, while $\forall x \in U (V(x) \subseteq U) \rightarrow \bigcup_{x \in U} V(x) \subseteq U$, therefore $U = \bigcup_{x \in U} V(x) \in \mathcal{T}$. \square

Theorem 1.3. Let (X, \mathcal{T}) be a topological space, \mathcal{U}_x is a neighbourhood system of point $x \in X$.

$$\forall U \in \mathcal{U}_x \forall V \in \mathcal{U}_x (U \cap V \in \mathcal{U}_x)$$

Proof. $\forall U \in \mathcal{U}_x \forall V \in \mathcal{U}_x \exists U_0 \in \mathcal{T} \exists V_0 \in \mathcal{T} (U_0 \subseteq U \wedge V_0 \subseteq V \wedge x \in U_0 \cap V_0)$,
By definition of topology, $\mathcal{T} \ni U_0 \cap V_0 \subseteq U \cap V$. \square

In history topologies were once built on neighbourhood systems. The following theorem shows the way.

Theorem 1.4. Let X be a set and $\forall x \in X$ a collection of subsets $\mathcal{U}_x \in \mathcal{P}(X)$ is appointed, satisfying:

- (1) $\forall x \in X (\mathcal{U}_x \neq \emptyset \wedge \forall U \in \mathcal{U}_x (x \in U))$;
- (2) $\forall x \in X \forall U \in \mathcal{U}_x \forall V \in \mathcal{U}_x (U \cap V \in \mathcal{U}_x)$;
- (3) $\forall x \in X \forall U \in \mathcal{U}_x \forall V \in \mathcal{P}(X) (U \subseteq V \rightarrow V \in \mathcal{U}_x)$;
- (4) $\forall x \in X \forall U \in \mathcal{U}_x \exists V \in \mathcal{U}_x (V \subseteq U \wedge \forall y \in V (V \in \mathcal{U}_y))$,

then there exists only one topology \mathcal{T} on X s.t. $\forall x \in X$, \mathcal{U}_x is the neighbourhood system of x in (X, \mathcal{T}) .

Proof. Let $\mathcal{T} = \{G \in \mathcal{P}(X) \mid \forall x \in G (G \in \mathcal{U}_x)\}$.

- a) Obviously $\emptyset \in \mathcal{T}$. Since the condition (1) and the condition (3) in theorem 1.4, $X \in \mathcal{T}$.
- b) Let $A, B \in \mathcal{T}$. Consider the condition (2) in theorem 1.4 applied to $x \in A \cap B$.
- c) Let $\forall i \in I (G_i \in \mathcal{T})$. $\forall x \in \bigcup_{i \in I} G_i$, there must exists a $i \in I$ s.t. $x \in G_i$ and $G_i \in \mathcal{U}_x$. Since the condition (3) in theorem 1.4, $G_i \subseteq \bigcup_{i \in I} G_i$ has implied $\bigcup_{i \in I} G_i \in \mathcal{U}_x$.

These tells that \mathcal{T} is a topology on X .

The condition (4) in theorem 1.4 tells that there always exists a $G \subset U$ for all $x \in X$ and $U \in \mathcal{U}_x$ s.t. $G \in \mathcal{T}$. Therefore \mathcal{U}_x must be a subset of the neighbourhood system of x .

For all neighbourhood U of $x \in X$, there must be a open neighbourhood subset $G \subseteq U$, which is also a member of \mathcal{U}_x . Since the condition (3) in theorem 1.4, $U \in \mathcal{U}_x$. Therefore the neighbourhood system of x must be a subset of \mathcal{U}_x .

Therefore, \mathcal{U}_x is the neighbourhood system of x .

Now prove the uniqueness. Let there be another topology \mathcal{T}' . Since theorem 1.2, $\forall U (G \in \mathcal{T}' \leftrightarrow \forall x \in G (G \in \mathcal{U}_x))$. Therefore $\mathcal{T}' = \mathcal{T}$. \square

1.4 Continuous Mappings

Definition 1.10. A mapping $f: X \rightarrow Y$, where X, Y is respectively equipped with topology $\mathcal{T}_X, \mathcal{T}_Y$, is said to be **continuous** at $x_0 \in X$ (let $y_0 = f(x_0) \in Y$), if $\forall U(y_0), \exists U(x_0)$ s.t. $f(U(x_0)) \subset U(y_0)$. It is **continuous** in X if it is continuous at each point $x \in X$.

The set of continuous mappings from X into Y can be denoted by $C(X, Y)$ or $C(X)$ when Y is clear.

It can be easily proved that an identify function $e_X: X \rightarrow X$ where X is equipped with a topology \mathcal{T} is a continuous function.

Theorem 1.5. (criterion of continuity)

Let (X, \mathcal{T}) , (Y, \mathcal{S}) be two topological space. A mapping $f: X \rightarrow Y$ is continuous iff

$$\forall V \in \mathcal{S} (\exists U \in \mathcal{T} (U = f^{-1}(V))).$$

Proof. \rightarrow : It is obvious if $f^{-1}(G_Y) = \emptyset$. If $f^{-1}(G_Y) \neq \emptyset$ and $\forall x_0 \in f^{-1}(G_Y)$, since $f \in C(X, Y)$, for $G_Y \in \mathcal{S}$, $\exists U(x_0)$ s.t. $f(U(x_0)) \subset G_Y$. Also notice that $f(U(x_0)) \subset G_Y \Rightarrow U(x_0) \subset f^{-1}(G_Y)$, therefore $f^{-1}(G_Y)$ is open (Theorem 1.2).

\leftarrow : $\forall x_0 \in X$, let $y_0 = f(x_0)$, $f^{-1}(U(y_0)) \in \mathcal{T}$ if $U(y_0) \in \mathcal{S} \cap \mathcal{U}_{y_0}$. Notice that $x_0 \in f^{-1}(U(y_0))$, $f^{-1}(U(y_0))$ is a neighbourhood of x_0 , therefore $f \in C(X, Y)$. \square

Theorem 1.6. Let (X, \mathcal{T}_X) , (Y, \mathcal{T}_Y) , (Z, \mathcal{T}_Z) be topological spaces. If $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are both continuous, $g \circ f: X \rightarrow Z$ is also continuous.

Proof.

$$\forall W \in \mathcal{T}_Z (g^{-1}(W) \in \mathcal{T}_Y) \rightarrow \forall W \in \mathcal{T}_Z (f^{-1}(g^{-1}(W)))$$

Since $f^{-1}(g^{-1}(W)) = (g \circ f)^{-1}(W)$, the theorem has been proved. \square

Definition 1.11. (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) are both topological spaces. A bijective mapping $f: X \rightarrow Y$ is a **homeomorphism** if $f \in C(X, Y) \wedge f^{-1} \in C(Y, X)$.

Definition 1.12. Two topological spaces (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) are said to be **homeomorphic** if there exists a homeomorphism $f: X \rightarrow Y$.

Homeomorphic topological spaces are identical with respect to their topological properties since the theorem 1.5 has shown that their open sets correspond to each other. In fact homeomorphic relations are equivalent relations.

1.5 Closure

Definition 1.13. Let X be a topological space and A be a subset of X . Let $x \in X$. If $\forall U \in \mathcal{U}_x (U \cap (A - \{x\}) \neq \emptyset)$, then x is called a **accumulation point**, **cluster point** or **limit point** of A . The set $A' := \{x \in X \mid x \text{ is an accumulation point of } A\}$ is called the **derived set** of A . A point $a \in A$ is called a **isolated point** of A if $a \notin A'$.

Theorem 1.7. Let X be a topological space and A, B be subsets of X . 1) $A \subseteq B \rightarrow A' \subseteq B'$; 2) $(A \cup B)' = A' \cup B'$; 3) $(A')' \subseteq A \cup A'$.

Proof. 1) When $A \subseteq B$, $U \cap (A - \{x\}) \subseteq U \cap (B - \{x\})$.

2) $(A \cup B)' = \{x \in X \mid \forall U \in \mathcal{U}_x (U \cap (A \cup B - \{x\}) \neq \emptyset)\}$. Also $U \cap (A \cup B - \{x\}) = U \cap (X - \{x\}) \cap (A \cup B) = (U \cap A - \{x\}) \cup (U \cap B - \{x\})$.

3) If $x \notin A \cup A'$, then $\exists G \in \mathcal{U}_x \cap \mathcal{T} (G \cap (A - \{x\}) = G \cap A = \emptyset)$. $\forall y \in G$, G itself is a neighbourhood of y that $G \cap (A - \{y\}) = G \cap A = \emptyset$, therefore $y \notin A'$. This means that G is a neighbourhood of x that $G \cap (A' - \{x\}) = G \cap A' = \emptyset$, i.e. $x \notin (A')'$. □

Definition 1.14. Let (X, \mathcal{T}) be a topological space and F be a subset of X . F is said to be **closed** iff $\mathcal{C}_X(F) \in \mathcal{T}$. The collection all closed sets is denoted by \mathcal{F} .

Theorem 1.8. Let (X, \mathcal{T}) be a topological space and F be a subset of X . F is closed iff $F' \subseteq F$.

Proof. \rightarrow : If $x \notin F$ then $x \in \mathcal{C}_X(F)$, which is open in (X, \mathcal{T}) . Then $\mathcal{C}_X(F)$ is a neighbourhood that $\mathcal{C}_X(F) \cap (F - \{x\}) = \mathcal{C}_X(F) \cap F = \emptyset$, i.e. $x \notin F'$.

\leftarrow : $\forall x \notin F (x \notin F')$, then there exists an open neighbourhood U of x that $U \cap F = \emptyset$, then $\mathcal{C}_X(F)$ is always a neighbourhood of its elements, since theorem 1.2, $\mathcal{C}_X(F) \in \mathcal{T}$. □

Definition 1.15. Let (X, \mathcal{T}) be a topological space and A be a subset of X . Set $\overline{A} := A \cup A'$ is called a **closure** of A .

Theorem 1.9. Let (X, \mathcal{T}) be a topological space and A be a subset of X . Let $x \in X$.

$$x \in \overline{A} \leftrightarrow \forall U \in \mathcal{U}(x) (U \cap A \neq \emptyset).$$

Proof. \rightarrow : If $x \in A$ then $\{x\} \subseteq U \cap A$, else if $x \in A'$ then $U \cap A \supset (U - \{x\}) \cap A \neq \emptyset$.

\leftarrow : If $\exists U \in \mathcal{U}(x) ((U - \{x\}) \cap A = \emptyset) \wedge x \notin A$, then there exists a $U \in \mathcal{U}(x)$ s.t. $U \cap A = \emptyset$. □

Theorem 1.10. Let (X, \mathcal{T}) be a topological space and A be a subset of X . A is closed in (X, \mathcal{T}) iff $A = \overline{A}$.

Proof. Since theorem 1.8, A is closed iff $A' \subseteq A$, which iff $A = A \cup A' = \overline{A}$. □

Corollary 1. Let (X, \mathcal{T}) be a topological space and A be a subset of X . \overline{A} is always closed.

Proof. Since (3) of theorem 1.7, $\overline{\overline{A}} = \overline{A}$. □

Lemma 3. Let (X, \mathcal{T}) be a topological space and A, B be subsets of X . $A \subseteq B \rightarrow \overline{A} \subseteq \overline{B}$.

Proof. $A \subseteq B \rightarrow A' \subseteq B'$ ((1) of theorem 1.7), so $A \cup A' \subseteq B \cup B'$, i.e. $\overline{A} \subseteq \overline{B}$. \square

We can say that the closure of a set is the smallest closed set containing it, as long as we prove the following theorem:

Theorem 1.11. *Let (X, \mathcal{T}) be a topological space and A be a subset of X .*

$$\overline{A} = \bigcap_{F \in \mathcal{T} \wedge A \subseteq F} F.$$

Proof. Since \overline{A} itself is closed (corollary 1), $\bigcap_{F \in \mathcal{T} \wedge A \subseteq F} F \subseteq \overline{A}$. On the other hand, $\bigcap_{F \in \mathcal{T} \wedge A \subseteq F} F$ is closed, so $\overline{\bigcap_{F \in \mathcal{T} \wedge A \subseteq F} F} = \bigcap_{F \in \mathcal{T} \wedge A \subseteq F} F$. Therefore, $A \subseteq \bigcap_{F \in \mathcal{T} \wedge A \subseteq F} F \rightarrow \overline{A} \subseteq \bigcap_{F \in \mathcal{T} \wedge A \subseteq F} F$ (Lemma 3). \square

Theorem 1.12. *Let (X, d) be a metric space and A be a non-empty subset of X .*

1) $\forall x \in X, x \in A' \leftrightarrow d(x, A - \{x\}) = 0$.

2) $\forall x \in X, x \in \overline{A} \leftrightarrow d(x, A) = 0$.

Proof. 1) We have $x \in A'$ iff $\forall \varepsilon \in \mathbb{R}_+ (B(x, \varepsilon) \cap (A - \{x\}) \neq \emptyset)$, which is established iff $\forall \varepsilon \in \mathbb{R}_+ \exists y \in A - \{x\} (d(x, y) < \varepsilon)$.

2) We only need to substitute $A - \{x\}$ with A in 1). \square

Theorem 1.13. *Let (X, \mathcal{T}) and (Y, \mathcal{S}) be two topological spaces, and $f: X \rightarrow Y$. Note the collections of closed sets in X and Y by $\mathcal{F}_X, \mathcal{F}_Y$. The statements below are equivalent:*

(1) $f \in C(X, Y)$.

(2) $\forall F \in \mathcal{F}_Y (f^{-1}(F) \in \mathcal{F}_X)$.

(3) $\forall A \in \mathcal{P}(X) (f(\overline{A}) \subseteq \overline{f(A)})$.

(4) $\forall B \in \mathcal{P}(Y) (\overline{f^{-1}(B)} \subseteq f^{-1}(\overline{B}))$.

Proof. (1) \rightarrow (2): Only to notice that $f^{-1}(\mathcal{C}_Y F) = \mathcal{C}_X f^{-1}(F)$.

(2) \rightarrow (3): $\forall A \in \mathcal{P}(X)$ we have $f(A) \subseteq \overline{f(A)}$, so $A \subseteq f^{-1}(\overline{f(A)})$. By (2) we know that $f^{-1}(\overline{f(A)})$ is closed, therefore $\overline{A} \subseteq f^{-1}(\overline{f(A)}) = \overline{f^{-1}(f(A))}$, so $f(\overline{A}) \subseteq \overline{f(A)}$.

(3) \rightarrow (4): By (3) we know that $\forall B \in \mathcal{P}(Y), f(\overline{f^{-1}(B)}) \subseteq \overline{f(f^{-1}(B))}$. Also $f(f^{-1}(B)) \subseteq B$ (equality satisfied when f is surjective), then $f(\overline{f^{-1}(B)}) \subseteq \overline{B}$, so $\overline{f^{-1}(B)} \subseteq f^{-1}(\overline{B})$.

(4) \rightarrow (1): $\forall G \in \mathcal{S}, \mathbb{C}_Y G \in \mathcal{F}_Y$, so by (3), we have

$$\overline{\mathbb{C}_X f^{-1}(G)} = \overline{f^{-1}(\mathbb{C}_Y G)} \subseteq f^{-1}(\overline{\mathbb{C}_Y G}) = f^{-1}(\mathbb{C}_Y G) = \mathbb{C}_X f^{-1}(G).$$

However by the definition of closure $\mathbb{C}_X f^{-1}(G) \subseteq \overline{\mathbb{C}_X f^{-1}(G)}$. Therefore $\mathbb{C}_X f^{-1}(G) = \overline{\mathbb{C}_X f^{-1}(G)}$, which means $\mathbb{C}_X f^{-1}(G)$ is closed (theorem 1.10), i.e. $f^{-1}(G)$ is open. \square

1.6 Interior Points and Boundary

Definition 1.16. Let (X, \mathcal{T}) be a topological space and A be a subset of X . We call x an **interior point** of A if A is a neighbourhood of x . We call x an **exterior point** of A if $\mathbb{C}_X(A)$ is a neighbourhood of x . The sets of all interior points of A is the **interior** of A , noted by $\text{int } A$.

Theorem 1.14. Let (X, \mathcal{T}) be a topological space and A be a subset of X . $\text{int } A = \mathbb{C}_X(\mathbb{C}_X(A))$, $\overline{A} = \mathbb{C}_X(\text{int } \mathbb{C}_X(A))$.

Proof. $x \in \text{int } A$ implies an open set G which is a subset of A and $x \in G$. The complement of G is closed, thus its closure is $\mathbb{C}_X(G)$ itself. $\mathbb{C}_X(A) \subseteq \mathbb{C}_X(G) \rightarrow \overline{\mathbb{C}_X(A)} \subseteq \mathbb{C}_X(G)$ (Lemma 3), therefore $x \notin \overline{\mathbb{C}_X(A)}$, which is the first equation to prove.

To prove the second one only need to replace the A with $\mathbb{C}_X(A)$ in the first equation. \square

Theorem 1.15. Let (X, \mathcal{T}) be a topological space and G be a subset of X .

$$G \in \mathcal{T} \leftrightarrow \text{int } G = G$$

Proof. G is open iff the complement of G is closed. And $\mathbb{C}_X(G) = \overline{\mathbb{C}_X(G)}$. The complement of the both size of this equation and Theorem 1.14 give the proof of the theorem. \square

With the propoties of closure and Theorem 1.14 the folowing statements should be easy to prove:

Theorem 1.16. Let (X, \mathcal{T}) be a topological space and A, B be subsets of X . $\text{int}(A \cap B) = \text{int } A \cap \text{int } B$, $\text{int}(\text{int } A) = \text{int } A$,

$$\text{int } A = \bigcup_{G \in \mathcal{T} \wedge G \subseteq A} G.$$

Therefore we can say that the interior of A is the largest open set contained in A .

Definition 1.17. Let (X, \mathcal{T}) be a topological space and A be a subset of X . A point x is said to be a **boundary point** of A if $\forall U \in \mathcal{U}(x) (U \cap A \neq \emptyset \wedge U \cap \mathbb{C}_X(A) \neq \emptyset)$. The set of all boundary points of A is called the **boundary** of A , noted by ∂A .

Theorem 1.17. Let (X, \mathcal{T}) be a topological space and A be a subset of X . (1) $\partial A = \overline{A} \cap \overline{\mathbb{C}_X(A)}$; (2) $\text{int } A = \overline{A} - \partial A$; (3) $\overline{A} = \text{int } A \cup \partial A$.

Proof. (1): Apply Theorem 1.9 to both A and $\mathbb{C}_X(A)$.

(2): $\text{int } A \cup \partial A = A \cup (\overline{A} \cap \overline{\mathbb{C}_X(A)}) = \overline{A} \cap (\text{int } A \cup \mathbb{C}_X(\text{int } A)) = \overline{A}$.

(3) $\overline{A} - \partial A = \overline{A} - (\overline{A} \cap \overline{\mathbb{C}_X(A)}) = \overline{A} \cap \mathbb{C}_X(\overline{\mathbb{C}_X(A)}) = \overline{A} \cap \text{int } A = \text{int } A$. \square

1.7 Basis

Definition 1.18. Let (X, \mathcal{T}) be a topological space and \mathcal{B} be a subset of \mathcal{T} . If $\forall G \in \mathcal{T} \exists \mathcal{B}_G \in \mathcal{P}(\mathcal{B})(G = \cup \mathcal{B}_G)$, then we call \mathcal{B} a **basis** or a **base** of the topology \mathcal{T} .

Theorem 1.18. Let (X, \mathcal{T}) be a topological space and \mathcal{B} be a subset of \mathcal{T} . \mathcal{B} is a basis of \mathcal{T} iff $\forall x \in X \forall U \in \mathcal{U}_x \exists B \in \mathcal{B}(x \in B \wedge B \subseteq U)$.

Proof. \rightarrow : $U \in \mathcal{U}_x$ implies an open set $G \in \mathcal{P}(U)$ that contains x , which is the union of elements in \mathcal{B} . Therefore there exists a $B \in \mathcal{B}$, which is the subset of U and it contains x .

\leftarrow : $\forall G \in \mathcal{T}$, it is a neighbourhood of all the points in G . For all x in G assign a $B_x \in \mathcal{B}$ which contains x and is the subset of G , so that G is the Union of all the B_x . \square

Theorem 1.19. Let \mathcal{B} be a basis of a topological space (X, \mathcal{T}) .

$$\forall B_1 \in \mathcal{B} \forall B_2 \in \mathcal{B} \forall x \in B_1 \cap B_2 \exists B \in \mathcal{B}(x \in B \subseteq B_1 \cap B_2).$$

Proof. By definition of topology and basis, B_1 and B_2 are both open and their intersection $B_1 \cap B_2$ is open as well. Then there exists a collection of sets in \mathcal{B} whose union is $B_1 \cap B_2$, there must be at least a set B that contains x . \square

The topology \mathcal{T} on X is determined if the basis \mathcal{B} is given, that is, if the union of \mathcal{B} is X and it satisfies Theorem 1.19, then \mathcal{T} , which is defined by the collection of the unions of B s in \mathcal{B} , is the *only* topology on X such that \mathcal{B} is a basis of it.

For example, **lower limit topology** \mathcal{T}_ℓ on \mathbb{R} is defined by giving a basis:

$$\mathcal{B}_\ell = \{[a, b) \mid a, b \in \mathbb{R} \wedge a < b\},$$

and $(\mathbb{R}, \mathcal{T}_\ell)$ is called the **lower limit topological space**, the **Sorgenfrey line** or the **arrow**, denoted by \mathbb{R}_ℓ .

It is obvious that $\mathcal{T} \subsetneq \mathcal{T}_\ell$.

Definition 1.19. Let (X, \mathcal{T}) be a topological space and \mathcal{S} be a subset of \mathcal{T} . If the collection \mathcal{B} of the finite intersections of the non-empty sets in \mathcal{S} is a basis of \mathcal{T} , i.e.

$$\mathcal{B} = \left\{ \bigcap_{i=1}^n S_i \mid S_i \in \mathcal{S}, i = 1, \dots, n, n \in \mathbb{N}_+ \right\},$$

then we call \mathcal{S} a **subbasis** or a **subbase** of \mathcal{T} .

2 Subspace

Index

- δ -ball, 3
- Symbols used
 - d , 2
 - \mathbb{H} , 2
 - \mathcal{T} , 4
- accumulation point, 7
- arrow, 11
- ball, 3
- base, 11
- basis, 11
- boundary, 10
- boundary point, 10
- Chebyshev metric, 2
- closed, 8
- closed set, 4
- closure, 8
- cluster point, 7
- cofinite space, 4
- continuous, 7
- countable complement space, 5
- criterion of continuity, 7
- derived set, 7
- discrete, 3
- discrete topology, 4
- distance, 2
- distant, 3
- exterior point, 10
- Hilbert space, 2
- homeomorphic, 7
- homeomorphism, 7
- interior, 10
- interior point, 10
- isolated point, 7
- limit point, 7
- lower limit topological space, 11
- lower limit topology, 11
- metric, 2
- metric space, 2
- metrizable space, 5
- metrization, 5
- neighbourhood, 5
- neighbourhood system, 5
- open neighbourhood, 5
- open set, 3, 4
- Schwarz inequality, 2
- Sorgenfrey line, 11
- subbase, 11
- subbasis, 11
- topological space, 4
- topology, 4
- topology induced by metric, 4
- trivial topology, 4