

Matrix Theory Assignment 1

Hrithik Raj

Abstract—This document contains the solution to problem No.66 from Lines and Planes

1 PROBLEM STATEMENT

$$\text{Simplify } \mathbf{z} = \left(\frac{1}{\begin{pmatrix} 1 \\ -4 \end{pmatrix}} - \frac{2}{\begin{pmatrix} 2 \\ 1 \end{pmatrix}} \right) \frac{\begin{pmatrix} 3 \\ -4 \end{pmatrix}}{\begin{pmatrix} 5 \\ 1 \end{pmatrix}}.$$

2 SOLUTION

In order to simplify the above equation, we need to find the multiplicative inverse of the sub - matrices.

We, first define the multiplicative inverse of a matrix of a complex number. Let \mathbf{T}_a be the matrix of the complex number a , then b is defined to be the multiplicative inverse of a if

$$\mathbf{T}_a \mathbf{T}_b = \mathbf{I} \quad (2.0.1)$$

$$\mathbf{b} = \mathbf{a}^{-1} = \begin{pmatrix} a_1 & -a_2 \\ a_2 & a_1 \end{pmatrix}^{-1} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (2.0.2)$$

$$= \frac{1}{\|a\|^2} \begin{pmatrix} a_1 \\ -a_2 \end{pmatrix} \quad (2.0.3)$$

$$\mathbf{z} = \left(\begin{pmatrix} 1 \\ -4 \end{pmatrix}^{-1} - 2 \begin{pmatrix} 2 \\ 1 \end{pmatrix}^{-1} \right) \begin{pmatrix} 3 \\ -4 \end{pmatrix} \frac{1}{\begin{pmatrix} 5 \\ 1 \end{pmatrix}} \quad (2.0.4)$$

Using Eq (2.0.3),

$$= \left(\frac{1}{17} \begin{pmatrix} 1 \\ 4 \end{pmatrix} - \frac{2}{5} \begin{pmatrix} 2 \\ -1 \end{pmatrix} \right) \begin{pmatrix} 3 \\ -4 \end{pmatrix} \frac{1}{26} \begin{pmatrix} 5 \\ 1 \end{pmatrix} \quad (2.0.5)$$

$$= \left(\begin{pmatrix} \frac{1}{17} \\ \frac{4}{17} \end{pmatrix} - \begin{pmatrix} \frac{4}{5} \\ \frac{-2}{5} \end{pmatrix} \right) \begin{pmatrix} 3 \\ -4 \end{pmatrix} \begin{pmatrix} \frac{5}{26} \\ \frac{1}{26} \end{pmatrix} \quad (2.0.6)$$

$$= \left(\begin{pmatrix} 0.058 \\ 0.235 \end{pmatrix} - \begin{pmatrix} 0.8 \\ -0.4 \end{pmatrix} \right) \begin{pmatrix} 3 \\ -4 \end{pmatrix} \begin{pmatrix} 0.192 \\ -0.038 \end{pmatrix} \quad (2.0.7)$$

$$= \begin{pmatrix} 0.058 - 0.8 \\ 0.235 + 0.4 \end{pmatrix} \begin{pmatrix} 3 \\ -4 \end{pmatrix} \begin{pmatrix} 0.192 \\ -0.038 \end{pmatrix} \quad (2.0.8)$$

$$= \begin{pmatrix} -0.742 \\ 0.635 \end{pmatrix} \begin{pmatrix} 3 \\ -4 \end{pmatrix} \begin{pmatrix} 0.192 \\ -0.038 \end{pmatrix} \quad (2.0.9)$$

Using equivalent matrices for the complex numbers,

$$= \begin{pmatrix} -0.742 & -0.635 \\ 0.635 & -0.742 \end{pmatrix} \begin{pmatrix} 3 & 4 \\ -4 & 3 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \begin{pmatrix} 0.192 \\ -0.038 \end{pmatrix} \quad (2.0.10)$$

Matrix multiplication gives,

$$= \begin{pmatrix} -2.226 + 2.54 & -2.968 - 1.905 \\ 1.905 + 2.968 & 2.54 - 2.226 \end{pmatrix} \begin{pmatrix} 0.192 \\ -0.038 \end{pmatrix} \quad (2.0.11)$$

$$= \begin{pmatrix} 0.314 & -4.873 \\ 4.873 & 0.314 \end{pmatrix} \begin{pmatrix} 0.192 \\ -0.038 \end{pmatrix} \quad (2.0.12)$$

$$= \begin{pmatrix} 0.314 \\ 4.873 \end{pmatrix} \begin{pmatrix} 0.192 \\ -0.038 \end{pmatrix} \quad (2.0.13)$$

Again using equivalent and multiplying,

$$= \begin{pmatrix} 0.314 & -4.873 \\ 4.873 & 0.314 \end{pmatrix} \begin{pmatrix} 0.192 & 0.038 \\ -0.038 & 0.192 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (2.0.14)$$

$$= \begin{pmatrix} 0.245 & -0.9236 \\ 0.9236 & 0.245 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (2.0.15)$$

$$\Rightarrow \mathbf{z} = \begin{pmatrix} 0.245 \\ 0.9236 \end{pmatrix} \quad (2.0.16)$$

Python Code:

```
https://github.com/Hrithikraj2/
MatrixTheory_EE5609/blob/master/
Assignment_1/codes/A1_code1.py
```

```
https://github.com/Hrithikraj2/
MatrixTheory_EE5609/blob/master/
Assignment_1/codes/A1_code4.py
```

Latex codes:

[https://github.com/Hrithikraj2/
MatrixTheory_EE5609/blob/master/
Assignment_1/latex/A1.tex](https://github.com/Hrithikraj2/MatrixTheory_EE5609/blob/master/Assignment_1/latex/A1.tex)