

Assignment-8

Hrithik Raj

Abstract—This document contains solution of Problem with Ramsey(4.1.4)

Download latex-tikz codes from

https://github.com/Hrithikraj2/MatrixTheory_EE5609/blob/master/Assignment_8/A8.tex

$$\mathbf{a}_j = \begin{pmatrix} a_{1j} \\ \vdots \\ a_{mj} \end{pmatrix} \in \mathfrak{R}^n \quad (2.0.9)$$

That gives us,

$$|C| = |\mathbf{c}_1, \dots, \mathbf{c}_m| \quad (2.0.10)$$

1 PROBLEM

If A is an $m \times n$ matrix, B is a $n \times m$ matrix and $n < m$, then Prove that AB is not invertible.

2 SOLUTION

Let us represent A and B as follows

$$A = (a_{ij}) \in \mathfrak{R}^{m \times n} \quad (2.0.1)$$

$$B = (b_{ij}) \in \mathfrak{R}^{n \times m} \quad (2.0.2)$$

Let C = AB which would be an $m \times m$ matrix.

$$C = (c_{ij}) \in \mathfrak{R}^{m \times m} \quad (2.0.3)$$

$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj} \quad (2.0.4)$$

for $j=1,2,\dots,m$. Let

$$\mathbf{c}_j = C_{ej} \in \mathfrak{R}^m \quad (2.0.5)$$

where (e_1, e_2, \dots, e_m) stands for the canonical basis of \mathfrak{R}^m . Then, we have

$$c_j = \begin{pmatrix} \sum_{k=1}^n a_{1k} b_{kj} \\ \vdots \\ \sum_{k=1}^n a_{mk} b_{kj} \end{pmatrix} \quad (2.0.6)$$

$$= \sum_{k=1}^n b_{kj} \begin{pmatrix} a_{1k} \\ \vdots \\ a_{mk} \end{pmatrix} \quad (2.0.7)$$

$$= \sum_{k=1}^n b_{kj} \mathbf{a}_k \quad (2.0.8)$$

$$= \sum_{k_1=1}^n \sum_{k_2=1}^n \dots \sum_{k_m=1}^n b_{k_1,1} b_{k_2,2} \dots b_{k_m,m} |\mathbf{a}_{k_1}, \dots, \mathbf{a}_{k_m}| \quad (2.0.11)$$

Since there are exactly n vectors \mathbf{a}_j and the determinant has $m > n$ entries, then we have

$$|\mathbf{a}_{k_1}, \dots, \mathbf{a}_{k_m}| = 0 \quad (2.0.12)$$

$$\forall k_1, \dots, k_m \in (1, 2, \dots, n) \quad (2.0.13)$$

Because atleast two entries must be equal. Hence

$$|C| = 0 \quad (2.0.14)$$

Since $|AB| = 0$. Therefore, AB is not invertible.