

Assignment-based Subjective Questions

1. **From your analysis of the categorical variables from the dataset, what could you infer about their effect on the dependent variable?**

Answer: The categorical variable in the dataset were a season, weathersit, holiday, mnth, yr and weekday. These were visualized using a boxplot. These variables had the following effect on our dependent variable:-

1. Season - The boxplot showed that spring season had least value of cnt whereas fall had maximum value of cnt. Summer and winter had intermediate value of cnt.

2. Weathersit - There are very less users when there is heavy rain/ snow indicating that this weather is extremely unfavorable. The highest count was seen when the weathersit was 'Clear, Partly Cloudy'.

3. Holiday - rentals reduced during the holiday.

4. Yr - The number of rentals in 2019 was more than 2018.


5. weekday – Monday and Friday saw the major number of users compared to other days.

2. **Why is it important to use drop_first=True during dummy variable creation?**

Answer: If we do not use drop_first = True, then n dummy variables will be created, and these predictors (n dummy variables) are themselves correlated which is known as multicollinearity.


3. **Looking at the pair-plot among the numerical variables, which one has the highest correlation with the target variable?**

Answer: "temp" and "atemp" are the two numerical variables which are highly correlated with the target variable (cnt)



4. How did you validate the assumptions of Linear Regression after building the model on the training set?

Answer:



Residuals distribution should follow normal distribution and centred around 0.(mean = 0).

We validate this assumption about residuals by plotting a distplot of residuals and see if residuals are following normal distribution or not. The above diagram shows that the residuals are distributed about mean = 0.

5. **Based on the final model, which are the top 3 features contributing significantly towards explaining the demand of the shared bikes?**

Answer:

The top 3 features are :-

1.temp - coefficient : 0.505

2.yr - coefficient : 0.239

3.season

General Subjective Questions


1. **Explain the linear regression algorithm in detail.**

Answer: Linear Regression is a type of supervised Machine Learning algorithm that is used for the prediction of numeric values. Linear Regression is the most basic form of regression analysis and the most commonly used predictive analysis model. Linear regression is based on the popular equation " $y = mx + c$ ".

It assumes that there is a linear relationship between the dependent variable(y) and the predictor(s)/independent variable(x). In regression, we calculate the best-fit line which describes the relationship between the independent and dependent variables. Regression is performed when the dependent variable is of continuous data type and Predictors or independent variables could be of any data type like continuous, nominal/categorical, etc. The regression method tries to find the best-fit line which shows the relationship between the dependent variable and predictors with the least error.

2. **Explain the Anscombe's quartet in detail.**

Answer: Anscombe's Quartet was developed by statistician Francis Anscombe. It includes four data sets that have almost identical statistical features, but they have a very different distribution and look totally different when plotted on a graph. It was developed to emphasize both the importance of graphing data before analyzing it and the effect of outliers and other influential observations on statistical properties.



3. What is Pearson's R?

Answer: Pearson's r is a numerical summary of the strength of the linear association between the variables. Its value ranges between -1 to +1. It shows the linear relationship between two sets of data. In simple terms, it tells us if we can draw a line graph to represent the data.

$r = 1$ means the data is perfectly linear with a positive slope.

$r = -1$ means the data is perfectly linear with a negative slope.

$r = 0$ means there is no linear association.

4. What is scaling? Why is scaling performed? What is the difference between normalized scaling and standardized scaling?

Answer: Feature scaling is a method used to normalize or standardize the range of independent variables or features of data. It is performed during the data preprocessing stage to deal with varying values in the dataset. If feature scaling is not done, then a machine learning algorithm tends to weigh greater values, higher and consider smaller values as the lower values, irrespective of the units of the values.

- Normalization is generally used when you know that the distribution of your data does not follow a Gaussian distribution. This can be useful in algorithms that do not assume any distribution of the data like K-Nearest Neighbors and Neural Networks.
- Standardization, on the other hand, can be helpful in cases where the data follows a Gaussian distribution. However, this does not have to be necessarily true. Also, unlike normalization, standardization does not have a bounding range. So, even if you have outliers in your data, they will not be affected by standardization.

5. You might have observed that sometimes the value of VIF is infinite. Why does this happen?

Answer: VIF - the variance inflation factor - The VIF gives how much the variance of the

coefficient estimate is being inflated by collinearity. $(VIF) = 1/(1-R_1^2)$. If there is a perfect correlation, then $VIF = \text{infinity}$. Where R_1^2 is the R-square value of that independent variable we want to check how well this independent variable is explained well by other independent variables- If that independent variable can be explained perfectly by other independent variables, then it will have perfect correlation and its R-squared value will be equal to 1. So, $VIF = 1/(1-1)$ which gives $VIF = 1/0$ which results in “infinity”.

6. What is a Q-Q plot? Explain the use and importance of a Q-Q plot in linear regression.

Answer: A q-q plot is a plot of the quantiles of the first data set against the quantiles of the second data set. It is used to compare the shapes of distributions. A Q-Q plot is a scatterplot created by plotting two sets of quantiles against one another. If both sets of quantiles came from the same distribution, we should see the points forming a line that's roughly straight.

The q-q plot is used to answer the following questions:

- Do two data sets come from populations with a common distribution?
- Do two data sets have a common location and scale?
- Do two data sets have similar distributional shapes?
- Do two data sets have similar tail behavior?