

Assignment 5

Hruday Beeravelli

Find Python Codes from below link

[https://github.com/Hruday-Beeravelli/
INTERNSHIP-IITH-1/blob/main/
Assignment4/A4.py](https://github.com/Hruday-Beeravelli/INTERNSHIP-IITH-1/blob/main/Assignment4/A4.py)

and Latex codes from below link

[https://github.com/Hruday-Beeravelli/
INTERNSHIP-IITH-1/blob/main/
Assignment4/A4.tex](https://github.com/Hruday-Beeravelli/INTERNSHIP-IITH-1/blob/main/Assignment4/A4.tex)

From (1.2.1)

Let

$$\mathbf{M} = \begin{pmatrix} \frac{9}{2} & \frac{15}{2} \\ -3 & -5 \end{pmatrix}^T \quad (1.2.6)$$

$$= \begin{pmatrix} \frac{9}{2} & -3 \\ \frac{15}{2} & -5 \end{pmatrix} \quad (1.2.7)$$

$$(1.2.8)$$

Row Reduction on \mathbf{M}

$$R_2 \leftarrow R_2 - \frac{5}{3} \times R_1$$

$$\mathbf{M} = \begin{pmatrix} \frac{9}{2} & -3 \\ 0 & 0 \end{pmatrix}$$

1 EXAMPLES 2

1.1 Question 11

Prove (by shewing that the area of the triangle formed by them is zero) that the following sets of three points are in a straight line $(-\frac{1}{2}, 3)$, $(-5, 6)$ and $(-8, 8)$

1.2 Solution

Rank of matrix method:

If rank of matrix is not full matrix after row reduction, then points are said to be collinear

$$((\mathbf{A} - \mathbf{B} \quad \mathbf{A} - \mathbf{C})^T) \quad (1.2.1)$$

$$\text{Let } \mathbf{A} = \begin{pmatrix} -\frac{1}{2} \\ 3 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} -5 \\ 6 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} -8 \\ 8 \end{pmatrix}$$

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} -\frac{1}{2} \\ 3 \end{pmatrix} - \begin{pmatrix} -5 \\ 6 \end{pmatrix} \quad (1.2.2)$$

$$= \begin{pmatrix} \frac{9}{2} \\ -3 \end{pmatrix} \quad (1.2.3)$$

$$\mathbf{A} - \mathbf{C} = \begin{pmatrix} -\frac{1}{2} \\ 3 \end{pmatrix} - \begin{pmatrix} -8 \\ 8 \end{pmatrix} \quad (1.2.4)$$

$$= \begin{pmatrix} \frac{15}{2} \\ -5 \end{pmatrix} \quad (1.2.5)$$

Since the rank of matrix \mathbf{M} is 1, which is not full rank, therefore the given points are collinear.




Fig. 0