References

- Aderhold, A., Husmeier, D., Lennon, J. J., Beale, C. M. & Smith, V. A. (2012). Hierarchical Bayesian models in ecology: reconstructing species interaction networks from non-homogeneous species abundance data. *Ecological Informatics*, 11, 55–64.
- Agarwal, D., Philips, J. M. & Venkatasubramanian, S. (2006). Hunting the bump: on maximizing statistical discrepancy. Proc. 17th Ann ACM-SIAM Symp. On Disc. Alg., pp. 1137–1146.
- Agnew, A. D. Q., Wilson, J. B. & Sykes, M. (1993). A vegetation switch as the cause of a forest/mire ecotone in New Zealand. *Journal of Vegetation Science*, 4, 273–278.
- Agresti, A. (1990). Categorical Data Analysis. New York: John Wiley & Sons, Inc.
- Aing, C., Halls, S., Oken, K., Dobrow, R., & Fieberg, J. (2011).
 A Bayesian hierarchical occupancy model for track surveys conducted in a series of linear, spatially correlated, sites.
 Journal of Applied Ecology, 48, 1508–1517.
- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, 19, 716–723.
- Albert, R. & Barabási, A.-L. (2002). Statistical mechanics of complex networks. Reviews of Modern Physics, 74, 47–97.
- Aldous, J. M. & Wilson, R. J. (2000). *Graphs and Applications:*An Introductory Approach. New York: Springer-Verlag.
- Allain, C. & Cloitre, M. (1991). Characterizing the lacunarity of random and deterministic fractal sets. *Physical Review A*, 44, 3552–3558.
- Allen, T. F. H. & Hoekstra, T. W. (1992). Toward a Unified Ecology. New York: University of Columbia Press.
- Almeida-Neto, M. & Guimaraes P. R. (2007). On nestedness analyses: rethinking matrix temperature and anti-nestedness. Oikos, 116, 716-722.
- Almeida-Neto, M., Guimaraes, P. & Guimaraes P. R. (2008).

 A consistent metric for nestedness analysis in ecological

- systems: reconciling concept and measurement. *Oikos*, **117**, 1227–1239.
- Alpargu, G. & Dutilleul, P. (2003). To be or not to be valid in testing the significance of the slope in simple quantitative linear models with autocorrelated errors. *Journal of Statis*tical Computation and Simulation, 73, 165–180.
- Alpargu, G. & Dutilleul, P. (2006). Stepwise regression in mixed quantitative linear models with autocorrelated errors. Communications in Statistics-Simulation and Computation, 35, 79-104.
- Amarasekare, P. (2009). Competition and coexistence in animal communities. In *The Princeton Guide to Ecology*. ed. S. A. Levin, pp. 196–201. Princeton: Princeton University Press.
- Andersen, M. (1992). Spatial analysis of two species interactions. Oecologia, 91, 134–140.
- Anderson, D. J. (1967). Studies on structure in plant communities. IV. Cyclical succession in *Dryas* communities from North-west Iceland. *Journal of Ecology*, **55**, 629–635.
- Anderson, M. J., Crist, T. O., Chase, J. M. et al. (2011). Navigating the multiple meanings of beta diversity: a roadmap for the practicing ecologist. Ecology Letters, 14, 19–28.
- Anderson, M. J., Ellingsen, K. E., & McArdle, B. H. (2006).Multivariate dispersion as a measure of beta diversity. *Ecology Letters*, 9, 683–693.
- Anselin, L. (1988) Spatial Econometrics: Methods and Models. New York: Kluwer Academic Publisher.
- Anselin, L. (1995). Local indicators of spatial association: LISA. Geographical Analysis, 27, 93–115.
- Anselin, L. (2001). Spatial econometrics. In: A Companion to Theoretical Econometrics, ed. B. Bultagi, pp. 310–330. Oxford: Basil Blackwell.
- Anselin, L. (2009). Spatial regression. In *The SAGE Handbook of Spatial Analysis*, eds. A. S. Fotheringham & P. A. Rogerson, pp. 255–275. London: SAGE Publications Ltd.
- Arbia, G., Benedetti, R. & Espa, G. (1996). Effects of the MAUP on image classification. *Geographical Systems*, 3, 123–141.
- Arnot, C., Fisher, P. F., Wadsworth, R. & Wellens, J. (2004).Landscape metrics with ecotones: pattern under uncertainty. *Landscape Ecology*, 19, 181–195.
- Atmar, W. & Patterson, B. D. (1993). The measure of order and disorder in the distribution of species in fragmented habitat. *Oecologia*, 96, 373–382.
- Aubréville, A. (1938). La forêt coloniale: Les forêts de l'Afrique occidentale française. Annals de l'Académie des Sciences Coloniales Paris, 9, 1–245.
- Aubréville, A. M. A. (1950-1951). La concept d'association dans la forêt dense equatoriale de la basse Côte

- d'Ivoire. Bulletin de la Société Botanique de France, 98, 145-158.
- Austin, M. P. (1987). Models for the analysis of species' response to environmental gradients. *Vegetatio*, **69**, 35–45.
- Austin, M. P. & Austin, B. O. (1980). Behaviour of experimental plant communities along a nutrient gradient. *Journal of Ecology*, 68, 891–918.
- Austin, M. P. & Smith, T. M. (1989). A new model for the continuum concept. Vegetatio, 83, 35-47.
- Austin, M. P., Nicholls, A. O., Doherty, M. D. & Meyers, J.A. (1994). Determining species response functions to an environmental gradient by means of a beta-function. *Journal of Vegetation Science*, 5, 215–228.
- Baddeley, A. J., Howard, C. V., Boyde, A. & Reid, S. (1987). Three-dimensional analysis of the spatial distribution of particles using the tandem-scanning reflected light microscope. *Acta Stereologica*, 6, 87–100.
- Bailey, T. C. & Gatrell, A. C. (1995). *Interactive Spatial Data Analysis*. Harlow: Longman Scientific & Technical.
- Baker, W. L. & Cai, Y. (1992). The r.le programs for multiscale analysis of landscape structure using the GRASS geographical information system. *Landscape Ecology*, 7, 291–302.
- Bang-Jensen, J. & Gutin G. (2002). Digraphs: Theory, Algorithms and Applications. London: Springer-Verlag.
- Barabási, A. (2009). Scale-free networks: a decade and beyond. *Science*, **325**, 412-413.
- Barabási, A., & Bonabeau, E. (2003). Scale-free networks. Scientific American, 288, 60-69.
- Barbier, N., Couteron, P., Lejoly, J., Deblauwe, V. & Lejeune, O. (2006). Self-organized vegetation patterning as a fingerprint of climate and human impact on semi-arid ecosystems. *Journal of Ecology*, **94**, 537–547.
- Barbujani, G. & Sokal, R. R. (1991). Zones of sharp genetic change in Europe are also linguistic boundaries. Proceedings of the National Academy of Sciences of the United States of America, 87, 1816–1819.
- Barbujani, G., Oden, N. N. & Sokal, R. R. (1989). Detecting regions of abrupt change in maps of biological variables. *Systematic Zoology*, 38, 376–389.
- Barot, S., Gignoux, J. & Menaut, J.-C. (1999). Demography of a savanna palm tree: predictions from comprehensive spatial pattern analysis. *Ecology*, 80, 1987–2005.
- Bartlett, M. S. (1935). Some aspects of the time correlation problem in regard to tests of significance. *Journal of the Royal Statistical Society*, 98, 536-543.
- Bartlett, M. S. (1948). Smoothing periodograms from time series with continuous spectra. *Nature*, 161, 686-687.

- Bascompte, J. (2009). Disentangling the web of life. *Science*, **325**, 416–419.
- Bascompte, J. & Solé, R. V. (1998). Spatiotemporal patterns in nature. Trends in Ecology and Evolution, 13, 173–174.
- Baselga, A. (2010) Partitioning the turnover and nestedness components of beta diversity. Global Ecology and Biogeography, 19, 134-143.
- Batschelet, E. (1981). Circular Statistics in Biology. London: Academic Press.
- Beale, C. M., Lennon, J. J., Yearsley, J. M, Brewer, M. J. & Elston, D. A. (2010). Regression analysis of spatial data. *Ecology Letters*, 13, 246–264.
- Beckage, B., Joseph, L., Belisle, P., Wolfson, D. B. & Platt, W. J. (2007). Bayesian change-point analyses in ecology. *New Phytologist*, 174, 456-467.
- Beguin, J., Martino, S., Rue, H. & Cumming, S. G. (2012). Hierarchical analysis of spatially autocorrelated data using integrated nested Laplace approximation. *Methods in Ecol*ogy and Evolution, 3, 921–929.
- Beiler, K. J., Durall, D. M., Simard, S. W., Maxwell, S. A. & Kretzer, A. M. (2009). Architecture of the wood-wide web: *Rhizopogon* spp. Genets link multiple Douglas-fir cohorts. *New Phytologist*, 185, 543–553.
- Beisner, B. E., Haydon, D. & Cuddington, K. L. (2003). Alternative stable states in ecology. Frontiers in Ecology and the Environment. 1, 376–382.
- Bekker, M. F. & Malanson, G. P. (2008). Linear forest patterns in subalpine environments. *Progress in Physical Geography*, 32, 635–653.
- Bell, G., Lechowicz, M. J., Appenzeller, A. et al. (1993). The spatial structure of the physical environment. *Oecologia*, 96, 114–121
- Bell, D. T., Plummer, J. A. & Taylor, S. K. (1993). Seed-germination ecology in southwestern Western-Australia. Botanical Review, 59, 24–73.
- Bellehumeur, C. & Legendre, P. (1997). Aggregation of sampling units: an analytical solution to predict variance. Geographical Analysis, 29, 258–266.
- Bellehumeur, C., Legendre, P. & Marcotte, D. (1997). Variance and spatial scales in a tropical rain forest: changing the size of sampling units. *Plant Ecology*, 130, 89–98.
- Benes, V. & Rataj, J. (2004). Stochastic Geometry: Selected Topics. Boston: Kluwer Academic Publisher.
- Bengtsson, J., Angelstam, P., Elmqvist, T. et al. (2003). Reserves, resilience and dynamic landscapes. Ambio, 32, 389–396.
- Bergman, C. M., Schaefer, J. A. & Luttich, S. N. (2000). Caribou movement as a correlated random walk. *Oecologia*, 123, 364–374.

- Bertness, M. D. & Calloway, R. (1994). Positive interactions in communities. *Trends in Ecology and Evolution*, 9, 191-193.
- Beyer H., Morales J., Murray D. & Fortin M-J. (2013). The effectiveness of Bayesian state-space models for estimating behavioural states from movement paths. *Methods in Ecol*ogy and Evolution, 4, 433–441.
- Bézier, P. (1977). Essai de définition numérique des courbes et des surfaces expérimentales. Thèse de doctorat ès-sciences, Université Pierre et Marie Curie, Paris.
- Bienert, A. Queck, R., Schmidt, A., Berhofer Ch., & Maas, H.-G. (2010). Voxel space analysis of terrestrial laser scans in forests for wind field modeling. *Int. Arch. Photogramm.*, *Rem. Sens. & Spat. Inf. Sci.*, 38, 92–97.
- Biggs, N. L., Lloyd, E. & Wilson, R. J. (1976). Graph Theory, 1736–1936. Oxford: Oxford University Press.
- Bini, L. M., Diniz, J. A. F., Rangel, T. F. L. V. et al. (2009). Coefficient shifts in geographical ecology: an empirical evaluation of spatial and non-spatial regression. *Ecography*, 32, 193–204.
- Bishop, Y. M. M., Fienberg, S. E. & Holland, P. W. (1975).
 Discrete Multivariate Analysis. Cambridge, MA: MIT Press.
- Bivand, R. (1980). A Monte Carlo study of correlation coefficient estimation with spatially autocorrelated observations. Quaestiones Geographicae, 6, 5–10.
- Bivand, R. S., Pebesma, E. J. & Gómez-Rubio, V. (2008).Applied Spatial Data Analysis with R. New York: Springer.
- Bjørnstad, O. N. & Falck, W. (2001). Nonparametric spatial covariance functions: estimation and testing. *Environmen*tal and Ecological Statistics, 8, 53–70.
- Bjørnstad, O. N., Ims, R. A. & Lambin, X. (1999a). Spatial population dynamics: analyzing patterns and processes of population synchrony. *Trends in Ecology and Evolution*, 14, 427–432.
- Bjørnstad, O. N., Stenseth, N. C. & Saitoh, T. (1999b). Synchrony and scaling in dynamics of voles and mice in northern Japan. *Ecology*, 80, 622–637.
- Blanchet, F. G., Legendre, P. & Borcard, D. (2008a). Modelling directional spatial processes in ecological data. *Ecological Modelling*, 215, 325–336.
- Blanchet, F. G., Legendre, P. & Borcard, D. (2008b). Forward selection of explanatory variables. *Ecology*, 89, 2623–2632.
- Blanchet, F. G., Legendre, P., Maranger, R., Monti, D. & Pepin, P. (2011). Modelling the effect of directional spatial ecological processes at different scales. *Oecologia*, 166, 357–368.
- Blonder, B. & Dornhaus, A. (2011). Time-ordered networks reveal limitations to information flow in ant colonies. *PLoS ONE*, 6(5), e20298.

- Bloomquist, E. W., Lemey, P. & Suchard, M. A. (2010). Three roads diverged? Routes to phylogeographic inference. *Trends in Ecology and Evolution*, **25**, 626–632.
- Blundell, G. M., Maier, J. A. K. & Debevec, E. M. (2001). Linear home ranges: effects of smoothing, sample size and autocorrelation on kernel estimates. *Ecological Monographs*, 71, 469–489.
- Bolker, B. M., Brooks, M. E., Clark, C. J. et al. (2009). Generalized linear mixed models: a practical guide for ecology and evolution. Trends in Ecology and Evolution, 24, 127-135.
- Bonabeau, E., Theraulaz, G., Deneubourg, J.-L., Aron, S. & Camazine, S. (1997) Self-organization in social insects. *Trends in Ecology and Evolution*, **12**, 188–193.
- Boots, B. N. (2002). Local measures of spatial association. Écoscience. 9. 168-176.
- Boots, B. N. (2003). Developing local measures of spatial association for categorical data. *Journal of Geographical* Systems, 5, 139–160.
- Borcard, D. & Legendre, P. (2002). All-scale spatial analysis of ecological data by means of principal coordinates of neighbour matrices. *Ecological Modelling*, 153, 51-68.
- Borcard, D. & Legendre P. (2012). Is the Mantel correlogram powerful enough to be useful in ecological analysis? A simulation study. *Ecology*, **93**, 1473–1481.
- Borcard, D., Legendre, P. & Drapeau, P. (1992). Partialling out the spatial component of ecological variation. *Ecology*, 73, 1045–1055
- Borcard, D., Gillet, F. & Legendre, P. (2011). *Numerical Ecology with R.* New York: Springer.
- Börger L., Franconi, N., Ferretti F. *et al.* (2006). An integrated approach to identify spatiotemporal and individual-level determinants of animal home range size. *The American Naturalist*, **168**, 471-485.
- Bossenbroek, J. M., Nekola, J. C. & Kraft, C. E. (2001). Prediction of long-distance dispersal using gravity models: zebra mussel invasion of inland lakes. *Ecological Applications*, 11, 1778–1788.
- Bourobou, D. N., Moussavou, H., Nzengue, E. et al. (2010). Applications for microsatellite markers in the management of forest trees. FAO Conference of Agricultural Biotechnology in Developing Countries, Guadalajara, 2010.
- Bowersox, M. A. & Brown, D. G. (2001) Measuring the abruptness of patchy ecotones: a simulation-based comparison of patch and edge metrics. *Plant Ecology*, **156**, 89–103.
- Box, G. & Jenkins, G. (1970). Time Series Analysis: Forecasting and Control. San Francisco: Holden-Day.
- Bradshaw, G. A. & Fortin, M.-J. (2000). Landscape heterogeneity effects on scaling and monitoring large areas using

- remote sensing data. Geographic Information Sciences, 6, 61-68
- Bradshaw, G. A. & Spies, T. (1992). Characterizing canopy gap structure in forests using wavelet analysis. *Journal of Ecol*ogy, 80, 205–215.
- Brandtberg, T. (1999). Automatic individual tree based analysis of high spatial resolution aerial images on naturally regenerated boreal forests. Canadian Journal of Forest Research, 29, 1464–1478.
- Bray, J. R. (1956). Gap phase replacement in a Maple-Basswood forest. *Ecology*, 37, 598-600.
- Brodie, C., Houle, G. & Fortin, M.-J. (1995). Development of a Populus balsamifera clone in subarctic Québec reconstructed from spatial analyses. Journal of Ecology, 83, 309–320.
- Brooker R. W., Maestre, F. T., Callaway, R. M. *et al.* (2008). Facilitation in plant communities: the past, the present, and the future. *Journal of Ecology*, **96**, 18–34.
- Brooks, C. P. (2006). Quantifying population substructure: extending the graph-theoretic approach. *Ecology*, 87, 864–872.
- Brown, D. G. (1998). Classification and boundary vagueness in mapping presettlement forest types. *International Journal* of Geographical Information Science, 12, 105–129.
- Brown, J. H. (1995). Macroecology. Chicago: University of Chicago Press.
- Brunt, J. W. & Conley, W. (1990). Behavior of a multivariate algorithm for ecological edge detection. *Ecological Modelling*, 49, 179–203.
- Burnham, K. P. & Anderson, D. R. (2002). Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach. Heidelberg and New York: Springer.
- Burrough, P. A. (1981). Fractal dimensions of landscapes and other environmental data. *Nature*, **294**, 240–242.
- Burrough, P. A. (1986). Principles of geographical information systems for land resources assessment. Monographs on Soil and Resources Survey No. 12. Oxford: Clarendon Press.
- Burrough, P. A. & Frank, A. (eds.) (1996). *Geographic Objects with Indeterminate Boundaries*. London: Taylor and Francis.
- Burrough, P. A. & McDonnell, R. A. (1998). Principles of Geographical Systems. Oxford: Oxford University Press.
- Cadenasso, M. L., Pickett, S. T. A., Weathers, K. C. et al. (2003).
 An interdisciplinary and synthetic approach to ecological boundaries. *Bioscience*, 53, 717–722.
- Cadman, M.D., Sutherland, D.A., Beck, G.G., Lepage, D. & Couturier, A.R. (eds.) (2007). Atlas of the Breeding Birds of Ontario, 2001–2005. Environment Canada, Ontario Field Ornithologists, Ontario Ministry of Natural Resources, and Ontario Nature, Toronto.

- Cain, M. L. (1989). The analysis of angular data in ecological field studies. *Ecology*, 70, 1540–1543.
- Cain, M. L. (1990). Models of clonal growth in Solidago altissima. Journal of Ecology, 78, 27-46.
- Cain, M. L. & Damman, H. (1997). Clonal growth and ramet performance in the woodland herb, Asarum canadense. Journal of Ecology, 85, 883–897.
- Cain, M. L., Pacala, S. W., Silander, J. A. & Fortin, M.-J. (1995).
 Neighborhood models of clonal growth in the white clover,
 Trifolium repens. American Naturalist, 145, 888–917.
- Campbell, J. E., Franklin, S. B., Gibson, D. J. & Newman, J. A. (1998). Permutation of two-term local quadrat variance analysis: general concepts for interpretation of peaks. *Jour-nal of Vegetation Science*, 9, 41–44.
- Candau, J.-N., Fleming, R. A. & Hopkin, A. (1998). Spatiotemporal patterns of large-scale defoliation caused by the spruce budworm in Ontario since 1941. *Canadian Journal* of Forest Research, 28, 1733–1741.
- Canny, J. (1986). A computational approach to edge detection. IEEE Transitional Pattern Analysis of Machine Intelligence, 8, 679–698.
- Cantwell, M. D. & Forman, T. T. (1993). Landscape graphs: ecological modeling with graph theory to detect configurations common to diverse landscapes. *Landscape Ecology*, 8, 239-255.
- Carl, G. & Kühn, I. (2008). Analyzing spatial ecological data using linear regression and wavelet analysis. Stochastic Environmental Research and Risk Assessment 22, 315–324.
- Carl, G. & Kühn, I. (2010). A wavelet-based extension of generalized linear models to remove the effect of spatial autocorrelation. Geographical Analysis, 42, 323–337.
- Carl, G., Dormann, C. F. & Kühn, I. (2008). A wavelet-based method to remove spatial autocorrelation in the analysis of species distributional data. Web Ecology, 8, 22–29.
- Carrer, M. & Urbinati, C. (2001). Spatial analysis of structural and tree-ring related parameters in a timberline forest in the Italian Alps. *Journal of Vegetation Science*, 12, 643–652.
- Casteigts, A., Flocchini, P., Quattrociocchi, W. & Santoro, N. (2011). Time-varying graphs and dynamic networks. 10th International Conference on *ad hoc* Networks and Wireless, July 2011.
- Cayuela, L., Benayas, J. M., Justel, A. & Salas-Rey, J. (2006). Modelling tree diversity in a highly fragmented tropical montane landscape. *Global Ecology and Biogeography*, 15, 602–613.
- Cerioli, A. (1997). Modified tests of independence in 2×2 tables with spatial data. *Biometrics*, **53**, 619-628.
- Cerioli, A. (2002). Testing mutual independence between two discrete-valued spatial processes: a correction to Pearson chi-squared. *Biometrics*, 58, 888–897.

- Chan, J., Bailey, J. & Leckie, C. (2008). Discovering correlated spatio-temporal changes in evolving graphs. *Knowledge* and Information Systems, 16, 53–96.
- Chao, A. (2004). Species richness estimation. In *Encyclopedia of Statistical Sciences*, eds. N. Balakrishnan, C. B. Read & A. Vidakovic. New York: Wiley.
- Chao, A., Chiu, C.-H. & Hsieh, T.C. (2012). Proposing a resolution to debates on diversity partitioning. *Ecology*, 93, 2037–2051.
- Chase, J. M., Kraft, N. B., Smith, K. G., Velland, M. & Inouye, B. D. (2011). Using null models to disentangle variation in community dissimilarity from variation in α-diversity. *Ecosphere*, 2, Article 24.
- Chatfield, C. (1975). The Analysis of Time Series: Theory and Practice. London: Chapman & Hall.
- Chilès, J.-P. & Delfiner, P. R. (2012). Geostatistics: Modeling Spatial Uncertainty, 2nd edn. New York: Wiley.
- Claramunt, C. & Thériault, M. (1997). Towards semantics for modeling spatio-temporal processes within GIS. In Advances in GIS Research II, eds. M. J. Kraak & M. Molenaar, pp. 47–63. London: Taylor & Francis.
- Clarke, K. R. & Warwick, R. M. (1998). A taxonomic distinctness index and its statistical properties. *Journal of Applied Ecology*, 35, 523–531.
- Claussen, D. L., Finkler, M. S. & Smith, M. M. (1997). Thread trailing of turtles: methods for evaluating spatial movements and pathway structure. *Canadian Journal of Zoology*, 75, 2120–2128.
- Clements, F. E. (1916). *Plant Succession*. Washington D.C.: Carnegie Institute.
- Cliff, A. D. & Ord, J. K. (1973). Spatial Autocorrelation. London: Pion.
- Cliff, A. D. & Ord, J. K. (1981). Spatial Processes: Models and Applications. London: Pion.
- Clifford, P., Richardson S. & Hémon, D. (1989). Assessing the significance of correlation between two spatial processes. *Biometrics*, 45, 123-134.
- Cochran, W. (1954). Some methods for strengthening the common χ^2 tests. *Biometrics*, **10**, 417–451.
- Codling, E. A, Plank, M. J. & Benhamou, S. (2008). Random walks models in biology. *Journal of the Royal Society Inter*face 5, 813–834.
- Cohn, R. D. (1999). Comparisons of multivariate relational structures in serially correlated data. *Journal of Agricultural*, *Biological & Environmental Statistics*, 4, 238–257.
- Collins, S. L., Glenn, S. M. & Roberts, D. W. (1993). The hierarchical continuum concept. *Journal of Vegetation Sci*ence, 4, 149–156.
- Colwell, R. K. (2009). Biodiversity: Concepts, patterns and measurement. *In The Princeton Guide to Ecology*, ed. S. A. Levin, pp. 257–263. Princeton: Princeton University Press.

- Condit, R., Ashton, P. S., Baker, P. et al. (2000). Spatial patterns in the distribution of tropical tree species. Science, 288, 1414–1417.
- Connell, J. H. & Slatyer, R. O. (1977). Mechanisms of succession in natural communities and their roles in community stability and organization. *American Naturalist*, 111, 1119–1144.
- Connor, E. F & Simberloff, D. (1984). Neutral models of species co-occurrence patterns. In *Ecological Communities:* Conceptual Issues And The Evidence, eds. D. R. Strong, Jr. et al., pp. 316–331. Princeton: Princeton University Press.
- Conover, W. J. (1980). Practical Nonparametric Statistics, 2nd edn. New York: Wiley.
- Couteron, P. & Ollier, S. S. (2005). A generalized, variogram based framework for multi-scale ordination. *Ecology*, 86, 828-834.
- Couwenberg, J., & Joosten, H. (2005). Self-organization in raised bog patterning: the origin of microtope zonation and mesotype diversity. *Journal of Ecology*, **93**, 1238–1248.
- Cox, D. R. (1955). Some statistical methods connected with series of events. *Journal of the Royal Statistical Society Series* B (Statistical Methodology), 17, 129–164.
- Craft, M. E., Volz, E., Packer, C. & Meyers, L. A. (2010). Disease transmission in territorial populations: the small-world network of Serengeti lions. *Journal of the Royal Society Interface*, doi: 10.1098/rsif.2010.0511.
- Crawley, M. J. (1986). Plant Ecology. Oxford: Blackwell.
- Cressie, N. A. C. (1991). Statistics for Spatial Data. New York: Wiley.
- Cressie, N. A. C. (1993). *Statistics for Spatial Data*, revised edn. New York: Wiley.
- Cressie, N. A. C. (1996). Change of support and the modifiable areal unit problem. *Geographical Systems*, 3, 159–180.
- Cressie, N. & Huang, H.-C. (1999). Classes of non-separable, spatio-temporal stationary covariance functions. *Journal of the American Statistical Association*, 94, 1330–1340.
- Cressie, N. & Wikle, C. K. (2011). Statistics for Spatio-Temporal Data. Hoboken NJ: Wiley.
- Cressie, N., Calder, C. A., Clark, J. S., Ver Hoef, J. M. & Wikle, C. K. (2009). Accounting for uncertainty in ecological analysis: the strengths and limitations of hierarchical statistical modeling. *Ecological Applications*, 19, 553–570.
- Crist, T. O., Veech, J. A., Gerring, J. C. & Summerville, K. S. (2003). Partitioning species diversity across landscapes and regions: a hierarchical analysis of alpha, beta, and gamma diversity. *American Naturalist*, **162**, 734–743.
- Csillag, F. & Kabos, S. (1996). Hierarchical decomposition of variance with applications in environmental mapping based on satellite images. *Mathematical Geology*, 28, 385–405.
- Csillag, F. & Kabos, S. (2002). Wavelets, boundaries, and the spatial analysis of landscape pattern. Écoscience, 9, 177-190.

- Csillag, F., Boots, B., Fortin, M.-J., Lowell, K. & Potvin, F. (2001). Multiscale characterization of boundaries and landscape ecological patterns. *Geomatica*, 55, 291–307.
- Currie, D. J. (2007). Disentangling the roles of environment and space in ecology. *Journal of Biogeography*, 34, 2009–2011.
- Cushman, S., McKelvey, K., Hayden, J. & Schwartz, M. (2006). Gene flow in complex landscapes: testing multiple hypotheses with causal modeling. *The American Naturalist*, **168**, 486–499.
- Cutler, N. A., Belyea, L. R. & Dugmore, A. J. (2008). The spatiotemporal dynamics of a primary succession. *Journal* of Ecology, 96, 231–246.
- Dale, M. R. T. (1977). Graph theoretical analysis of the phytosociological structure of plant communities: the theoretical basis. *Vegetatio*, 34, 137–154.
- Dale, M. R. T. (1979). The analysis of patterns of zonation and phytosociological structure of seaweed communities. Ph. D. Thesis. Halifax, N.S.: Dalhousie University.
- Dale, M. R. T. (1985). Graph theoretical methods for comparing phytosociological structures. Vegetatio, 63, 79–88.
- Dale, M. R. T. (1986). Overlap and spacing of species' ranges on an environmental gradient. Oikos, 47, 303–308.
- Dale, M. R. T. (1988). The spacing and intermingling of species boundaries on an environmental gradient. *Oikos*, 53, 351-356.
- Dale, M. R. T. (1995). Spatial pattern in communities of crustose saxicolous lichens. *Lichenologist*. 27, 495–503.
- Dale, M. R. T. (1999). Spatial Pattern Analysis in Plant Ecology. Cambridge: Cambridge University Press.
- Dale, M. R. T. (2000). Lacunarity analysis of spatial pattern: a comparison. *Landscape Ecology*, 15, 467-478.
- Dale, M. R. T. & Blundon, D. J. (1991). Quadrat covariance analysis and the scales of interspecific association during primary succession. *Journal of Vegetation Science*, 2, 103–112.
- Dale, M. R. T. & Fortin, M.-J. (2002). Spatial autocorrelation and statistical tests in ecology. *Écoscience*, **9**, 162–167.
- Dale, M. R. T. & Fortin, M.-J. (2009). Spatial autocorrelation and statistical tests: some solutions. *Journal of Agricultural*, *Biological, and Environmental Statistics*, 14, 188–206.
- Dale, M. R. T. & Fortin, M.-J. (2010). From graphs to spatial graphs. Annual Review of Ecology, Evolution, and Systematics, 41, 21–38.
- Dale, M. R. T. & MacIsaac, D. A. (1989). New methods for the analysis of spatial pattern in vegetation. *Journal of Ecology*, 77, 78–91.
- Dale, M. R. T. & Mah, M. (1998). The use of wavelets for spatial pattern analysis in ecology. *Journal of Vegetation Science*, 9, 805–814.
- Dale, M. R. T. & Powell, R. D. (1994). Scales of segregation and aggregation of plants of different kinds. *Canadian Journal of Botany*, 72, 448-453.

- Dale, M. R. T. & Powell, R. D. (2001). A new method for characterizing point patterns in plant ecology. *Journal of Vegetation Science*, 12, 597–608.
- Dale, M. R. T. & Zbigniewicz, M. W. (1995). The evaluation of multi-species pattern. *Journal of Vegetation Science*, 6, 391–398.
- Dale, M. R. T. & Zbigniewicz, M. W. (1997). Spatial pattern in boreal shrub communities: effects of a peak in herbivore density. *Canadian Journal of Botany*, 75, 1342–1348.
- Dale, M. R. T., Blundon, D. J., MacIsaac, D. A. & Thomas, A. G. (1991). Multiple species effects and spatial autocorrelation in detecting species associations. *Journal of Vegetation Science*, 2, 635–642.
- Dale, M. R. T., Dixon, P., Fortin, M.-J. et al. (2002). The conceptual and mathematical relationships among methods for spatial analysis. Ecography, 25, 558–577.
- Darling, D. A. (1953). On a class of problems related to the random division of an interval. Annals of Mathematical Statistics, 24, 239–253.
- Darlington, P. J. (1957). Zoogeography: The Geographical Distribution of Animals. New York: Wiley.
- Daubechies, I. (1992). *Ten Lectures on Wavelets*. Philadelphia: Society for Industrial and Applied Mathematics.
- Daubechies, I. (1993). Different Perspectives on Wavelets.
 Providence, RI: American Mathematical Society.
- Davidar, P., Yogananad, K., Ganesh, T. & Devy, S. (2002). Distributions of forest birds and butterflies in the Andaman islands, Bay of Bengal: nested patterns and processes. *Eco-graphy*, 25, 5–16.
- de Jong, P., Aarssen, L. W. & Turkington, R. (1980). The analysis of contact sampling data. *Oecologia*, 45, 322–324.
- de Solla, S. R., Bonduriansky, R. & Brooks, R. J. (1999). Eliminating autocorrelation reduces biological relevance of home range estimates. *Journal of Animal Ecology*, 68, 221–234.
- Deblauwe, V., Couteron, P., Lejeune, O., Bogaert, J. & Barbier, N. (2011). Environmental modulation of self-organized periodic vegetation patterns in Sudan. *Ecography*, 34, 990– 1001.
- Del Mondo, G., Stell, J.G., Claramunt, C. & Thibaud, R. (2010).
 A graph model for spatio-temporal evolution. *Journal of Universal Computer Science*, 16, 1452–1477.
- Delmelle, E. & Goovaerts, P. (2009). Second-phase sampling designs for non-stationary spatial variables. *Geoderma*, 153, 205–216.
- Demattei, C. & Cucala, L. (2011). Multiple spatio-temporal cluster detection for case event data: an ordering-based approach. Communications in Statistics - Theory & Methods, 40, 358-372.

- Dennis, B., Desharnais, R. A., Cushing, J. M., Henson, S. M. & Costantino, R. F. (2001). Estimating chaos and complex dynamics in an insect population. *Ecological Monographs*, 71, 277–303.
- Deutsch, C. V. & Journel, A. G. (1992). GSLIB. Geostatistical Software Library and User's Guide. New York: Oxford University Press.
- Dieckmann, U., Law, R. & Metz, J. A. J. (eds.) (2000). The Geometry of Ecological Interactions. Cambridge: Cambridge University Press.
- Dietz, E. J. (1983). Permutation tests for association between two distance matrices. Systematic Zoology, 32, 21–26.
- Diggle, P. J. (1979). Statistical methods for spatial point patterns in ecology. In *Spatial and Temporal Analysis in Ecol*ogy, eds. R. M. Cormack & J. K. Ord, pp. 95–150. Fairland, MD: International Cooperative Publishing House.
- Diggle, P. J. (1983). Statistical Analysis of Spatial Point Patterns. London: Academic Press.
- Diggle, P. J. (2003) Statistical Analysis of Spatial Point Patterns, 2nd edition. Hodder Arnold.
- Diggle, P. J. & Chetwynd A. G. (1991). Second-order analysis of spatial clustering for inhomogeneous populations. *Biometrics*, 47, 1155–1163.
- Diggle, P. J. & Justiniano, P. (2010). Model-based Geostatistics. New York: Springer.
- Diggle, P. J. & Ribeiro, P. J. (2007). *Model-Based Geostatistics*. Springer, New York, USA.
- Diggle P. J., Chetwynd, A. G., Haggkvist, R. & Morris, S. (1995).Second-order analysis of space-time clustering. Statististical Methods in Medical Research, 4, 124-136.
- Diks, C., Takens, F. & DeGoede, J. (1997). Spatio-temporal chaos: a solvable model. *Physica D*, **104**, 269–285.
- Diniz-Filho, J. A. F. & Bini, L. M. (2005). Modelling geographical patterns in species richness using eigenvector-based spatial filters. Global Ecology and Biogeography, 14, 177-185.
- Diniz-Filho, J. A. F., Bini, L. M. & Hawkins, B. A. (2003). Spatial autocorrelation and red herrings in geographical ecology. *Global Ecology and Biogeography*, 12, 53–64.
- Diniz-Filho, J. A. F., Nabout, J. C., Telles, M. P. D., Soares, T. N. & Rangel, T. (2009). A review of techniques for spatial modeling in geographical, conservation and landscape genetics. *Genetics and Molecular Biology*, 32, 203–211.
- Dixon, P. M. (2002). Nearest-neighbor contingency table analysis of spatial segregation for several species. *Écoscience*, **9**, 142–151.
- Doak, P. (2000). Population consequences of restricted dispersal for an insect herbivore in a subdivided habitat. *Ecol*ogy, 81, 1828–1841.

- Doebeli, M. & Ruxton, G. D. (1998). Stabilization through spatial pattern formation in metapopulations with longrange dispersal. *Proceedings of the Royal Society of London* Series B, 265, 1325–1332.
- Dong, P. (2009). Lacunarity analysis of raster datas ets and 1D, 2D, and 3D point patterns. *Computers & Geosciences*, 35, 2100-2110.
- Doob, J. L. (1996). The development of rigor in mathematical probability (1900–1950). The American Mathematical Monthly, 103, 586–595.
- Dormann, C. F. (2006). Effects of incorporating spatial autocorrelation into the analysis of species distribution data. Global Ecology and Biogeography, 16, 129-138.
- Dormann, C. F. (2009). Response to Comment on "Methods to account for spatial autocorrelation in the analysis of species distributional data: a review". *Ecography*, **32**, 379–381.
- Dormann, C. F., McPherson, J. M., Araújo, M. B. et al. (2007). Methods to account for spatial autocorrelation in the analysis of species distributional data: a review. Ecography, 30, 609–628.
- Dos Santos, D. A., Fernandez, H. R., Cuezzo, M. G. & Dominguez, E. (2008). Sympatry inference and network analysis in biogeography. Systematic Biology, 57, 432-448.
- Doty, M. S. & Archer, J. G. (1950). An experimental test of the tide factor hypothesis. *American Journal of Botany*, 37, 458–464.
- Doupanloup, I., Schneider, S. & Excoffier, L. (2002). A simulated annealing approach to define the genetic structure of populations. *Molecular Ecology*, 11, 2571–2581.
- Drake D. A. R. & Mandrak N. E. (2010). Least-cost transportation networks predict spatial interaction of invasion vectors. *Ecological Applications*, 20, 2286–2299.
- Dray, S. (2011). A new perspecive about Moran's *I* coefficient: spatial autocorrelation as a linear regression problem. *Geo-graphical Analysis*, 43, 127–141.
- Dray, S., Legendre P. & Peres-Neto P. R. (2006). Spatial modeling: a comprehensive framework for principal coordinate analysis of neighbor matrices (PCNM). *Ecological Modelling*, 196, 483–493.
- Dray, S., Pélissier, R., Couteron, P. et al. (2012). Community ecology in the age of multivariate multiscale spatial analysis. Ecological Monographs, 82, 257-275.
- Duczmal, L., Kulldorff, M. & Huang, L. (2006). Evaluation of spatial scan statistics for irregularly shaped clusters. *Journal* of Computational and Graphical Statistics, 15, 428–442.
- Dufresne, F., Bourget, E. & Bernatchez, L. (2002). Differential patterns of spatial divergence in microsatellite and allozyme alleles: further evidence for locus-specific selection in the acorn barnacle, Semibalanus balanoides? Molecular Ecology, 11, 113–123.

- Dungan, J. L. (2001). Scaling up and scaling down: relevance of the support effect on remote sensing of vegetation. In *Modelling Scale in Geographical Information Science*, eds.
 N. J. Tate & P. M. Atkinson, pp. 231–235. New York: Wiley.
- Dungan, J. L., Perry, J. N., Dale, M. R. T. et al. (2002). A balanced view of scale in spatial statistical analysis. *Ecography*, 25, 626–640.
- Dutilleul, P. (1993a). Spatial heterogeneity and the design of ecological field experiments. *Ecology*, 74, 1646–1658.
- Dutilleul, P. (1993b). Modifying the t test for assessing the correlation between two spatial processes. *Biometrics*, 49, 305–314.
- Dutilleul, P. R. L. (2011). Spatio-temporal Heterogeneity: Concepts and Analyses. Cambridge: Cambridge University Press.
- Dutilleul, P., Stockwell, J. D., Frigon, D. & Legendre, P. (2000).
 The Mantel test versus Pearson's correlation analysis: assessment of the differences for biological and environmental studies. *Journal of Agricultural, Biological and Environmental Statistics*, 5, 131–150.
- Dyer, R. J. & Nason, J. D. (2004). Population graphs: the graph theoretic shape of genetic structure. *Molecular Ecology*, 13, 1713–1727.
- Eddy, S. R. (2004). What is Bayesian statistics? *Nature Biotechnology*, 22, 1177–1178.
- Edgington, E. S. (1995). *Randomization Tests*, 3rd edn. New York: Marcel Dekker.
- Edwards, G. & Fortin, M.-J. (2001). Cognitive view of spatial uncertainty. In *Spatial Uncertainty in Ecology: Implications for Remote Sensing and GIS Applications*, eds. C. Hunsaker, M. Goodchild, M. Friedl & T. Case, pp. 133–157. New York: Springer-Verlag.
- Edwards, G. & Lowell, K. E. (1996). Modeling uncertainty in photointerpreted boundaries. *Photogrammetric Engineering* and Remote Sensing, 62, 337–391.
- Efron, B. & Tibshirani, R. J. (1993). An Introduction to the Bootstrap. New York: Chapman & Hall.
- Egbert, G. D. & Lettenmaier, D. P. (1986). Stochastic modeling of space-time structure of atmospheric chemical deposition. Water Resources Research, 22, 165-179.
- Ellner, S. & Turchin, P. (1995). Chaos in a noisy world: new methods and evidence from time-series analysis. *American Naturalist*, 145, 343–375.
- Epperson, B. K. (2003). Covariances among join-count spatial autocorrelation measures. *Theoretical Population Biology*, 64, 81-87.
- Erdös, P. & Rényi, A. (1960). On the evolution of random graphs. Publications of the Mathematical Institute of the Hungarian Academy of Science, 5, 17-61.

- Escudero, A., Iriondo, J. M. & Torres, M. E. (2003). Spatial analysis of genetic diversity as a tool for plant conservation. *Biological Conservation*, 113, 351–365.
- Evans, J. P. & Cain, M. L. (1995). A spatially explicit test of foraging behaviour in a clonal plant. *Ecology*, 76, 1147–1155.
- Everhart, S. E., Askew, A., Seymour, L., Holb, I. J. & Scherm, H. (2011). Characterization of three-dimensional spatial aggregation and association patterns of brown rot symptoms within intensively mapped sour cherry trees. *Annals of Botany*, 108, 1195–1202.
- Excoffier, L., Smouse, P. & Quattro, J. (1992). Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics*, 131, 479–491.
- Fagan, W. F. (2002). Connectivity, fragmentation, and extinction risk in dendritic metapopulations. *Ecology*, 83, 3243–3249.
- Fagan, W. F., Fortin, M.-J. & Soykan, C. (2003). Integrating edge detection and dynamic modeling in quantitative analyses of ecological boundaries. *BioScience*, 53, 730-738.
- Faghih, F. & Smith, M. (2002). Combining spatial and scale-space techniques for edge detection to provide a spatially adaptive wavelet-based noise filtering algorithm. *IEEE Transactions on Image Processing*, 11, 1069–1071.
- Fall, A., Fortin, M.-J., Manseau, M. & O'Brien, D. (2007). Spatial graphs: principles and applications for habitat connectivity. *Ecosystems*, 10, 448–461.
- Fattorini, S. (2002). Biogeography of the tenebrionid beetles (Coleoptera, Tenebrionidae) on the Aegean Islands (Greece). *Journal of Biogeography*, **29**, 49–67.
- Fattorini, S. (2007). Non-randomness in the species-area relationship: testing the underlying mechanisms. *Oikos*, 116, 678-689
- Feeley, K. J., Gillespie, T. W., Lebbin, D. J. & Walter, H. S. (2007). Species characteristics associated with extinction vulnerability and nestedness rankings of birds in tropical forest fragments. *Animal Conservation*, 10, 493–501.
- Felsenstein, J. (1975). A pain in the torus: some difficulties with models of isolation by distance. *The American Naturalist*, 109, 359–368.
- Fischer, J. & Lindenmayer, D. B. (2007). Landscape modification and habitat fragmentation: a synthesis. *Global Ecology* and Biogeography, 16, 265–280.
- Fisher, R. A. (1932). Statistical Methods for Research Workers, 4th edn. London: Oliver & Boyd.
- Fisher, R. A. (1935). The mathematical distributions used in the common tests of significance. *Econometrica*, 3, 353–365.
- Fisher, R. A. (1970). Statistical Methods for Research Workers. Edinburgh: Oliver & Boyd.

- Fitzpatrick, M. C., Preisser, E. L., Porter, A. et al. (2010). Ecological boundary detection using Bayesian areal wombling. Ecology, 91, 3448–3455.
- Foody, G. M. (2004). Spatial nonstationarity and scaledependency in the relationship between species richness and environmental determinants for the sub-Saharan endemic avifauna. Global Ecology and Biogeography, 13, 315-320.
- Fortin, M.-J. (1992). Detection of ecotones: definition and scaling factors. Ph.D. thesis, Department of Ecology and Evolution, State University of New York at Stony Brook, 258 pages.
- Fortin, M.-J. (1994). Edge detection algorithms for twodimensional ecological data. *Ecology*, 75, 956–965.
- Fortin, M.-J. (1997). Effects of data types on vegetation boundary delineation. Canadian Journal of Forest Research, 27, 1851–1858.
- Fortin, M.-J. (1999a). Effects of sampling unit resolution on the estimation of the spatial autocorrelation. Écoscience, 6, 636-641
- Fortin, M.-J. (1999b). The effects of quadrat size and data measurement on the detection of boundaries. *Journal of Vegetation Science*, 10, 43–50.
- Fortin, M.-J. & Dale, M. R. T. (2005). Spatial Analysis: A Guide for Ecologists. Cambridge: Cambridge University Press.
- Fortin, M.-J. & Dale, M. R. T. (2009). Spatial autocorrelation in ecological studies: a legacy of solutions and myths. *Geo-graphical Analysis*, 41, 392–397.
- Fortin, M.-J. & Drapeau, P. (1995). Delineation of ecological boundaries: comparisons of approaches and significance tests. Oikos, 72, 323–332.
- Fortin, M.-J. & Gurevitch, J. (2001). Mantel tests: spatial structure in field experiments. In *Design and Analysis of Ecological Experiments*, 2nd edn, eds. S. M. Scheiner & J. Gurevitch, pp. 308–326. Oxford: Oxford University Press.
- Fortin, M.-J. & Jacquez, G. M. (2000). Randomization tests and spatially autocorrelated data. *Bulletin of the Ecological Soci*ety of America, 81, 201–205.
- Fortin, M.-J. & Melles, S. J. (2009). Avian spatial responses to forest spatial heterogeneity at the landscape level: conceptual and statistical challenges. In *Real World Ecology: Large-Scale and Long-Term Case Studies and Method*, eds. S. Miao, S. Carstenn & M. Nungesser, pp. 137–160. New York: Springer.
- Fortin, M.-J. & Payette, S. (2002). How to test the significance of the relation between spatially autocorrelated data at the landscape scale: a case study using fire and forest maps. *Écoscience*, **9**, 213–218.

- Fortin, M.-J., Boots, B., Csillag, F. & Remmel, T. K. (2003). On the role of spatial stochastic models in understanding landscape indices in ecology. *Oikos*, 102, 203–212.
- Fortin, M.-J., Drapeau, P. & Jacquez, G. M. (1996). Quantification of the spatial co-occurrences of ecological boundaries. *Oikos*, 77, 51–60.
- Fortin, M.-J., Drapeau, P. & Legendre, P. (1989). Spatial autocorrelation and sampling design in plant ecology. *Vegetatio*, 83, 209–222.
- Fortin, M.-J., Jacquez, G. M. & Shipley, B. (2002). Computerintense methods. In *Encyclopedia of Environmetrics*, eds. A. El-Shaarawi & W. W. Pesgorsch, pp. 399–402. Chishester: Wiley.
- Fortin, M.-J., Jacquez, G. M. & Shipley, B. (2012b). Computerintense methods. In *Encyclopedia of Environmetrics*, 2nd edition, eds. A. H. El-Shaarawi & W.W. Piegorsch, pp. 489-493. Chichester: Wiley.
- Fortin M.-J., James, P. M. A., MacKenzie, A., Melles, S.J. &. Rayfield, B. (2012a). Spatial statistics, spatial regression, and graph theory in ecology. *Spatial Statistics*, 1, 100–109.
- Fortin, M.-J., Keitt, T. H., Maurer, B. A., Taper, M. L., Kaufman, D. M. & Blackburn, T. M. (2005). Species ranges and distributional limits: pattern analysis and statistical issues. *Oikos.* 108, 7–17.
- Fortin, M.-J., Marineau, K. & Payette, S. (1999). Spatial vegetation diversity index along a postfire successional gradient in the northern boreal forest. *Écoscience*, 6, 204–213.
- Fortin, M.-J., Olson, R. J., Ferson, S. et al. (2000). Issues related to the detection of boundaries. Landscape Ecology, 15, 453– 466
- Fortuna, M. A. & Bascompte, J. (2008). The network approach in ecology. In *Unity in Diversity: Reflections on Ecology after* the Legacy of Ramon Margalef, eds. F. Valladares, A. Camacho, A. Elosegi, M. Estrada, J. M. Gili, C. Gracia & J. C. Senar, pp. 371–393. Madrid: Fundación BBVA.
- Fortuna, M. A., Garcia, C., Guimarães, P. R. & Bascompte, J. (2008). Spatial mating networks in insect-pollinated plants. *Ecology Letters*, 11, 490-498.
- Fortuna, M. A., Gómez-Rodriguez, C. & Bascompte, J. (2006). Spatial network structure and amphibian persistence in stochastic environments. *Proceedings of the Royal Society of London Series B, Biological Sciences*, 273, 1429–1434.
- Fotheringham, A. S. (2009). Geographically weighted regression. In *The SAGE Handbook of Spatial Analysis*, eds. A. S. Fotheringham & P. A. Rogerson, pp. 243–253. London: SAGE Publications Ltd.
- Fotheringham, A. S. & Rogerson, P. A. eds. (2009). *The SAGE Handbook of Spatial Analysis*. Sage.

- Fotheringham, A. S. & Zhan, F. B. (1996). A comparison of three exploratory methods for cluster detection in spatial point patterns. *Geographical Analysis*, 28, 200–218.
- Fotheringham, A. S., Brunsdon, C. & Charlton, M. (2000). *Quantitative Geography: Perspectives on Spatial Data Analysis.* London: Sage Publications.
- Fotheringham, A. S., Brunsdon, C. & Charlton, M. (2002).

 Geographically Weighted Regression: The Analysis of
 Spatially Varying Relationships. Chichester: Wiley.
- Fowler, H. W. & Fowler, F. G. (1976). The Concise Oxford Dictionary of Current English, 6th edn, ed. J. B. Sykes. Oxford: Oxford University Press.
- Foxall, R. & Baddeley, A. (2002). Nonparametric measures of association between a spatial point process and a random set, with geological applications. *Applied Statistics*, 51, 165– 182.
- Franco, M. & Harper, J. (1988). Competition and the formation of spatial pattern in spacing gradients: an example using Kochia scoparia. Journal of Ecology, 76, 959-974.
- Frelich, L. (2002). Forest Dynamics and Disturbance Regimes: Studies from Temperate Evergreen-Deciduous Forests. New York: Cambridge University Press.
- Friedman, M. (1937). The use of ranks to avoid the assumption of normality implicit in the analysis of variance. *Journal of the American Statistical Association*, **32**, 675–701.
- Fukushima, Y., Hiura, T. & Tanabe, S. I. (1998). Accuracy of the MacArthur-Horn method for estimating a foliage profile. Agricultural and Forest Methodology, 92, 203–210.
- Gabriel, K. R. & Sokal, R. R. (1969). A new statistical approach to geographic variation analysis. *Systematic Ecology*, 18, 259–270.
- Galiano, E. F. (1982). Pattern detection in plant populations through the analysis of plant-to-all-plants distances. *Vegetatio*, 49, 39-43.
- Galiano, E. F. (1983). Detection of multi-species patterns in plant populations. *Vegetatio*, 53, 129–138.
- García Molinos, J. & Donohue, I. (2011). Temporal variability within disturbance events regulates their effects on natural communities. *Oecologia*, 166, 795–806.
- Garroway, C. J., Bowman, J., Carr, D. & Wilson, P. J. (2008).Applications of graph theory to landscape genetics. *Evolutionary Applications*, 1, 620–630.
- Gaston, K. J. & Blackburn, T. M. (2000). Patterns and Processes in Macroecology. Malden: Blackwell Publishing Company.
- Gauch, H. G. & Whittaker, R. H. (1972). Coenocline simulation. *Ecology*, 53, 446-451

- Gaucherel, C., Burel, F. & Baudry, J. (2007). Multiscale and surface pattern analysis of the effect of landscape pattern on carabid beetles distribution. *Ecological Indicators*, 7, 598– 609.
- Gauthier, O. & Lapointe, F.-J. (2007). Seeing the trees for the network: consensus, information content, and superphylogenies. Systematic Biology, 56, 345–355.
- Gavrikov, V. & Stoyan, D. (1995). The use of marked point processes in ecological and environmental forest studies. *Environmental and Ecological Statistics*, 2, 331–344.
- Geary, R. C. (1954). The contiguity ratio and statistical mapping. The Incorporated Statistician, 5, 115–145.
- Gelfand, A. E. (2012). Hierarchical modeling for spatial data problems. Spatial Statistics, 1, 30–39.
- Gellert, W., Küstner, H., Hellwich, M. & Kästner, H. (1977). The VNR Concise Encyclopedia of Mathematics. New York: Van Nostrand Reinhold.
- Gering, J. C., Crist, T. O. & Veech, J. A. (2003). Additive partitioning of species diversity across multiple spatial scales: Implications for regional conservation of biodiversity. *Conservation Biology*, 17, 488–499.
- Gerrard, D. J. (1969). Competition quotient: a new measure of the competition affecting individual forest trees. *Research Bulletin 20*. Agricultural Experiment Station, Michigan State University.
- Getis, A. (1991). Spatial interaction and spatial autocorrelation: a cross product approach. *Environment and Planning A*, 23, 1269–1277.
- Getis, A. & Boots, B. (1978). Models of Spatial Processes: An Approach to the Study of Point, Line and Area Patterns. Cambridge: Cambridge University Press.
- Getis, A. & Franklin, J. (1987). Second-order neighborhood analysis of mapped point patterns. *Ecology*, 68, 473–477.
- Getis A. & Ord, J. K. (1992). The analysis of spatial association by use of distance statistics. *Geographical Analysis*, 24, 189– 206.
- Getis, A. & Ord, J. K. (1996). Local spatial statistics: an overview. In *Spatial Analysis: Modelling in a GIS Environment*, eds. P. Longley & M. Batty, pp. 261–277. Cambridge: GeoInformation International.
- Gignoux, J., Camille, D. & Sebastien, B. (1999). Comparing the performances of Diggle's tests of spatial randomness for small samples with and without edge-effect correction: application to ecological data. *Biometrics*, 55, 156-164.
- Gilbert, B. & Bennett, J. R. (2010). Partitioning variation in ecological communities: do the numbers add up? *Journal of Applied Ecology*, 47, 1071–1082.
- Gill, J. (2002). Bayesian Methods: A Social and Behavioral Sciences Approach. Boca Raton: Chapman & Hall/CRC.

- Glaz, J., Naus, J. & Wallenstein, S. (2001). Scan Statistics. New York: Springer-Verlag.
- Gleason, H. A. (1927). Further views of the succession concept. *Ecology* 8, 299–326.
- Gleik, J. (1987). Chaos: Making a New Science. New York: Viking.
- Glenn-Lewin, D. C. & van der Maarel, E. (1992). Patterns and processes of vegetation dynamics. In *Plant Succession: Theory and Prediction*, eds. D. C. Glenn-Lewin, R. K. Peet & T. T. Veblen, pp. 11–59. London: Chapman & Hall.
- Glenn-Lewin, D. C., Peet, R. K. & Veblen, T. T. eds. (1992).
 Plant Succession: Theory and Prediction. London: Chapman & Hall.
- Gonzalez-Andujar, J. L. & Perry, J. N. (1993). Chaos, metapopulations and dispersal. *Ecological Modelling*, 65, 255–263.
- Good, P. (1993). Permutation Tests: A Practical Guide to Resampling Methods for Hypothesis Testing. New York: Springer-Verlag.
- Goovaerts, P. (1997). Geostatistics for Natural Resources Evaluation. New York: Oxford University Press.
- Goovaerts, P. & Jacquez, G. M. (2004). Accounting for regional background and population size in the detection of spatial clusters and outliers using geostatistical filtering and spatial neutral models: the case of lung cancer in Long Island, New York. *International Journal of Health Geographics*, 3, 14.
- Goovaerts, P. & Jacquez, G. M. (2005). Detection of temporal changes in the spatial distribution of cancer rates using local Moran's *I* and geostatistically simulated spatial neutral models. *Journal of Geographical Systems*, 7, 137–159.
- Gordon, A. D. (1999). Classification. *Monographs on Statistics* and Applied Probability 82, 2nd edn. London: Chapman & Hall/CRC
- Goreaud, F. & Pélissier, R. (1999). On explicit formulas of edge effect correction for Ripley's K-function. *Journal of Vegeta*tion Science, 10, 433–438.
- Gosz, J. R. (1993). Ecotone hierarchies. Ecological Applications, 3, 368-376.
- Gotelli, N. J. (2004). A taxonomic wish-list for community ecology. Philosophical Transactions of the Royal Society of London Series B-Biological Sciences, 359, 585–597.
- Goulard, M., Pagès, L. & Cabanettes, A. (1995). Marked point process: using correlation functions to explore a spatial data set. *Biometrical Journal*, 37, 837–853.
- Grant, E. H.C., Lowe, W. H. & Fagan, W. F. (2007). Living in the branches: population dynamics and ecological processes in dendritic networks. *Ecology Letters*, 10, 165–175.
- Grant, P. R., & Schluter. D. (1984). Interspecific competition inferred from patterns of guild structure. In *Ecological Com*munities: Conceptual Issues and the Evidence, eds. D. R.

- Strong, Jr. *et al.*, pp. 201–233. Princeton: Princeton University Press.
- Greig-Smith, P. (1961). Data on pattern within plant communities. I. The analysis of pattern. *Journal of Ecology*, 49, 695–702.
- Greig-Smith, P. (1983). Quantitative Plant Ecology, 3rd edn. Berkeley: University of California Press.
- Griffith, D. A. (1981). Interdependence in space and time: numerical and interpretative considerations. In *Dynamic Spatial Models*, eds. D. A. Griffith & R. D. MacKinnon, pp. 258–287. Netherlands: Sijthoff & Noordhoff.
- Griffith, D. A. (1992). What is spatial autocorrelation? Reflection on the past 25 years of spatial statistics. L'Espace géographique, 3, 265–280.
- Griffith, D. A. (1996). Spatial autocorrelation and eigenfunctions of the geographic weights matrix accompanying georeferenced data. *Canadian Geographer*, 40, 351–367.
- Griffith, D. A. (2008). Spatial-filtering-based contributions to a critique of geographically weighted regression (GWR). Environment and Planning A, 40, 2751-2769.
- Griffith, D. A. & Paelinck, J. H. P. (2009). Specifying a joint space- and time-lag using a bivariate Poisson distribution. *Journal of Geographical Systems*, 11, 23–36.
- Griffith, D. A. & Peres-Neto P. R. (2006). Spatial modeling in ecology: the flexibility of eigenfunction spatial analyses. *Ecology*, 87, 2603–2613.
- Grillet, M.-E., Barrera, R., Martínez, J.-E., Berti, J. & Fortin, M.-J. (2010a). Disentangling the effect of local and global spatial variation on a mosquito-borne infection in a neotropical heterogeneous environment. *American Journal of Tropical Medicine and Hygiene*, 82, 194–201.
- Grillet, M.-E., Jordan, G. J. & Fortin, M.-J. (2010b). State transition detection in the spatio-temporal incidence of malaria. Spatial and Spatio-Temporal Epidemiology Journal, 1, 251–259.
- Gross, T. & Blasius, B. (2008). Adaptive coevolutionary networks: a review. *Journal of the Royal Society, Interface* 5, 259, 271
- Guillot, G. & Rousset, F. (2013). Dismantling the Mantel tests.

 Methods in Ecology and Evolution, 4, 336–344.
- Guillot, G., Leblois, R., Coulon, A. & Frantz, A. C. (2009). Statistical methods in spatial genetics. *Molecular Ecology*, 18, 4734–4756.
- Guo, Q. & Kelly, M. (2004). Interpretation of scale in paired quadrat variance methods. *Journal of Vegetation Science*, 15, 763–770.
- Gustafson, E. J. (1998). Quantifying landscape spatial pattern: what is the state of the art? *Ecosystems*, 1, 143–156.
- Haas, S. E., Hooten, M. B, Rizzo, D. M. & Meentemeyer, R. K. (2011). Forest species diversity reduces disease risk in a

- generalist plant pathogen invasion. *Ecology Letters*, **14**, 1108–1116.
- Haase, P. (1995). Spatial pattern analysis in ecology based on Ripley's K-function: introduction and methods of edge correction. *Journal of Vegetation Science*, 6, 575–582.
- Haining, R. P. (1978). The moving average model for spatial interaction. Transactions of the Institute for British Geographers, NS3, 202–225.
- Haining, R. P. (1990). Spatial Data Analysis in the Social and Environmental Sciences. Cambridge: Cambridge University Press.
- Haining, R. P. (1991). Bivariate correlation with spatial data. Geographical Analysis, 23, 210-227.
- Haining, R. P. (2003). Spatial Data Analysis: Theory and Practice. Cambridge: Cambridge University Press.
- Hall, K. R. & Maruca, S. L. (2001). Mapping a forest mosaic: a comparison of vegetation and songbird distributions using geographic boundary analysis. *Plant Ecology*, **156**, 105–120.
- Halley, J. D. & Winkler, D. A. (2008). Consistent concepts of self-organization and self-assembly. *Complexity*, 14, 10-17.
- Halley, J. M., Hartley, S., Kallimanis, A. S. et al. (2004). Uses and abuses of fractal methodology in ecology. Ecology Letters, 7, 254–271.
- Handcock, R. & Csillag, F. (2002). Ecoregionalization assessment: spatio-temporal analysis of net primary production across Ontario. Écoscience, 9, 219-230.
- Hansen, A. & di Castri, F. (1992). Landscape Boundaries: Consequences for Biotic Diversity and Ecological Flows. New York: Springer-Verlag.
- Hanski, I. (2009). Metapopulations and spatial population processes. In *The Princeton Guide to Ecology*, ed. S. A. Levin, pp. 177–185. Princeton: Princeton University Press.
- Hanski, I. & Woiwod, I. P. (1993). Spatial synchrony in the dynamics of moth and aphid populations. *Journal of Animal Ecology*, 62, 656–668.
- Harary, F. (1967). Topological concepts in graph theory. In A Seminar on Graph Theory, ed. F. Harary, pp. 13–24. New York: Holt, Rinehart & Winston.
- Harary, F. (1969). Graph Theory. Reading, MA: Addison-Wesley.
- Harary, F. & Gupta, G. (1997). Dynamic graph models. Mathematical and Computer Modelling, 25, 79–87.
- Hargrove, W. W., Hoffman, F. M. & Schwartz, P. M. (2002).
 A fractal landscape realizer for generating synthetic maps.
 Conservation Ecology, 6, 2. Online www.consecol.org/vol6/iss1/art2.
- Harper, K. A., Macdonald, S. E., Burton, P. et al. (2005). Edge influence on forest structure and composition in fragmented landscapes. Conservation Biology, 19, 768-782.

- Harrison, P. J., Hanski, I. & Ovaskainen, O. (2011). Bayesian state-space modeling of metapopulation dynamics in the Glanville fritillary butterfly. *Ecological Monographs*, 8, 581-598.
- Harrison, S., Davies, K. F., Safford, H. D. & Veirs, J. H. (2006).
 Beta diversity and the scale-dependence of the productivity-diversity relationship: a test in the Californian serpentine flora. *Journal of Ecology*, 94, 110–117.
- Harvey, E. & Miller, T. E. (1996). Variance in composition of inquiline communities in leaves of *Sarracenia purpurea* L. on multiple spatial scales. *Oecologia*, **108**, 562–566.
- Hastings, A. (2009). Biological chaos and complex dynamics. In *The Princeton Guide to Ecology*, ed. S. A. Levin, pp. 172–176. Princeton: Princeton University Press.
- Heagerty, P. J. & Lumley, T. (2000). Window subsampling of estimating functions with application to regression models. *Journal of the American Statistical Society*, 95, 197–211.
- Hedley, S. L. & Buckland, S. T. (2004). Spatial models for line transect sampling. *Journal of Agricultural, Biological and Environmental Statistics*, 9, 181–199.
- Heino, J. & Muotka, T. (2005). Highly nested snail and clam assemblages in boreal lake littorals: Roles of isolation, area, and habitat suitability. *Ecoscience*, 12, 141-146.
- Hendry, R. J. & McGlade, J. M. (1995). The role of memory in ecological systems. Proceedings of the Royal Society of London Series B, Biological Sciences, 259, 153–159.
- Henebry, G. M. (1995). Spatial model error analysis using autocorrelation indices. *Ecological Modelling*, **82**, 75–91.
- Hengl, T., Heuvelink, G. B. M. & Rossiter, D. G. (2007). About regression-kriging: from equations to case studies. *Computers and Geosciences*, 33, 1301–1315.
- Hengl, T., Heuvelink, G. B. M. & Stein, A. (2004). A generic framework for spatial prediction of soil variables based on regression-kriging. *Geoderma*, 120, 75–93.
- Higashi, M., Yamamura, N., Abe, T. & Burns, T. P. (1991). Why don't all termite species have a sterile worker caste? Proceedings of the Royal Society of London Series B, Biological Sciences, 246, 25–29.
- Hill, M. O. (1973). The intensity of spatial pattern in plant communities. *Journal of Ecology*, 61, 225–235.
- Hoagland, B. W. & Collins S. L. (1997). Heterogeneity in short-grass prairie vegetation: the role of playa lakes. *Journal of Vegetation Science*, 8, 277–286.
- Holden, Z. A. & Evans, J. S. (2010). Using fuzzy C-means and local autocorrelation to cluster satellite-inferred burn severity classes. *International Journal of Wildland Fire*, 19, 853–860.
- Holland, M. M., Risser, P. G. & Naiman, R. J. (eds.) (1991). Ecotones. New York: Chapman & Hall.

- Holyoak, M., Leibold, M. A. & Holt, R. D. eds. (2005). Metacommunities: Spatial Dynamics and Ecological Communities. Chicago: University of Chicago Press.
- Hope, A. C. A. (1968). A simplified Monte Carlo significance test procedure. *Journal of the Royal Statistical Society B*, 30, 582–598.
- Hopkins, B. (1957). Pattern in the plant community. *Journal of Ecology*, **45**, 451–463.
- Hothorn, T., Müller, J., Schröder, B., Kneib, T. & Brandl, R. (2011). Decomposing environmental, spatial, and spatiotemporal components of species distributions. *Ecological Monographs*, 81, 329–347.
- Hubert, L. J., & Golledge, R. G. (1981). A heuristic method for the comparison of related structures. *Journal of Mathematical Psychology*, 23, 214–226.
- Hudak, A. T., Lefsky, M. A., Cohen, W. B. & Berterretche, M. (2002). Integration of LiDAR and Landsat ETM plus data for estimating and mapping forest canopy height. *Remote Sensing of Environment*, 82, 397–416.
- Hufkens K., Scheunders, P. & Ceulemans, R. (2009). Ecotones in vegetation ecology: methodologies and definitions revisited. *Ecological Research*, 24, 977–986.
- Hughes, J.S., Fortin, M.-J., Nealis, V. & Régnière, J. (2014).Pollen cone production in jack pine: spatial and temporal patterns subject to natural disturbance by the jack pine budworm. Candeain Journal of Forest Research, 44, 195–211.
- Hui, C. (2009). On the scaling patterns of species spatial distribution and association. *Journal of Theoretical Biology*, 261, 481-487.
- Hui, C., Veldtman, R. & McGeoch, M. (2010). Measures, perceptions and scaling patterns of aggregated species distributions. *Ecography*, 33, 95-102.
- Hunsaker C., Goodchild, M., Friedl, M. & Case, T., eds. (2001).
 Spatial Uncertainty in Ecology. Implications for Remote Sensing and GIS Applications. New York: Springer-Verlag.
- Hurlbert, S. H. (1984). Pseudoreplication and the design of ecological field experiments. *Ecological Monographs*, 54, 187–211.
- Huson, D. H. & Bryant, D. (2006). Application of phylogenetic networks in evolutionary studies. *Molecular Biology and Evolution*, 23, 254–267.
- Huston, M. A. (1994). Biological Diversity: The Coexistence of Species on Changing Landscapes. New York: Cambridge University Press.
- Ichinose, S. (2001). Transient structures of wave patterns arising in the wave regeneration of subalpine coniferous forests. *Physics Reviews*, E64, 061903.
- Illian, J., Penttinen, A. & Stoyan, D. (2008). Statistical Analysis and Modeling of Spatial Point Patterns. Chichester: Wiley.

- Ings, T. C., Montoya, J. M., Bascompte, J. et al. (2009). Ecological networks-beyond food webs. *Journal of Animal Ecology*, 78, 253-269. Not seen, author?
- Isaaks, E. H. & Srivastava, R. M. (1989). An Introduction to Applied Geostatistics. Oxford: Oxford University Press.
- Ives, A. R. & Zhu, J. (2006). Statistics for correlated data: phylogenies, space, and time. *Ecological Applications*, 16: 20–32.
- Iyengar, V. S. (2005). Space-time clusters with flexible shapes. Morbidity and Mortality Weekly Report, 54 (Supplement), 71-76.
- Jacquemyn, H., Honnay, O. & Pailler, T. (2007). Range size variation, nestedness, and species turnover of orchid species along an altitudinal gradient on Reunion Island: implications for conservation. *Biological Conservation*, 136, 388–397.
- Jacquez, G. M. (1995). The map comparison problem: tests for the overlap of geographical boundaries. *Statistical Medicine*, 14, 2343–2361.
- Jacquez, G. M. (1996). A k nearest neighbour test for spacetime interaction. Statistical Medicine, 15, 1935–1949.
- Jacquez, G. M., Kaufmann, A. & Goovaerts, P. (2008). Boundaries, links and clusters: a new paradigm in spatial analysis? Environmental and Ecological Statistics, 15, 403–419.
- Jacquez, G. M., Maruca, S. L. & Fortin, M.-J. (2000). From fields to objects: areview of geographic boundary analysis *Journal of Geographical Systems*, 2, 221–241.
- James P. M. A. & Fortin, M.-J. (2012). Ecosystems and spatial patterns. In *Encyclopedia of Sustainability Science and Tech*nology, ed. R. A. Meyers, pp. 3326–3342. Springer Science +Business Media.
- James, P. M. A., Fleming, R. A. & Fortin, M.-J. (2010). Identifying significant scale-specific spatial boundaries using wavelets and null models: spruce budworm defoliation in Ontario, Canada as a case study. *Landscape Ecology*, 25, 873–887.
- James, P. M. A., Fortin, M.-J., Fall, A., Kneeshaw, D. & Messier, C. (2007). The effects of spatial legacies following shifting management practices and fire on boreal forest age structure. *Ecosystems*, 10, 1261–1277.
- James P. M. A., Sturtevant, B. R., Townsend, P., Wolter, P. & Fortin, M.-J. (2011). Two-dimensional wavelet analysis of spruce budworm host basal area in the Border Lakes landscape. *Ecological Applications*, 21, 2197–2209.
- Jelinski, D. E. & Wu, J. (1996). The modifiable areal unit problem and implications for landscape ecology. *Landscape Ecology*, 3, 129–140.
- Jenkins, G. M. & Watts, D. G. (1968). Spectral Analysis and Its Applications. San Francisco: Holden-Day.

- Jetz, W., Rahbek, C. & Lichstein, J. W. (2005). Local and global approaches to spatial data analysis in ecology. *Global Ecology & Biogeography*, 14, 97–98.
- Johnson, E. A. (1992). Fire and Vegetation Dynamics. Cambridge: Cambridge University Press.
- Johnson, J. B. & Omland, K. S. (2004). Model selection in ecology and evolution. *Trends in Ecology & Evolution*, 19, 101-108.
- Johnston, C. A., Pastor, J. & Pinay, G. (1992). Quantitative methods for studying landscape boundaries. In *Landscape Boundaries*, eds. A. J. Hansen & F. di Castri, pp. 107–125. New York: Springer-Verlag.
- Jombart, T., Devillard, S., Dufour, A. B. & Pontier, D. (2008).Revealing cryptic spatial patterns in genetic variability by a new multivariate method. *Heredity*, 101, 92–103.
- Jombart, T., Dray, S. & Dufour, A. B. (2009). Finding essential scales of spatial variation in ecological data: A multivariate approach. *Ecography*, 32, 161–168.
- Jones, D. D. (1977). Catastrophe theory applied to ecological systems. Simulation, 29, 1-15.
- Jones, E. W. (1945). The structure and reproduction of the virgin forest of the north temperate zone. *New Phytologist*, 44, 130-148.
- Jongman, R. H., ter Braak, C. J. F. & Van Tongeren, O. F. R. eds. (1995). *Data Analysis in Community and Landscape Ecology*. Cambridge: Cambridge University Press.
- Jordan, G. J., Fortin, M.-J. & Lertzman, K. P. (2008). Spatial pattern and persistence of historical fire boundaries in southern interior British Columbia. *Environmental and Ecological Statistics*, 15, 523–535.
- Jost, L. (2006). Entropy and diversity. Oikos, 113, 363-375.
- Jost, L. (2007). Partitioning diversity into alpha and beta components. *Ecology*, 88, 2427–2439.
- Journel, A. G. & Huijbregts, C. (1978). Mining Geostatistics. London: Academia Press.
- Juhasz-Nagy, P (1993). Notes on compositional diversity. Hydrobiologia, 249, 173-182.
- Kabos, S. & Csillag, F. (2002). The analysis of spatial association of nominal data on regular lattice by join-count-statistic without first-order homogeneity. *Computers and Geosciences*, 28, 901–910.
- Kamarianakis, Y. & Prastacos, P. (2003). Spatial time-series modeling: a review of the proposed methodologies. Regional Economics Applications Laboratory (REAL) Discussion Papers, REAL 03-T-19.
- Kareiva, P. & Shigesada, N. (1983). Analyzing insect movement as a correlated random walk. *Oecologia*, 56, 234–238.
- Keddy, P. A. (1991). Working with heterogeneity: an operator's guide to environmental gradients. In *Ecological*

- *Heterogeneity*, eds. J. Kolasa and S. T. A. Pickett. New York: Springer Verlag.
- Keitt, T. H. & Fischer, J. (2006). Detection of sale-specific community dynamics using wavelets. *Ecology*, 87, 2895– 2904.
- Keitt, T. H. & Urban, D. L. (2005). Scale-specific inference using wavelet. *Ecology*, 86, 2497–2504.
- Keitt, T. H., Urban, D. L. & Milne, B. T. (1997). Detecting critical scales in fragmented landscapes. *Conservation Ecology* [online], 1, 4. URL: http://www.consecol.org/vol1/iss1/art4/.
- Kelly, R. P., Oliver T. A. & Sivasundar, A. (2010). A method for detecting population genetic structure in diverse, high gene-flow species. *Journal of Heredity* 101, 423–436.
- Kempe, D., Kleinberg, J. & Kumar, A. (2002). Connectivity and inference problems in temporal networks. *Journal of Com*puter and System Sciences, 64, 820–842.
- Kenkel, N. C. (1993). Modeling Markovian dependence in populations of *Aralia nudicaulis*. Ecology, 74, 1700–1706.
- Kenkel, N. C. & Walker, D. J. (1993). Fractals and ecology. Abstracta Botanica, 17, 53-70.
- Kenkel, N. C., Hendrie, M. L. & Bella, I. E. (1997). A long-term study of *Pinus banksiana* population dynamics. *Journal of Vegetation Science*, 8, 241–254.
- Kidd, D. M. & Ritchie, M. G. (2006). Phylogeographic information systems: putting the geography into phylogeography. *Journal of Biogeography*, 33, 1851-1865.
- Kimura, M. & Crow, J. F. (1964). Number of alleles that can be maintained in a finite population. *Genetics*, 49, 725–738.
- Kingman, J. F. C. (1993). Poisson Processes. Oxford: Oxford University Press.
- Kirkpatrick, D. G. & Radke, J. D. (1985) A framework for computational morphology. Computational Geometry, Machine Intelligence and Pattern Recognition, 2, 217–248.
- Kneitel, J. M. & Miller, T. E. (2003) Dispersal rates affect species composition in metacommunities of *Sarracenia* purpurea inquilines. American Naturalist, 162, 165–171.
- Knox, E. G. (1964). The detection of space-time interactions. Applied Statistics, 13, 25–29.
- Koen, E. L., Garroway, C. J., Wilson, P. J. & Bowman, J. (2010). The effect of map boundary on estimates of landscape resistance to animal movement. *PLoS One* 5(7): e11785.
- Koenig, W. D. & Knops, J. M. (1998). Testing for spatial autocorrelation in ecological time series. *Ecography*, 21, 423–429.
- Kolaczyk, E. D. (2009). Statistical Analysis of Network Data. New York: Springer.
- Koleff, P., Gaston, K. J. & Lennon, J. J. (2003). Measuring beta diversity for presence absence data. *Journal of Animal Ecol*ogy, 72, 367–382.

- König, D., Carvajal-Gonzalez, S., Downs, A. M., Vassy, J. & Rigaut, J. P. (1991). Modeling and analysis of 3-D arrangements of particles by point processes with examples of application to biological data obtained by confocal scanning light microscopy. *Journal of Microscopy*, 161, 405-433.
- Kost, C., de Olieira, E. G., Knoch, T. A. & Wirth, R. (2005). Spatio-temporal permanence and plasticity of foraging trails in young and mature leaf-cutting ant colonies (*Atta* spp.). *Journal of Tropical Ecology*, 21, 677–688.
- Kostakos, V. (2009). Temporal graphs. Physica A, 388, 1007– 1023.
- Koutaniemi, L. (1999). Twenty-one years of string movements on the Liippasuo aapa mire, Finland. Boreas, 28, 521–530.
- Krebs, C. J. (2002). Ecology: The Experimental Analysis of Distribution and Abundance: Hands-On Field Package, 5th edn. San Francisco, CA: Prentice Hall.
- Krebs, C. J., Boutin, S. & Boonstra, R. (2001). Ecosystem Dynamics of the Boreal Forest: The Kluane Project. Oxford: Oxford University Press.
- Krige, D. G. (1966). Two-dimensional weighted moving average trend surfaces for ore-evaluation. *Journal of the South Africa Institute of Mining and Metallurgy*, 66, 13–38.
- Kulldorff, M. (1997). A spatial scan statistic. Communications in statistics: Theory and Methods. 26: 1481–1496.
- Kulldorff, M., Athas, W., Feuer, E., Miller, B. A. & Key, C. (1998). Evaluating cluster alarms: a space-time scan statistic and brain cancer in Los Alamos. *American Journal of Public Health*, 88, 1377–1380.
- Kuldorff, M., Heffernan, R., Hartman, J., Assunção, R. M. & Mostashari, F. (2005). A space-time permutation scan statistic for the early detection of disease outbreaks. *PLoS Medicine*, 2, 216–224.
- Lahiri, S. N. (1999). Theoretical comparisons of block bootstrap methods. *The Annals of Statistics*, 27, 386-404.
- Lambin, X., Elston, D. A., Petty, S. J. & MacKinnon, J. L. (1998).
 Spatial asynchrony and periodic traveling waves in cyclic populations of field voles. *Proceedings of the Royal Society of London Series B*, 265, 1491–1496.
- Lamour, A., Termorshuizen, A. J., Volker, D. & Jeger, M. J. (2007). Network formation by rhizomorphs of *Armillaria lutea* in natural soil: their description and ecological significance. FEMS *Microbial Ecology*, 62, 222–232.
- Lande, R. (1996). Statistics and partitioning of species diversity, and similarity among multiple communities. Oikos, 76, 5–13.
- Lande, R., Engen, S. & Sather, B. E. (2003). Stochastic Population Dynamics in Ecology and Conservation. Oxford: Oxford University Press.
- Lantuéjoul, C. (2002). Geostatistical Simulation, Models, and Algorithms. Berlin: Springer.

- Law, R., Herben, T. & Dieckmann, U. (1997). Non-manipulative estimates of competition coefficients in a montane grassland community. *Journal of Ecology*, 85, 505–517.
- Ledo, A., Condés, S. & Montes, F. (2011). Intertype mark correlation function: A new tool for the analysis of species interactions. *Ecological Modelling*, 222, 2580–2587.
- Leduc, A., Drapeau, P., Bergeron, Y. & Legendre, P. (1992). Study of spatial components of forest cover using partial Mantel tests and path analysis. *Journal of Vegetation Science*, 3, 69–78.
- Legendre, P. (1993). Spatial autocorrelation: trouble or new paradigm? *Ecology*, 74, 1659-1673.
- Legendre, P. (2000). Comparison of permutation methods for the partial correlation and partial Mantel tests. *Journal of Statistical Computer Simulation*, 67, 37–73.
- Legendre, P. (2008). Studying beta diversity: ecological variation partitioning by multiple regression and canonical analysis. *Journal of Plant Ecology*, 1, 3–8.
- Legendre, P. & De Cáceres, M. (2013). Beta diversity as the variance of community data: dissimilarity coefficients and partitioning. *Ecology Letters*, 16, 951-963.
- Legendre, P. & Fortin, M.-J. (1989). Spatial pattern and ecological analysis. *Vegetatio*, 80, 107–138.
- Legendre, P. & Fortin, M.-J. (2010). Comparison of the Mantel test and alternative approaches for detecting complex multivariate relationships in the spatial analysis of genetic data. *Molecular Ecology Resources*, 10, 831–844.
- Legendre, P. & Legendre, L. (1998). *Numerical Ecology*, 2nd English edn. Amsterdam: Elsevier.
- Legendre, P. & Legendre, L. (2012). Numerical Ecology, 3rd English edn. Amsterdam: Elsevier.
- Legendre, P. & Troussellier, M. (1988). Aquatic heterotrophic bacteria: modeling in the presence of spatial autocorrelation. *Limnology and Oceanography*, 33, 1055-1067.
- Legendre, P., Borcard, D., & Peres-Neto, P.R. (2005). Analyzing beta diversity: partitioning the spatial variation of community composition data. *Ecological Monographs*, 75, 435–450.
- Legendre, P., Dale, M. R. T., Fortin, M.-J., Casgrain, P. & Gurevitch, J. (2004). Effects of spatial structures on the results of field experiments. *Ecology*, 85, 3202–3214.
- Legendre, P., Dale, M. R. T., Fortin, M.-J. et al. (2002). The consequences of spatial structure for the design and analysis of ecological field surveys. Ecography, 25, 601-615.
- Legendre, P., Lapointe, F.-J. & Casgrain, P. (1994). Modeling brain evolution from behavior: a permutational regression approach. *Evolution*, 48, 1487–1499.
- Legendre, P., Sokal, R. R., Oden, N. L., Vaudor, A. & Kim, J. (1990). Analysis of variance with spatial autocorrelation in

- both the variable and the classification criterion. *Journal of Classification*, **7**, 53–75.
- Leibold, M. A. (2009). Spatial and metacommunity dynamics in biodiversity. In *The Princeton Guide to Ecology*, ed. S. A. Levin, pp. 312–319. Princeton: Princeton University Press.
- Leitão, A. B., Miller, J., Ahern, J. & McGarigal, K. (2006). Measuring Landscapes: A Planner's Handbook. Washington, D.C.: Island Press.
- Lejeune, O. & Tlidi, M. (1999). A model for the explanation of vegetation stripes (tiger bush). *Journal of Vegetation Sci*ence, 10, 201–208.
- Lele, S. (1991). Jackknifing linear estimating equations: asymptotic theory and applications in stochastic processes. *Journal of the Royal Statistical Society B*, 53, 253–267.
- Lennon, J. J. (2000). Red-shifts and red herrings in geographical ecology. *Ecography* 23, 101–113.
- Lesne, A. (2006). Complex networks: from graph theory to biology. Letters in Mathematical Physics, 78, 235–262.
- Leung, B., Bossenbroek, J. M. & Lodge, D. M. (2006). Boats, pathways, and aquatic biological invasions: estimating dispersal potential with gravity models. *Biology Invasions*, 8, 241–254.
- Leung, Y. (1987). On the imprecision of boundaries. Geographical Analysis, 19, 125–151.
- Levin, S. A. (1992). The problem of pattern and scale in ecology. *Ecology*. **73**, 1943–1967.
- Levin, S. A. (2000). Multiple scales and the maintenance of biodiversity. *Ecosystems*, 3, 498–506.
- Levin, S. A. ed. (2009). The Princeton Guide to Ecology. Princeton: Princeton University Press.
- Li, H. & Reynolds, J. F. (1995). On definition and quantification of heterogeneity. *Oikos*, 73, 280-284.
- Liang, S. D., Banerjee, S. & Carlin, B. P. (2009) Bayesian wombling for spatial point processes. *Biometrics*, 65, 1243–1253.
- Lichstein, J. W. (2007). Multiple regression on distance matrices: A multivariate spatial analysis tool. *Plant Ecology*, **188**, 117–131.
- Lin, K.-P., Long, Z.-H. & Ou, B. (2011). The size and power of bootstrap tests for spatial dependence in a linear regression model. *Computational Economics* 38, 153–171.
- Lindeberg, T. (1994). Scale-Space Theory in Computer Vision. Dordrecht: Kluwer.
- Little, L. R. & Dale, M. R. T. (1999) A method for analysing spatio-temporal pattern in plant establishment, tested on a Populus balsamifera clone. Journal of Ecology, 87, 620–627.
- Liu, C. (2001). A comparison of five distance-based methods for pattern analysis. *Journal of Vegetation Science*, 12, 411–416.
- Loh, J. M. (2008). A valid and fast spatial bootstrap for correlation functions. *The Astrophysical Journal*. 681, 726–734.

- Lomnicki, Z. A. & Zaremba, S. K. (1955). Some applications of zero-one processes. *Journal of the Royal Statistical Society*, ser. B, 17, 243–255.
- Lomolino, M. V. (2006). Investigating causality of nestedness of insular communities: selective immigrations or extinctions? *Journal of Biogeography*, 23, 699–703.
- Loreau, M. (2010). *From Populations to Ecosystems*. Princeton: Princeton University Press.
- Lotwick, H. W. & Silverman, B. W. (1982). Methods for analysing spatial processes of several types of points. *Journal of the Royal Statistical Society B*, 44, 406-413.
- Lovett Doust, J. (1981). Population dynamics and local specialization in a clonal perennial (*Ranunculus repens*): I. The dynamics of ramets in contrasting habitats. *Journal of Ecol*ogy, 69, 743–755.
- Lowell, K. (1997). Effect(s) of the "no-same-color-touching" constraint on the join-count statistic: a simulation study. *Geographical Analysis*, 29, 339–353.
- Lu, H., & Carlin, B. P. (2005). Bayesian areal wombling for geographical boundary analysis. *Geographical Analysis*, 37, 265–285.
- Ludwig, J. A. & Cornelius, J. M. (1987). Locating discontinuities along ecological gradients. *Ecology*, 68, 448–450.
- Ludwig, J. A. & Goodall, D. W. (1978). A comparison of paired with blocked-quadrat variance methods for the analysis of spatial pattern. *Vegetatio*, 38, 49–59.
- Ludwig, J. A. & Reynolds, J. F. (1988). *Statistical Ecology*. New York: Wiley.
- MacArthur, R. H. (1965). Patterns of species diversity. Biological Reviews of the Cambridge Philosophical Society, 40, 510-533.
- Mack, R. N., & Harper, J. L. (1997). Interference in dune annuals - spatial pattern and neighborhood effects. *Journal* of *Ecology*, 65, 345–363.
- MacKinnon, J. L., Petty, S. J., Elston, D. A. et al. (2001). Scale invariant spatio-temporal patterns of field vole density. *Journal of Animal Ecology*, 70, 101–111.
- Magurran, A. E. (2004). *Measuring Biological Diversity*. Oxford: Blackwell.
- Magurran, A. E. & McGill, B. J. eds. (2010). Biological Diversity: Frontiers in Measurement and Assessment. Oxford: Oxford University Press.
- Mahecha, M. D. & Schmidtlein, S. (2008). Revealing biogeographical patterns by nonlinear ordinations and derived anisotropic spatial filters. Global Ecology & Biogeography, 17, 284–296.
- Mahli, Y. & Román-Cuesta, R. M. (2008). Analysis of lacunarity and scales of spatial homogeneity in IKONOS images of Amazonian tropical forest canopies. *Remote Sensing of the Environment.* 112, 2074–2087.

- Mamaghani, M. J., Andersson, M. & Krieger, P. (2010). Spatial point pattern analysis of neurons using Ripley's K-function in 3D. Frontiers in Neuroinformatics, 4, Article 9. doi: 10.3389/fninf.2010.00009.
- Mandelbrot, B. B. (1983). *The Fractal Geometry of Nature*. New York: Freeman.
- Manly, B. F. J. (1997). Randomization, Bootstrap, and Monte Carlo Methods in Biology, 2nd edn. London: Chapman & Hall
- Manly, B. F. J. (2006). Randomization, Bootstrap, and Monte Carlo Methods in Biology, 3rd edn. Boca Raton: Chapman & Hall/CRC.
- Manly, B. F. J., McDonald, L. L., Thomas, D. L., McDonald,
 T. L. & Erickson, W. P. (2002). Resource Selection by
 Animals: Statistical Design and Analysis for Field Studies,
 2nd edn. Dordrecht: Kluwer Academic.
- Mantel, N. (1967). The detection of disease clustering and a generalized regression approach. Cancer Research, 27, 209–220.
- Marj, T., Tuia, D. & Ratle, F. (2006). Detection of clusters using space-time scan statistics. *International Journal of Wildland Fire*, 18, 830-836.
- Marquet, P. A. (2009). Macroecological perspectives on communities and ecosystems. In *The Princeton Guide to Ecology*, ed. Levin, S. A., pp. 257–63. Princeton: Princeton University Press.
- Marr, D. & Hildreth, E. (1980). Theory of edge detection. Proceedings of the Royal Society of London, Series B, 207, 187–217.
- Mason, O. & Verwoerd, M. (2007). Graph theory and networks in biology. *IET Systems Biology*, 1, 89–119.
- Matheron, G. (1970). La théeorie des variables réegionaliséees, et ses applications. Les Cahiers du Centre de Morphologie Mathéematique de Fontainebleau. Fontainebleau: Fascicule 5.
- Matula, D. W. & Sokal, R. R. (1980). Properties of Gabriel graphs relevent to geographic variation research and the clustering of points in the plane. *Geographical Analysis*, 12, 205–222
- May, R. M. (1977). Thresholds and breakpoints in ecosystems with a multiplicity of stable states. *Nature*, 269, 471-477.
- McAbendroth, L., Foggo, A., Rundle, S. D. & Bilton, D. (2005).Unravelling nestedness and spatial pattern in pond assemblages. *Journal of Animal Ecology*, 74, 41–44.
- McBratney, A. B. & de Gruijter, J. J. (1992). A continuum approach to soil classification by modified fuzzy *k*-means with extra grades. *Journal of Soils Science*, **43**, 159–176.
- McCoy, E. D., Bell, S. S. & Walters, K. (1986). Identifying biotic boundaries along environmental gradients. *Ecology*, **67**, 749–759.
- McGarigal, K. & Marks, B. J. (1995). FRAGSTATS: spatial pattern analysis program for quantifying landscape structure.

- US Forest Service General Technical Report PNW 351. Corvallis, OR: US Forest Service.
- McIntire, E. J. B. & Fajardo, A. (2009). Beyond description: the active and effective way to infer processes from spatial patterns. *Ecology*, **90**, 46–56.
- McIntire, E. J. B. & Fortin, M.-J. (2006) Structure and function of wildfire and mountain pine beetle forest boundaries. *Ecography*, 29, 309–318.
- McKnight, M. W., White, P. S., McDonald, R. I. et al. (2007).Putting beta-diversity on the map: broad-scale congruence and coincidence in the extremes. PLoS Biology, 5, 2424–2432.
- McRae B. H., Dickson B. G., Keitt T. H. & Shah V. B. (2008).Using circuit theory to model connectivity in ecology, evolution, and conservation. *Ecology*, 89, 2712–24.
- Melián, C. J., Bascompte, J. & Jordano, P. (2005). Spatial structure and dynamics in a marine food web. In *Aquatic Food Webs: An Ecosystem Approach*, eds. A. Belgrano, U. Scharler, J. Dunne & R. E. Ulanowicz, pp. 19–24. New York: Oxford University Press.
- Melles, S. J., Badzinski, D., Fortin, M.-J., Csillag F. & Lindsay, K. (2009). Disentangling habitat and social drivers of nesting patterns in songbirds. *Landscape Ecology*, 24, 519–531.
- Melles, S. J., Fortin, M.-J., Lindsay, K. & Badzinski, D. (2011).
 Expanding northward: influence of climate change, forest connectivity, and population processes on a threatened species' range shift. Global Change Biology, 17, 17–31.
- Meyers, L. A. (2007). Contact network epidemiology: bond percolation applied to infectious disease prediction and control. *Bulletin of the American Mathematical Society*, **44**, 63_86
- Mielke Jr., P. W. & Berry, K. J. (2007) *Permutation Methods.*A Distance Function Approach, 2nd edn. New York: Springer.
- Miller, M. P. (2005). Alleles In Space (AIS): computer software for the joint analysis of inter-individual spatial and genetic information. *Journal of Heredity*, 96, 722–724.
- Miller, M. P., Haig, S. M. & Wagner, R. S. (2006). Phylogeography and spatial genetic structure of the Southern torrent salamander: implications for conservation and management. *Journal of Heredity*, 97, 561–570.
- Millspaugh, J. J., Skalski, J. R., Kernohan, B. J. et al. (1998).Some comments on spatial independence in studies of resource selection. Wildlife Society Bulletin, 26, 232–236.
- Milne, B. T. (1992). Spatial aggregation and neutral models in fractal landscapes. *American Naturalist*, 139, 32–57.
- Minasny, B. & McBratney, A. B. (2005). The Matérn function as a general model for soil variograms. *Geoderma*, 128, 192–207.

- Minchin, P. R. (1989). Montane vegetation of the Mt. Field Massif, Tasmania – a test of some hypotheses about properties of community patterns. *Vegetatio*, 83, 97–110.
- Minta, S. C. (1992). Tests of spatial and temporal interaction among animals. *Ecological Applications*, 2, 178–188.
- Mithen, R., Harper, J. L. & Weiner, J. (1984). Growth and mortality of individual plants as a function of "available area". *Oecologia*, 62, 57–60.
- Mizon, G. E. (1995). A simple message for autocorrelation correctors: don't. *Journal of Econometrics*, 69, 267–289.
- Mladenoff, D. J., & Baker, W. L. (1999). Progress and future directions in spatial modeling of forest landscapes. Chapter 13 in Advances in Spatial Modeling of Forest Landscape Change: Approaches and Applications, eds. D. J. Mladenoff & W. L. Baker. Cambridge: Cambridge University Press.
- Moorcroft, P., & Lewis, M. A. (2006). Mechanistic Home Range Analysis. Princeton: Princeton University Press.
- Moran, P. A. P. (1947). The random division of an interval. Journal of the Royal Statistical Society, Supplement, 9, 92–98.
- Moran, P. A. P. (1948). The interpretation of statistical maps. Journal of the Royal Statistical Society B, 10, 243–251.
- Morrison, L. W. (2013). Nestedness in insular floras: spatiotemporal variation and underlying mechanisms. *Journal of Plant Ecology*. doi: 10.1093/pe/rtt002.
- Morsdorf, F., Meier, E., Kötz, B. *et al.* (2004). LIDAR-based geometric reconstruction of boreal type forest stands at single tree level for forest and wildland fire management. *Remote Sensing of Environment*, **92**, 353–362.
- Moss, R., Elston, D. A. & Watson, A. (2000). Spatial asynchrony and demographic traveling waves during red grouse population cycles. *Ecology*, 81, 981–989.
- Mouquet, N. & Loreau, M. (2002). Coexistence in metacommunities: the regional similarity hypothesis. *American Naturalist*, 159, 420–426.
- Mugglestone, M. A. (1996). The role of tessellation methods in the analysis of three-dimensional spatial point patterns. *Journal of Agricultural, Biological & Environmental Statis*tics, 1, 141–153.
- Mugglestone, M. A. & Renshaw, E. (1996). A practical guide to the spectral analysis of spatial point processes. *Computational Statistics & Data Analysis*, 21, 43–65.
- Munguia-Rosas, M. A. & Sosa, V. J. (2008). Nurse plants vs. nurse objects: Effects of woody plants and rocky cavities on the recruitment of the *Pilosocereus leucocephalus* columnar cactus. *Annals of Botany*, 101, 175–185.
- Murakami, M. & Hirao, T. (2010). Lizard predation alters the effect of habitat area on the species richness of insect assemblages on Bahamian isles. *Diversity and Distributions*, 16, 952–958.

- Murphy, M., Dezzani, R., Pilliod, D. & Storfer, A. (2010). Landscape genetics of high mountain frog metapopulations. *Molecular Ecology*, 19, 3634–3649.
- Nekola, J. C. & White P. S. (1999). The distance decay of similarity in biogeography and ecology. *Journal of Biogeog*raphy, 26, 867–878.
- Nestel, D. & Klein, M. (1995). Geostatistical analysis of leafhopper (Homoptera: Cicadellidae) colonization and spread of deciduous orchards. *Environmental Entomology*, 24, 1032–1039.
- Neu, C. W., Byers, C. R., Peek, J. M. & Boy, V. (1974). A technique for analysis of utilization-availability data. *Journal of Wildlife Management*, 38, 541–545.
- Newman, M. E. J. (2006). Modularity and community structure in networks. Proceedings of the National Academy of Sciences, 103, 8577–8582.
- Ng, K. K. S., Lee, S. L. & Koh, C. L. (2004). Spatial structure and genetic diversity of two tropical tree species with contrasting breeding systems and different ploidy levels. *Molecular Ecology*, 13, 657–669.
- Niklas, K. J. & Norstog, K. (1984). Aerodynamics and pollen grain depositional patterns on cycad megastrobili: implications on the reproduction of three cycad genera (Cycas, Dioon, and Zamia). *Botanical Gazette*, 145, 92–104.
- Nikoloski, Z., Grimbs, S., May, P. & Selbig, J. (2008). Metabolic networks are NP-hard to reconstruct. *Journal of Theoretical Biology*, 254, 807–816.
- Nosil, P. & Rundle, H. (2009). Ecological speciation: Natural selection and the formation of new species. In *The Prince*ton Guide to Ecology, ed. S. A. Levin, pp. 134–142. Princeton: Princeton University Press.
- Noy-Meir, I. & Anderson, D. (1971). Multiple pattern analysis or multiscale ordination: towards a vegetation hologram. In *Statistical Ecology, vol. 3: Populations, Ecosystems, and Systems Analysis*, eds. G. P. Patil, E. C. Pielou & W. E. Waters, pp. 207–232. University Park: Pennsylvania State University Press.
- O'Brien D., Manseau, M., Fall, A. & Fortin, M.-J. (2006). Testing the importance of spatial configuration of winter habitat for woodland caribou: an application of graph theory. *Biological Conservation*, **130**, 70-83.
- O'Neill, R. V., Hunsaker, C. T., Timmins, S. P. et al. (1996). Scale problems in reporting landscape pattern at the regional scale. *Landscape Ecology*, 11, 169–180.
- O'Neill, R. V., Krummel, J. R., Gardner, R. H. *et al.* (1988). Indices of landscape pattern. *Landscape Ecology*, **1**, 153–162.
- O'Neill, R. V., Riitters, K. H., Wickham, J. D. & Jones, K. B. (1999). Landscape pattern metrics and regional assessment. *Ecosystem Health*, **5**, 225–233.

- O'Sullivan, D. & Unwin, D. (2003). Geographic Information Analysis. New Jersey: Wiley.
- O'Sullivan, D. & Unwin, D. (2010). Geographic Information Analysis, 2nd edn. New Jersey: Wiley.
- Oden, N. L. (1984). Assessing the significance of a spatial correlogram. *Geographical Analysis*, **16**, 1–16.
- Oden, N. L. & Sokal, R. R. (1986). Directional autocorrelation: an extension of spatial correlograms to two dimensions. Systematic Zoology, 35, 608–617.
- Oden, N. L. & Sokal, R. R. (1992) An investigation of 3-matrix permutation tests. *Journal of Classification*, **9**, 275–290.
- Oden, N. L., Sokal, R. R., Fortin, M.-J. & Goebl, H. (1993).
 Categorical wombling: detecting regions of significant change in spatially located categorical variables. *Geographical Analysis*, 25, 315–336.
- Ohman, M. E. (1990). The demographic effects of diel vertical migration by zooplankton. *Ecological Monographs*, **60**, 257–282.
- Okabe, A. & Satoh, T. (2009). Spatial analysis on a network. In The SAGE Handbook on Spatial Analysis, eds. A. S. Fotheringham & P. A. Rogers, pp. 443–464. London: SAGE.
- Okabe, A. & Yamada, I. (2001). The K-function method on a network and its computational implementation. Geographical Analysis, 33, 270-290.
- Okabe, A., Boots, B. & Sugihara, K. (1992). Spatial Tesselations: Concepts and Applications of Voronoi Diagrams. New York: Wiley
- Okabe, A., Boots, B., Sugihara, K. & Chiu, S. N. (2000). Spatial tessellations-Concepts and Applications of Voronoi Diagrams, 2nd edn. Chichester: Wiley.
- Okabe, A., Okunuki, K.-I. & Shiode, S. (2006). SANET: A toolbox for spatial analysis on a network. *Geographical Analysis*, 38, 57–66.
- Olson, J. S. (1958). Lake Michigan dune development. 2. Plants as agents and tools in geomorphology. *Journal of Geology*, 66, 345–351.
- OMNR (Ontario Ministry of Natural Resources) (1998).

 Ontario Provincial Landcover Database. Peterborough:
 Ontario Ministry of Natural Resources.
- Opdam, P. & Wascher, D. (2004). Climate change meets habitat fragmentation: linking landscape and biogeographical scale levels in research and conservation. *Biological Conservation*, 117, 285–297.
- Openshaw, S. (1984). *The Modifiable Areal Unit Problem*. Norwich: Geo Books.
- Openshaw S. (1987) An automated geographical analysis system. *Environment and Planning A*, **19**, 431–436.
- Osborne, P. E. & Suarez-Seoane, S. (2002). Should data be partitioned spatially before building large-scale distribution models? *Ecological Modelling*, **157**, 249–259.

- Osborne, P. E., Giles M., Foody, G. M. & Suárez-Seoane, S. (2007). Non-stationarity and local approaches to modelling the distributions of wildlife. *Diversity and Distributions*, **13**, 313–323.
- Ostendorf, B. & Reynolds, J. F. (1998). A model of arctic tundra vegetation derived from topographic gradients. *Landscape Ecology*, **13**, 187–201.
- Otis, D. L. (1997). Analysis of habitat selection studies with multiple patches within cover types. *Journal of Wildlife Management*, 61, 1016–1022.
- Otis, D. L. & White, G.C. (1999). Autocorrelation of location estimates and the analysis of radiotracking data. *Journal of Wildlife Management*, 63, 1039–1044.
- Pacala, S. W. (1997). Dynamics of plant communities. In *Plant Ecology*, 2nd edn, ed. M. C. Crawley, pp. 532–555. Blackwell Scientific. UK.
- Pagel, J. & Schurr, F. M. (2012). Forecasting species ranges by statistical estimation of ecological niches and spatial population dynamics. *Global Ecology and Biogeography*, 21, 293–304.
- Pakeman, R. J. & Small, J. L. (2005). The role of the seed bank, seed rain and the timing of disturbance in gap regeneration. *Journal of Vegetation Science*, 16, 121–130.
- Paley, C. J., Taraskin, S. & Elliott, S. (2007) Temporal and dimensional effects in evolutionary graph theory. *Physical Review Letters*, 98, 098103.
- Palmer, M. W. (1992). The coexistence of species in fractal landscapes. American Naturalist, 139, 375–397.
- Palmer, M. W. (1993). Putting things in even better order: the advantages of canonical correspondence analysis. *Ecology*, 74, 2215–2230.
- Parrott, L., Proulx, R. & Thibert-Plante, X. (2008). Threedimensional metrics for the analysis of spatiotemporal data in ecology. *Ecological Informatics*, 3, 343–353.
- Pascual, M. & Dunne, J. A. eds. (2006). Ecological Networks: Linking Structure to Dynamics in Food Webs. New York: Oxford University Press.
- Patil, G. P. & Taillie, C. (2004). Upper level set scan statistic for detecting arbitrarily shaped hotspots. *Environmental and Ecological Statistics*, 11, 183–197.
- Patterson, B. D. & Atmar, W. (1986). Nested subsets and the structure of insular mammalian faunas and archipelagoes. *Biological Journal of the Linnean Society*, 28, 65–82.
- Pausas, J. G., Bonet, A., Maestre, F. T. & Climent, A. (2006). The role of the perch effect on the nucleation process in Mediterranean semi-arid oldfields. *Acta Oecologia*, 29, 346–352.
- Peet, R. K. (1974). The measurement of species diversity.

 Annual Review of Ecology and Systematics, 5, 285-307.
- Pelletier, B., Fyles, J. W. & Dutilleul, P. (1999). Tree species control and spatial structure of forest floor properties in a mixed-species stand. *Écoscience*, 6, 79–91.

- Penny, D., Foulds, L. R. & Hendy, M. D. (1982). Testing the theory of evolution by comparing phylogenetic trees constructed from five different protein sequences. *Nature*, 297, 197–200.
- Penrose, M. (2003). *Random Geometric Graphs*. Oxford: Oxford University Press.
- Penttinen, A. K., Stoyan, D. & Henttonen, H. M. (1992).
 Marked point processes in forest statistics. Forest Science,
 38, 806–824.
- Peralta, P. & Mather, P. (2000). An analysis of deforestation patterns in the extractive reserves of Acre, Amazonia, from satellite imagery: a landscape ecological approach. *International Journal of Remote Sensing*, 21, 2555–2570.
- Peres-Neto, P. R. (2006). A unified strategy for estimating and controlling spatial, temporal and phylogenetic autocorrelation in ecological models. *Oecologia Brasiliensis*, 10, 105–119.
- Peres-Neto, P. R. & Legendre, P. (2010). Estimating and controlling for spatial structure in the study of ecological communities. Global Ecology and Biogeography, 19, 174-184.
- Perez, S. I., Diniz-Filho, J. A. F., Bernal, V. & Gonzalez, P. N. (2010). Spatial regression techniques for inter-population data: studying the relationships between morphological and environmental variation. *Journal of Evolutionary Biology*, 2, 237–248.
- Perry, J. N. (1994). Chaotic dynamics can generate Taylor's power-law. Proceedings of the Royal Society of London Series B, Biological Sciences, 257, 221–226.
- Perry, J. N. (1995). Spatial analysis of distance indices. *Journal of Animal Ecology*, 64, 303–314.
- Perry, J. N. (1996). Simulating spatial patterns of counts in agriculture and ecology. Computers and Electronics in Agriculture, 15, 93–109.
- Perry, J. N. (1999). Red-blue plots for detecting clusters in count data. *Ecology Letters*, 2, 106-113.
- Perry, J. N., Smith, R. H., Woiwod, I. P. & Morse, D.R. (eds.) (2000). Chaos in Real Data: The Analysis of Non-Linear Dynamics from Short Ecological Time Series, 225pp. Dordrecht: Kluwer Academic.
- Perry, J. N., Woiwod, I. P. & Hanski, I. (1993). Using responsesurface methodology to detect chaos in ecological time series. Oikos, 68, 329-339.
- Peters, V. S. (2003). Keystone processes affect succession in boreal mixed woods: the relationship between masting in white spruce and fire history. Ph.D. Thesis, University of Alberta.
- Peterson, E. E., Ver Hoef, J. M., Isaak, D. J. et al. (2013). Modeling dendritic ecological networks in space: an integrated network perspective. *Ecology Letters*, 16, 707-719.

- Peterson, G. D. (2002). Contagious disturbance, ecological memory, and the emergence of landscape pattern. *Ecosystems*, 5, 329–338.
- Petrovskii, S. V. & Malchow, H. (2001). Wave of chaos: new mechanism of pattern formation in spatio-temporal population dynamics. *Theoretical Population Biology*, 59, 157– 174.
- Pfeifer, P. E. & Deutsch, S. J. (1980). Identification and interpretation of first order space-time ARMA models. *Technometrics*, 22, 397–408.
- Philibert, M. D., Fortin, M.-J. & Csillag, F. (2008). Spatial structure effects on the detection of patches boundaries using local operators. *Environmental and Ecological Statis*tics, 15, 447–467.
- Pielou, D. P. & Pielou, E. C. (1968). The associations of species of infrequent occurrence: the insect and spider fauna of *Polyporus betulinus* (Bulliard) Fires. *Journal of Theoretical Biology*, 21, 202–216.
- Pielou, E. C. (1959). The use of point-to-plant distances in the study of the pattern of plant populations. *Journal of Ecology*, 47, 607–613.
- Pielou, E. C. (1975a). Ecological models on an environmental gradient. In *Applied Statistics*, ed. R. P. Gupta, pp. 261–269. Holland: North-Holland Publishing Company.
- Pielou, E. C. (1975b). *Ecological Diversity*. New York: John Wilev.
- Pielou, E. C. (1977a). The latitudinal spans of seaweed species and their patterns of overlap. *Journal of Biogeography*, 4, 299–311.
- Pielou, E. C. (1977b). *Mathematical Ecology*. New York: Wiley.Pielou, E. C. (1978). Latitudinal overlap of seaweed species: evidence for quasi-sympatric speciation. *Journal of Biogeography*, 5, 227–238.
- Pielou, E. C. (1979). Quick method of determining the diversity of foraminiferal assemblages. *Journal of Paleontology*, 53, 1237-1242.
- Pinhiero, J. C. & Bates, D.M. (2000) Mixed-effects Models in S and S-Plus. New York: Springer.
- Pinto, N. & Keitt, T. H. (2009). Beyond the least-cost path: evaluating corridor redundancy using a graph-theoretic approach. *Landscape Ecology*, 24, 253–266.
- Pitas, I. (2000). Digital Image Processing Algorithms and Applications. New York: Wiley-Interscience.
- Plante, M., Lowell, L., Potvin, F., Boots, B. & Fortin, M.-J. (2004). Studying deer habitat on Anticosti Island, Québec: relating animal occurrences and forest map information. *Ecological Modelling*, 174, 387–399.
- Plant, R. E. (2012). Spatial Data Analysis in Ecology and Agriculture using R. Boca Raton: CRC Press. Taylor & Francis Group.

- Plotkin, J. B., Potts, M. D., Leslie, N. et al. (2000). Species-area curves, spatial aggregation, and habitat specialization in tropical forests. *Journal of Theoretical Biology*, 207, 81–99.
- Plotnick, R. E., Gardner, R. H., Hargrove, W. W., Pretegaard, K. & Perlmutter, M. (1996). Lacunarity analysis: a general technique for the analysis of spatial patterns. *Physical Review E*, 53, 5461–5468.
- Polakowska, A. E., Fortin, M.-J. & Couturier, A. (2012). Quantifying the spatial relationship between bird species' distributions and landscape feature boundaries in southern Ontario, Canada. *Landscape Ecology*, 27, 1481–1493.
- Pollard, J. H. (1971). On distance estimators of density in randomly distributed forests. *Biometrics*, 27, 991-1002.
- Porteus, B. T. (1987). The mutual independence hypothesis for categorical data in complex sampling schemes. *Biometrika*, **74**, 857–862.
- Poulin, R. & Guegan, J. F. (2000). Nestedness, anti-nestedness, and the relationship between prevalence and intensity in ectoparasite assemblages of marine fish: a spatial model of species coexistence. *International Journal for Parasitology*, 30, 1147-1152.
- Proulx, S. R., Promislow, D. E. L. & Phillip, P.C. (2005). Network thinking in ecology and evolution. *Trends in Ecology and Evolution*, 20, 345–352.
- Qi, Y. & Wu, J. (1996). Effects of changing scale on the results of landscape pattern analysis using spatial autocorrelation indices. *Landscape Ecology*, 11, 39–50.
- Rahbek, C. (2005). The role of spatial scale and the perception of large-scale species-richness patterns. *Ecology Letters*, 8, 224–239.
- Rangel, T. F., Diniz, J. A. F. & Bini, L.M. (2010). SAM: a comprehensive application for Spatial Analysis in Macroecology. *Ecography*, 33, 46–50.
- Ranta, E., Kaitala, V., Lindström, J. & Helle, E. (1997). The Moran effect and synchrony in population dynamics. *Oikos*, 78, 136–142.
- Rao, C. R. (1964). The use and interpretation of principal component analysis in applied research. Sankhyaá, Ser., A26, 329-358.
- Rayfield, B., Fortin, M.-J. & Fall, A. (2010). The sensitivity of least-cost habitat graphs to relative cost surface values. *Landscape Ecology*, 25, 519–532.
- Rayfield, B., Fortin, M.-J. & Fall, A. (2011). Connectivity for conservation: a framework to classify habitat network statistics. *Ecology*, 92, 847–858.
- Read S., Bath, P., Willett, P. & Maheswaran, R. (2011).
 Measuring the spatial accuracy of the spatial scan statistic. Spatial and Spatio-temporal Epidemiology, 2, 69–78.
- Redding, T. E., Hope, G. D., Fortin, M.-J., Schmidt, M. G. & Bailey, W.G. (2003). Spatial patterns of soil temperature and

- moisture across subalpine forest-clearcut edges in the southern interior of British Columbia. *Canadian Journal of Soil Science*, **83**, 121–130.
- Reich, R. M., Metzger, K. L. & Bonham, C.D. (1997). Application of permutation procedures for comparing multispecies point patterns of grassland plants. *Grassland Sci*ence, 43, 189–195.
- Remmel, T. K. & Csillag, F. (2003). When are two landscape pattern indices significantly different? *Journal of Geographical Systems*, 5, 331–351.
- Remmel, T. K. & Fortin, M.-J. (2013). Categorical, classfocused map patterns: Characterization and comparison. *Landscape Ecology*, 28, 1587–1599.
- Remmert, H. (1991). The Mosaic-cycle Concept of Ecosystems. Berlin: Spring-Verlag.
- Renshaw, E. & Ford, E. D. (1984). The description of spatial pattern using two-dimensional spectral analysis. *Vegetatio*, 56, 75–85.
- Richardson, M. C., Fortin, M.-J. & Branfireun, B.A. (2009). Hydrogeomorphic edge detection and delineation of land-scape functional units from LiDAR digital elevation models. Water Resources Research, 45, W10441, doi:10.1029/2008WR007518.
- Ricotta, C. (2005). Additive partitioning of Rao's quadratic diversity: a hierarchical approach. *Ecological Modelling*, 183, 365–371.
- Ricotta, C. (2006). Towards a complex, plural and dynamic approach to diversity: rejoinder to Myers and Patil, Podani, and Sarkar. Acta Biotheoretica, 54, 141-146.
- Ricotta, C. & Anand, M. (2006). Spatial complexity of ecological communities: Bridging the gap between probabilistic and non-probabilistic uncertainty measures. *Ecological Modelling*, 197, 59–66.
- Ricotta, C. & Marignani, M. (2007). Computing beta-diversity with Rao's quadratic entropy: a change of perspective. *Diversity and Distributions*, 13, 237-241.
- Rietkerk, M., Dekker, S. C., Wassen, M. J., Verkroost, A. W. M. & Bierkens, M.F.P. (2004). A putative mechanism for bog patterning. *The American Naturalist*, **163**, 700–708.
- Riitters, K. H., & Coulston, J. W. (2005). Hot spots of perforated forest in the eastern United States. *Environmental Manage*ment, 35, 483–492.
- Ripley, B. D. (1976). The second-order analysis of stationary point processes. *Journal of Applied Probability*, 13, 255–266.
- Ripley, B. D. (1978). Spectral analysis and the analysis of pattern in plant communities. *Journal of Ecology*, 66, 965– 981.
- Ripley, B. D. (1981). Spatial Processes. New York: Wiley.
- Ripley, B. D. (1988). Statistical Inference for Spatial Processes. Cambridge: Cambridge University Press.

- Robertson, C., Nelson, T. A., Boots, B. & Wulder, M.A. (2007).
 STAMP: Spatial-temporal analysis of moving polygons.
 Journal of Geographical Systems, 9, 207–227.
- Rooney, S. M., Wolfe, A. & Hayden, T.J. (1998). Autocorrelated data in telemetry studies: time to independence and the problem of behavioural effects. *Mammal Review*, 29, 89–98.
- Rosenberg, M. S. & Anderson, C. D. (2011). PASSaGE: Pattern Analysis, Spatial Statistics and Geographic Exegesis. Version 2. Methods in Ecology and Evolution, 2, 229–232.
- Rosenzweig, M. L. (1995). Species Diversity in Space and Time. Cambridge: Cambridge University Press.
- Rossi, R. E., Mulla, D. J., Journel, A. J. & Franz, E.H. (1992). Geostatistical tools for modeling and interpreting ecological spatial dependence. *Ecological Monographs*, 62, 277–314.
- Rousset, F. (1996). Equilibrium values of measures of population subdivision for stepwise mutation processes. *Genetics*, 142, 1357–1362.
- Rozenfeld, A. F., Arnaud-Haond S., Hernández-García, E. et al. (2008). Network analysis identifies weak and strong links in a metapopulation system. Proceedings of the National Academy of Sciences, 105, 18 824–18 829.
- Ruxton, G. D. (1993). Linked populations can still be chaotic. *Oikos*, **68**, 347–348.
- Sadahiro, Y. & Umemura, M. (2002). A computational approach for the analysis of changes in polygon distributions. *Journal of Geographical Systems*, 3, 137–154.
- Salvatori, V., Skidmore, A. K., Corsi, F. & Van der Meer, F. (1999). Estimating tenporal independence of radiotelemetry data on animal activity. *Journal of Theoretical Biology*, 198, 567–574.
- Sanuy, D. & Bovet, P. (1997). A comparative study on the paths of five anura species. *Behavioural Processes*, 41, 193–199.
- Sarkar, A. (2009). An Introduction to Biodiversity Informatics. New York: World Scientific Publishing.
- Saunders, S. C., Brosofske, K. D., Chen, J., Drummer, T. D. & Gustafson E. J. (2005). Identifying scales of pattern in ecological data: a comparison of lacunarity, spectral and wavelet analyses. *Ecological Complexity*, 2, 87–105.
- Schabenberger, O. & Gotway, C. A. (2005). Statistical Methods for Spatial Data Analysis. London: Chapman and Hall.
- Scheiner, S. M., Cox, S. B., Willig, M. et al. (2000). Species richness, species-area curves and Simpson's paradox. Evolutionary Ecology Research, 2, 791–802.
- Schladitz, K. & Baddeley, A. J. (2000). A third order point process characteristic. Scandinavian Journal of Statistics, 27, 657-671.
- Schroeder, B. & Seppelt, R. (2006). Analysis of pattern-process interactions based on landscape models-Overview, general concepts, and methodological issues. *Ecological Modelling*, 199, 505-516.

- Schroeder, M. (1991). Fractals, Chaos, Power Laws. New York: Freeman.
- Schultz, C. B. & Crone, E.E. (2001). Edge-mediated dispersal behavior in a prairie butterfly. *Ecology*, **82**, 1879–1892.
- Schulz, J. P. (1960). Ecological Studies on Rain Forest in Northern Surinam. Verhandelingen der Koninklijke Nederlandse Akademie van Wetenschappen, Afdeling Natuurkunde.
- Scrosati, R. & Heaven, C. (2007). Spatial trends in community richness, diversity and evenness across rocky intertidal environmental stress gradients in eastern Canada. *Marine Ecogy Progress Series*, 342, 1–14.
- Selmi, S. & Boulinier, T. (2001). Ecological biogeography of Southern Ocean islands: the importance of considering spatial issues. *American Naturalist*, **158**, 426-437.
- Setzer, R. W. (1985). Spatio-temporal patterns of mortality in Pemphigus populicaulis and P. populitransversus on cottonwoods. Oecologia, 67, 310-321.
- Sherratt, J. A. (2001). Periodic travelling waves in cyclic predator-prey systems. *Ecology Letters*, 4, 30-37.
- Sherratt, T. N., Lambin, X., Petty, S. J. et al. (2000). Use of coupled oscillator models to understand synchrony and traveling waves in populations of the field vole Microtus agrestis in northern England. Journal of Applied Ecology, 37, 148–158.
- Shih, F. Y. (2009). Image Processing and Mathematical Morphology: Fundamentals and Applications. Boca Raton: CRC Press.
- Shimatani, K. (2001). Multivariate point processes and spatial variation of species diversity. Forest Ecology and Management, 142, 215–229.
- Shimatani, K. & Kubota, Y. (2004). Spatial analysis for continuously changing point patterns along a gradient and its application to an Abies sachalinensis population. Ecological Modelling, 180, 359–369.
- Shipley, B. & Keddy, P. R. (1987). The individualistic and community-unit concepts as falsifiable hypotheses. *Vegeta*tio, 69, 47-55.
- Silvertown, J., Dodd, M., Gowing, D., Lawson, C. & McConaway, K. (2006). Phylogeny and the hierarchical organization of plant diversity. *Ecology*, 87, S39–S49.
- Simard, Y., Marcotte, D. & Naraghi, K. (2003). Three-dimensional acoustic mapping and simulation of krill distribution in the Saguenay-St. Lawrence Marine Park whale feeding ground. *Aquatic Living Resources*, 16, 137-144.
- Simberloff, D. (1979). Nearest neighbor assessments of spatial configurations of circles rather than points. *Ecology*, 60, 679–685.

- Simberloff, D. & Martin, J. L. (1991). Nestedness of insular avifaunas-Simple summary statistics masking complex species patterns. *Ornis Fennica*, 68, 178–192.
- Smith, R. L. & Robinson, P. J. (1995). A Bayesian approach to modelling spatial-temporal precipitation data. *Proceedings* of the Sixth International Meeting on Statistical Climatology, Galway, Ireland.
- Smouse, P. E., Long, J. C. & Sokal, R.R. (1986). Multiple regression and correlation extensions of the Mantel test of matrix correspondence. *Systematic Zoology*, 35, 627–632.
- Soininen, J., McDonald, R., & Hillibrand, H. (2007). The distance decay of similarity in ecological communities. *Ecography* 30, 3–12.
- Sokal, R. R. & Oden, N.L. (1978). Spatial autocorrelation in biology. 1. Methodology. Biological Journal of the Linnean Society, 10, 199–228.
- Sokal, R. R. & Rohlf, F.J. (1981). *Biometry*, 2nd edn. San Francisco: Freeman.
- Sokal, R. R. & Rohlf, F.J. (1995). Biometry, 3rd edn. San Francisco: Freeman.
- Sokal, R. R. & Wartenberg, D.E. (1983). A test of spatial autocorrelation using an isolation-by-distance model. *Genetics*, 105, 219–237.
- Sokal, R. R., Oden, N. L. & Thomson, B.A. (1988). Genetic changes across language boundaries in Europe. *American Journal of Physical Anthropology*, 76, 337–361.
- Sokal, R. R., Oden, N. L. & Thomson, B.A. (1998). Local spatial autocorrelation in biological variables. *Biological Journal of* the Linnean Society, 65, 41–62.
- Sokal, R. R., Oden, N. L., Thomson, B. A. & Kim, J. (1993). Testing for regional differences in means: distinguishing inherent from spurious spatial autocorrelation by restricted randomization. *Geographical Analysis*, 25, 199–210.
- Solé, R. (2007). Scaling laws in the drier. *Nature*, 449, 151–153.
 Solé, R. & Bascompte, J. (2006). *Self-Organization in Complex Ecosystems*. Princeton: Princeton University Press.
- Solé, R. V. & Bascompte, J. (1995). Measuring chaos from spatial information. *Journal of Theoretical Biology*, 175, 139–147.
- Solow, A. R. (1989). Bootstrapping sparsely sampled spatial point patterns. *Ecology*, 70, 379–382.
- Spear, S. Balkenhol N., Fortin M.-J., McRae B. & Scribner K. (2010). Use of resistance surfaces for landscape genetic studies: considerations for parameterization and analysis. *Molecular Ecology*, 19, 3576–3591.
- Spooner, P. G., Lunt, I. D., Okabe, A. & Shiode, S. (2004). Spatial analysis of roadside *Acacia* populations on a road network using the network *K*-function. *Landscape Ecology*, 19, 491–499.

- Sprugel, D. G. (1976). Dynamic structure of wave-regenerated Abies balsamea forests in the northeastern United States. Journal of Ecology, 64, 889–911.
- St.-Louis, V., Fortin, M.-J. & Desrochers, A. (2004). Association between microhabitat and territory boundaries of two forest songbirds. *Landscape Ecology*, 19, 591–601.
- Stohlgren, T. J. (2007). Measuring Plant Diversity: Lessons from the Field. New York: Oxford University Press.
- Stone, L. & Ezrati, S. (1996). Chaos, cycles and spatiotemporal dynamics in plant ecology. *Journal of Ecology*, 84, 279–291.
- Stoyan, D. & Penttinen, A. (2000). Recent applications of point process methods in forestry statistics. *Statistical Science*, 15, 61–78.
- Stoyan, D., Kendall, W. S., & Mecke, J. (1995). Stochastic Geometry and its Applications, 2nd edn. Chichester: Wiley.
- Strogatz, S. H. (2001). Exploring complex networks. *Nature*, 410, 268–276.
- Sutcliffe, O. L., Thomas, C. D. & Moss, D. (1996). Spatial synchrony and asynchrony in butterfly population dynamics. *Journal of Animal Ecology*, 65, 85–95.
- Suzuki, S. N., Kachi, N. & Suzuki, J.-I. (2008). Development of local size hierarchy causes regular spacing of trees in an even-aged *Abies* forest: analyses using spatial autocorrelation and the mark correlation function. *Annals of Botany*, 102, 435–441.
- Swihart, R. K. & Slade, N.A. (1985). Testing for independence of observations in animal movements. *Ecology*, 66, 1176– 1184
- Swihart, R. K. & Slade, N.A. (1986). The importance of statistical power when testing for independence in animal movements. *Ecology*, 67, 255–258.
- Takahashi, K., Kulldorff, M., Tango, T. & Yih, K (2008). A flexibly shaped space-time scan statistic for disease outbreak detection and monitoring. *International Journal of Health Geographics* 7:14. doi:10.1186/1476-072X-7-14.
- Tang, J. Musolesi, M., Mascolo, C. & Latora, V. (2009). Temporal Distance Metrics for Social Network Analysis. Proceedings of the 2nd ACM SIGCOMM workshop on Online Social Networks. WOSN'09. http://doi.acm.org/10.1145/1592665.1592674.
- Tango, T. (2010). Statistical Methods for Disease Clustering. New York: Springer.
- Tavaré, S. (1983). Serial dependence in contingency tables. Journal of the Royal Statistical Society B, 45, 100-106.
- Tavaré, S. & Altham, P. M. E. (1983). Serial dependence of observations leading to contingency tables, and corrections to chi-squared statistics. *Biometrika*, 70, 139-144.

- Taylor, P. D., Day, T. & Wild, G. (2007). Evolution of cooperation in a finite homogeneous graph. *Nature*, 447, 469-742.
- ter Braak, C. J. F. (1986). Canonical correspondence analysis: a new eigenvector technique for multivariate direct gradient analysis. *Ecology*, 67, 1167–1179.
- Thiery, J. M., Dherbes, J. M., & Valentin, C. (1995). A model simulation the genesis of banded vegetation patterns in Niger. *Journal of Ecology*, 83, 497–507.
- Thomas, D. L. & Taylor, E.J. (1990). Study designs and tests for comparing resource use and availability. *Journal of Wildlife Management*, 54, 322-330.
- Tischendorf, L. (2001). Can landscape indices predict ecological processes consistently? *Landscape Ecology*, 16, 235–54.
- Tobin, P. C. & Bjørnstad, O. N. (2003). Spatial dynamics and cross-correlation in a transient predator-prey system. *Jour*nal of Animal Ecology, 72, 460–467.
- Tobler, W. F. (1970). A computer movie simulating urban growth in the Detroit region. *Economic Geography*, 46, 234–240.
- Todd, K. W., Csillag, F. & Atkinson, P.M. (2003). Three-dimensional mapping of light transmittance and foliage distribution using lidar. *Canadian Journal of Remote Sensing*, 29, 544–555.
- Torrence, C. & Compo G. P. (1998). A practical guide to wavelet analysis. *Bulletin of the American Meteorological* Society, 79, 61–78.
- Toussaint, G. T. (1980). The relative neighbourhood graph of a finite planar set. *Pattern Recognition*, **12**, 261–268.
- Tuia, D., Ratle, F., Lasaponara, R., Tleesca, L & Kanevski, M. (2008). Scan statistics analysis of forest fire clusters. Communications in Nonlinear Science and Numerical Simulation, 13, 1689–1694.
- Tuomisto, H. (2010a). A diversity of beta diversities: straightening up a concept gone awry. Part 1. Defining beta diversity as a function of alpha and gamma diversity. *Eco-graphy*, 33, 2–22.
- Tuomisto, H. (2010b). A diversity of beta diversities: straightening up a concept gone awry. Part 2. Quantifying beta diversity and related phenomena. *Ecography*, 33, 23-45.
- Tuomisto, H. & Ruokolainen, K. (2006). Analyzing or explaining beta diversity? Understanding the targets of different methods of analysis. *Ecology*, 87, 2697–2708.
- Turchin, P. (1996). Fractal analyses of animal movement: a critique. Ecology, 77, 2086–2090.
- Turchin, P. (1998). Quantitative Analysis of Movement. Sunderland: Sinauer.

- Turchin, P. & Taylor, A. D. (1992). Complex dynamics in ecological time series. *Ecology*, 73, 289–305.
- Turner, M. G., Gardner, R. H. & O'Neill, R. V. (2003). Landscape Ecology in Theory and Practice: Pattern and Process. New York: Springer-Verlag.
- Ulam, S. (1962). Patterns of growth of figures: mathematical aspects. In: Proceedings of Symposia in Applied Mathematics, XIV: Mathematical Problems in the Biological Sciences. American Mathematical Society, 61-65.
- Ulrich, W., Almeida-Neto, M. & Gotelli, N.J. (2008).
 A consumer's guide to nestedness. Oikos 118, 3–17.
- Underwood, A. J. (1978a). The detection of nonrandom patterns of distribution of species along a gradient. *Oecologia* (Berl.), 36, 317-326.
- Underwood, A. J. (1978b). A refutation of critical tide levels as determinants of the structure of the intertidal communities on British shores. *Oecologia (Berl.)*, 37, 261–276.
- Upton, G. J. G. & Fingleton, B. (1985). Spatial Data Analysis by Example, vol. I: Point Pattern and Quantitative Data. New York: Wiley.
- Upton, G. J. G. & Fingleton, B. (1989). Spatial Data Analysis by Example, vol. II: Categorical and Directional Data. New York: Wiley.
- Urban, D. L. & Keitt, T. (2001). Landscape connectivity: a graph-theoretic perspective. *Ecology*, 82, 1205–1218.
- Urban, D. L., Minor, E. S., Treml, E. A. & Schick, R.S. (2009).
 Graph models of habitat mosaics. *Ecology Letters*, 12, 260–273.
- Vaček, S. & Lepš, J. (1996). Spatial dynamics of forest decline: the role of neighbouring trees. *Journal of Vegetation Science*, 7, 789–798.
- Valcu, M. & Kempenaers, B. (2010). Is spatial autocorrelation an intrinsic property of territory size? *Oecologia*, 162, 609–615.
- van Es, H. M. & van Es, C. L. (1993). The spatial nature of randomization and its effects on the outcome of field experiments. *Agronomy Journal*, 85, 420-428.
- van Langevelde, F., van der Knaap, W. G. M. & Claassen, G.D.H. (1998). Comparing connectivity in landscape networks. Environmental Planning B: Planning and Design, 25, 849–863. Not seen, author?
- van Lieshout, M. N. M. & Baddeley, A.J. (1999). Indices of dependence between types in multivariate point patterns. *Scandinavian Journal of Statistics*, 26, 511–532.
- Van Strien, M. J., Keller, D. & Holderegger, R. (2012). A new analytical approach to landscape genetic modelling: leastcost transect analysis and linear mixed models. *Molecular Ecology*, 21, 4010–4023.
- Vekemans, X. & Hardy, O. J. (2004). New insights from finescale spatial genetic structure analyses in plant populations. *Molecular Ecology*, 13, 921–935.

- Velland, M. (2001). Do commonly used indices of β-diversity measure species turnover? *Journal of Vegetation Science*, 12, 545–552.
- Venables, W. N. & Ripley, B.D. (2002). *Modern Applied Statistics with S*, 4th edn. New York: Springer-Verlag.
- Vepakomma, U. & Fortin, M.-J. (2010). Scale-specific effects of disturbance and environment on vegetation patterns of a boreal landscape. *Proceedings of SilviLaser* 2010.
- Vepakomma, U., Kneeshaw, D. & Fortin, M.-J. (2012). Spatial contiguity and continuity of disturbance in boreal forests: gap persistence, expansion, shrinkage and displacement. *Journal of Ecology*, **100**, 1257–1268.
- Ver Hoef, J. M. & Glenn-Lewin, D. C. (1989). Multiscale ordination: a method for detecting pattern at several scales. Vegetatio, 82, 59-67.
- Ver Hoef, J. M., Peterson, E. E. & Theobald, D. (2006). Spatial statistical models that use flow and stream distance. *Envir*onmental and Ecological Statistics, 13, 449–464.
- Viljugrein, H., Lingjærde, O. C., Stenseth, N. C. & Boyce, M.S. (2001). Spatio-temporal patterns of mink and muskrat in Canada during a quarter century. *Journal of Animal Ecol*ogy, 70, 671–682.
- Volford, A., Izsák, F., Ripszám, M. & Lagzi, I. (2007). Pattern formation and self-organization in a simple precipitation system. *Langmuir*, 23, 961–994.
- Wackernagel, H. (2010). Multivariate Geostatistics, 3rd edn. New York: Springer, 403 pages.
- Wagner, H. H. (2003). Spatial covariance in plant communities: integrating ordination, geostatistics, and variance testing. *Ecology*, 84, 1045–1057.
- Wagner, H. H. (2004). Direct multi-scale ordination with canonical correspondence analysis. *Ecology*, 85, 342–351.
- Wagner, H. H. & Fortin, M.-J. (2005). Spatial analysis of landscapes: concepts and statistics. *Ecology*, 86, 1975–1987.
- Wagner H. H., Holderegger, R., Werth, S. et al. (2005). Variogram analysis of the spatial genetic structure of continuous populations using multilocus microsatellite data. Genetics, 169, 1739–1752.
- Wakefield, J. C., Kelsall, J. E. & Morris, S.E. (2000). Clustering, cluster detection, and spatial variation in risk. In *Spatial Epidemiology: Methods and Applications*, eds. P. Elliott, J. C. Wakefield, N. G. Best & D. J. Briggs, pp. 128–152. Oxford: Oxford University Press.
- Wallerman, J., Joyce, S., Vencatasawmy, C. P. & Olsson, H. (2002). Prediction of forest stem volume using kriging adapted to detected edges. *Canadian Journal of Forest Research*, 32, 509–518.
- Wang, Y.-H., Yang, K.-C., Bridgman, C. L. & Lin, L.-K. (2008).
 Habitat suitability modeling to correlate gene flow with landscape connectivity. *Landscape Ecology*, 23, 989–1000.

- Wartenberg, D. (1985). Multivariate spatial autocorrelation: a method for explanatory geographical analysis. *Geographical Analysis*, 17, 263–283.
- Wasserman, L. (2004). All of Statistics. New York: Springer.
- Watt, A. S. (1947). Pattern and process in the plant community. *Journal of Ecology*, 35, 1–22.
- Watts, D. J. (1999). Small Worlds: The Dynamics of Networks Between Order and Randomness. Princeton: Princeton University Press.
- Watts, D. J. & Strongtz, S.H. (1998). Collective dynamics of 'small-world' networks. *Nature*, 393 (6684), 440–442.
- Weber, K. T., Burcham, M. & Marcum, C.L. (2001). Assessing independence of animal locations with association matrices. *Journal of Rangeland Management*, 54, 21–24.
- Webster, R. (1973). Automatic soil-boundary location from transect data. *Mathematical Geology*, 5, 27–37.
- Webster, R. & Oliver, M. A. (2001). Geostatistics for Environmental Scientists. Chichester: Wiley.
- Webster, R. & Oliver, M. A. (2007). Geostatistics for Environmental Scientists. Wiley. 330 pages.
- Webster's (1989). Webster's Encyclopaedic Unabridged Dictionary of the English Language. New Jersey: Gramercy Books.
- Weishampel, J. F., Godin, J. R. & Henebry, G.M. (2001). Pantropical dynamics of "intact" rain forest canopy structure. Global Ecology and Biogeography, 10, 389–397.
- West, D. B. (2001). *Introduction to Graph Theory*, 2nd edn. Upper Saddle River, NJ: Prentice Hall.
- Westoby, M. (2006). Phylogenetic ecology at world scale, a new fusion between ecology and evolution. *Ecology*, 87, S163–S166.
- White, E. P., Morgan Ernset, S.K., Adler, P. B., Hurlberts, A. H. & Lyons, S.K. (2011). Integrating spatial and temporal approaches to understanding species richness. *Philosophical Transactions of the Royal Society B—Biological Sciences*, 365, 3633–3643.
- White, G. C. & Garrott, R.A. (1990). *Analysis of Wildlife Radio-Tracking Data*. New York: Academic Press.
- Whittaker, R. H. (1975). Communities and Ecosystems, 2nd edn. New York: MacMillan.
- Whittaker, R. H. (1977). Evolution of species diversity in land communities. Evolutionary Biology, 10, 1-67.
- Whittle, P. (1954). On stationary processes in the plane. *Biometrika*, 41, 434–449.
- Wiens, J. A. (1989). Spatial scaling in ecology. Functional Ecology, 3, 385–397.
- Wiens, J. A., Crist, T. O. & Milne, B.T. (1993). On quantifying insect movements. *Environmental Entomology*, 22, 709-715.
- Wikle, C. K. (2003). Hierarchical Bayesian models for predicting the spread of ecological processes. *Ecology*, 84, 1382–1394.

- Wilbur, H. M. (1997). Experimental ecology of food webs: complex systems in temporary ponds. *Ecology*, 78, 2279–2302.
- Williams, B. (1992). The measurement of sinuosity in correlated random walks. *Journal of Theoretical Biology*, 155, 437–442.
- Wilson, J. B. (1988). The effect of initial advantage on the course of plant competition. *Oikos*, 51, 19–24.
- Wilson, J. B. & Agnew, A.D.Q. (1992). Positive-feedback switches in plant communities. Advances in Ecological Research, 23, 263–336.
- Wilson, M. V. & Mohler, C.L. (1983). Measuring compositional change along gradients. *Vegetatio*, 54, 129–141.
- Wilson, M. V. & Shmida, A. (1984). Measuring beta diversity with presence-absence data. *Journal of Ecology*, 72, 1055– 1064.
- Windle, M. J. S., Rose, G. A., Devillers, R. & Fortin, M.-J. (2010). Exploring spatial non-stationarity of fisheries survey data using geographically weighted regression (GWR): an example from the Northwest Atlantic. *ICES Journal of Marine Science*, 67, 145–54.
- Wolfram, S. (2002). A New Kind of Science. Champagne: Wolfram Media.
- Womble, W. H. (1951). Differential systematics. Science, 114, 315–322.
- Woodcock, C. & Strahler, A. (1987). The factor of scale in remote sensing. Remote Sensing of Environment, 21, 311–322.
- Woollons, R. C. (1998). Even-aged stand mortality estimation through a two-step regression process. Forest Ecology and Management, 105, 189–195.
- Wright, D. H. & Reeves, J.H. (1992). On the meaning and measurement of nestedness of species assemblages. *Oecologia*, 92, 416–428.
- Wright, F. (1990). The effective number of codons used in a gene. *Gene*, **87**, 23–29.
- Wright, S. (1931). Statistical theory of evolution. *Journal of the American Statistical Association*, 26, 201–208.
- Wu, J. (2004). Effects of changing scale on landscape pattern analysis: scaling relations. *Landscape Ecology*, 19, 125–138.
- Wu, J. & Qi, Y. (2000). Dealing with scale in landscape analysis: an overview. Geographic Information Sciences, 6, 1–5.
- Wu, J. G., Shen, W. J., Sun, W. Z. & Tueller, P.T. (2002). Empirical patterns of the effects of changing scale on landscape metrics. *Landscape Ecology*, 17, 761–782.
- Wu, X. B., Thuro, T. L. & Whisenant, S.G. (2000). Fragmentation and changes in hydrologic function of tiger bush landscapes, south-west Niger. *Journal of Ecology*, 88, 790–800.
- Wulder, M. A. & Boots, B. (1998). Local spatial autocorrelation characteristics of remotely sensed imagery assessed with the Getis statistic. *International Journal of Remote Sensing*, 19, 2223–2331.

- Wulder, M. A., White, J. C., Coops, N. C., Nelson, T. & Boots, B. (2007). Using local spatial autocorrelation to compare outputs from a forest growth model. *Ecological Modelling*, 209, 264–276.
- Wulff, B. L. & Webb, K.L. (1969). Intertidal zonation of marine algae at Gloucester Point, Virginia. *Chesapeake Science*, 10, 29–35.
- Yarranton, G. A. (1966). A plotless method of sampling vegetation. *Journal of Ecology*, 59, 224–237.
- Yarranton, G. A. & Morrison, R.G. (1974). Spatial dynamics of primary succession: nucleation. *Journal of Ecology*, 62, 417–428.

- Young, C. G., Dale, M. R. T. & Henry, G.H.R. (1999). Spatial pattern of vegetation in high Arctic sedge meadows. *Écoscience*, **6**, 556-564.
- Zadeh, L. A. (1965). Fuzzy sets. Information and Control, 8, 338–353.
- Zapala, M. A. & Schork, N.J. (2006). Multivariate regression analysis of distance matrices for testing associations between gene expression patterns and related variables. Proceedings of the National Academy of Sciences of the United States of America, 103, 19 430-19 435.
- Zar, J. H. (1984). Biostatistical Analysis, 2nd edn. Englewood Cliffs, NJ: Prentice-Hall.