

# STA2201-lab1

Kerry Hu

## # Lab Exercises

1. Plot the ratio of male to female mortality rates over time for ages 10,20,30 and 40 (different color for each age) and change the theme.

```
library(tidyverse)
```

```
-- Attaching packages ----- tidyverse 1.3.2 --
v ggplot2 3.4.0      v purrr   0.3.4
v tibble  3.1.8      v dplyr  1.0.10
v tidyr   1.2.1      v stringr 1.5.0
v readr   2.1.3      v forcats 0.5.2
-- Conflicts ----- tidyverse_conflicts() --
x dplyr::filter() masks stats::filter()
x dplyr::lag()     masks stats::lag()
```

```
dm <- read_table("https://www.prdh.umontreal.ca/BDLC/data/ont/Mx_1x1.txt", skip = 2, col_t
```


Warning: 494 parsing failures.

row	col	expected	actual
108	Female	no trailing characters	. 'https://www.prdh.umontreal.ca/BDLC/data/ont/Mx_1x1
109	Female	no trailing characters	. 'https://www.prdh.umontreal.ca/BDLC/data/ont/Mx_1x1
110	Female	no trailing characters	. 'https://www.prdh.umontreal.ca/BDLC/data/ont/Mx_1x1
110	Male	no trailing characters	. 'https://www.prdh.umontreal.ca/BDLC/data/ont/Mx_1x1
110	Total	no trailing characters	. 'https://www.prdh.umontreal.ca/BDLC/data/ont/Mx_1x1
...	...	...	...

See problems(...) for more details.

```
dp <- dm |>
  filter(Age==10|Age==20|Age==30|Age==40)|>
  mutate(mf_ratio = Male/Female)|>select(Year,Age,mf_ratio)
dp|> ggplot(aes(x=Year,y=mf_ratio,color = Age))+geom_line()+labs(title = "10,20,30,40 ye
```

10,20,30,40 year old MF mortality ratio over time



- Find the age that has the highest female mortality rate each year.

```
summary_max <- dm |> group_by(Year)|>select(Year,Age,Female)|>
  summarize(max_mortality = max(Female, na.rm = TRUE))
df_list1 = list(dm,summary_max)
dmf<- df_list1 |> reduce(left_join, by='Year') |>select(Year,Age,Female,max_mortality)
dmf <- dmf |>
  mutate_at(3, ~replace_na(.,0))
Year=rep(0,200)
Age=rep(0,200)
YearAge1<-data.frame(Year,Age)

j=1
for(i in 1:10989){
  if(dmf$Female[i]==dmf$max_mortality[i]) {YearAge1$Age[j]<-dmf$Age[i];YearAge1$Year[j]<-d
  }
}
```

```
YearAge<-head(YearAge1,102)  
YearAge
```

	Year	Age
1	1921	106
2	1922	98
3	1923	104
4	1924	107
5	1925	98
6	1926	106
7	1927	106
8	1928	104
9	1929	104
10	1930	105
11	1931	104
12	1932	105
13	1933	104
14	1934	106
15	1935	104
16	1936	106
17	1937	105
18	1938	104
19	1939	105
20	1940	104
21	1941	105
22	1942	104
23	1943	105
24	1944	98
25	1945	104
26	1946	105
27	1947	104
28	1948	99
29	1949	102
30	1950	102
31	1951	110+
32	1952	107
33	1953	106
34	1954	110+
35	1955	107
36	1956	110+
37	1957	107
38	1958	110+

39	1959	108
40	1960	107
41	1961	106
42	1962	108
43	1963	109
44	1964	109
45	1965	109
46	1966	105
47	1967	107
48	1968	97
49	1969	109
50	1970	107
51	1971	107
52	1972	107
53	1973	105
54	1974	107
55	1974	109
56	1975	108
57	1975	110+
58	1976	106
59	1976	108
60	1977	103
61	1978	109
62	1979	109
63	1980	110+
64	1981	107
65	1982	109
66	1983	110+
67	1984	110+
68	1985	110+
69	1986	109
70	1987	109
71	1988	108
72	1989	108
73	1990	108
74	1991	103
75	1992	108
76	1993	109
77	1994	109
78	1995	107
79	1996	109
80	1997	107
81	1998	110+

```

82 1999 110+
83 2000 106
84 2001 110+
85 2002 107
86 2003 109
87 2004 108
88 2005 108
89 2006 110+
90 2007 107
91 2008 109
92 2009 110+
93 2010 108
94 2011 110+
95 2012 109
96 2013 110+
97 2014 110+
98 2015 110+
99 2016 110+
100 2017 110+
101 2018 110+
102 2019 110+

```

3. Use the ‘`summarize(across())`’ syntax to calculate the standard deviation of mortality rates by age for the Male, Female and Total populations.

```

summary_SD <- dm |> group_by(Age)|>
  summarize(across(Female:Total, sd))
summary_SD

```

```

# A tibble: 111 x 4
  Age      Female      Male      Total
<chr>    <dbl>    <dbl>    <dbl>
1 0      0.0256    0.0330    0.0294
2 1      0.00352  0.00396  0.00374
3 10     0.000474  0.000561  0.000509
4 100    0.0928    0.138     0.0729
5 101    0.125     0.158     0.0995
6 102    0.143     0.214     0.114
7 103    0.252     0.371     0.208
8 104    0.449     NA         0.363
9 105    NA        NA         NA
10 106    NA        NA         NA

```

```
# ... with 101 more rows
```

4. The Canadian HMD also provides population sizes over time (<https://www.prhd.umontreal.ca/BDLC/data/ont>). Use these to calculate the population weighted average mortality rate separately for males and females, for every year. Make a nice line plot showing the result (with meaningful labels/titles) and briefly comment on what you see (1 sentence). Hint: 'left\_join' will probably be useful here.

```
df <- read_table("https://www.prhd.umontreal.ca/BDLC/data/ont/Population.txt", skip = 2, col_types = "d")
df_list1 = list(dm, df)
dmf <- df_list1 |> reduce(left_join, by=c('Year', 'Age'))
dmf <- dmf |> mutate(weighed_F = Female.x * Female.y, weighed_M = Male.x * Male.y) |> select(-Total.x, -Total.y)
dmfsum <- dmf |> group_by(Year) |> summarize(total_F = sum(Female.y, na.rm = TRUE), total_M = sum(Male.y, na.rm = TRUE))
df_list2 = list(dmf, dmfsum)
dmf <- df_list2 |> reduce(left_join, by='Year')
dmf
```

```
# A tibble: 10,989 x 10
```

	Year	Age	Female.x	Male.x	Female.y	Male.y	weighed_F	weighed_M	total_F	total_M
	<dbl>	<chr>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>
1	1921	0	0.0978	0.129	30157.	31530.	2948.	4070.	1.46e6	1.48e6
2	1921	1	0.0129	0.0144	30391.	31319.	394.	452.	1.46e6	1.48e6
3	1921	2	0.00521	0.00737	30962.	31785.	161.	234.	1.46e6	1.48e6
4	1921	3	0.00471	0.00457	31306.	32031.	147.	146.	1.46e6	1.48e6
5	1921	4	0.00461	0.00433	31364.	32046.	145.	139.	1.46e6	1.48e6
6	1921	5	0.00372	0.00361	31175.	31847.	116.	115.	1.46e6	1.48e6
7	1921	6	0.00265	0.00393	30808.	31466.	81.7	124.	1.46e6	1.48e6
8	1921	7	0.00295	0.00351	30295.	30922.	89.5	108.	1.46e6	1.48e6
9	1921	8	0.00237	0.00285	29660.	30270.	70.4	86.4	1.46e6	1.48e6
10	1921	9	0.00198	0.00255	28923	29494.	57.4	75.3	1.46e6	1.48e6


```
# ... with 10,979 more rows, and abbreviated variable names 1: weighed_F,
# 2: weighed_M
```

```
dmf <- dmf |> mutate(weighed_rate_F = Female.x * Female.y / total_F, weighed_rate_M = Male.x * Male.y / total_M)

summary_mean <- dmf |> group_by(Year) |>
  summarize(mean_mortality_f = mean(weighed_rate_F, na.rm = TRUE),
            mean_mortality_m = mean(weighed_rate_M, na.rm = TRUE))

summary_mean |> ggplot(aes(x = Year)) + geom_line(aes(y = mean_mortality_f), colour = "firebrick4") +
  labs(title = "Female & Male ave_weighted mortality rates over time, Ontario",
```

```
y = "Average Weighted Mortality rate") + scale_color_manual("",values = c("Female"=
```



The trend of average weighted mortality rates over time of both gender was going down until 2000 year and after 2000 year, the trend became a little higher and stable.