Horizon-dependent Trade-offs and Undercoverage in Probabilistic Movement Forecasting

Public review revision, 14 October 2026; authorship unconfirmed

Abstract

A movement forecast can improve its distributional score without locating the target more accurately or providing reliable uncertainty regions. We examine these distinctions in an existing evaluation of fitted stochastic movement predictors applied to historical GPX position recordings with unverified carrier identities and transport modes, using 46 retrospectively selected recording-hash blocks, five simulation seeds and 512 particles per ensemble. At nominal 1 and 5 minutes, deterministic constant-speed extrapolation has lower energy score (ES) than the reference stochastic predictor Full; at 15 and 30 minutes the ordering reverses. Averaging the four targets gives ES 477.85 m for Full and 824.73 m for inertia, concealing that reversal. Within the stochastic procedures, a 300 s observation-interval variant reduces average ES by 23.40 m but increases final mean-position error by 9.64 m relative to its named control. Full's nominal 90% prediction disk covers only 46.52% of 30-minute targets. These are conditional descriptive findings: participant independence and physical clock provenance remain unverified, and the cohort was selected using future-route map support. Required forecast failures prevent terrain-family inference. The study shows why horizon-specific score, mean-position error and region coverage must be reported together; it does not establish stochastic-model superiority over a simple probabilistic baseline or operational reliability.

1 Introduction

Movement prediction is often summarized by a single accuracy measure, yet three distinct tasks are involved: assigning a distribution to future positions, locating a representative position, and describing a region that contains the realized target at its stated frequency. A gain on one task need not imply a gain on the others. This distinction matters when a probabilistic forecast is interpreted as a reliable search region.

Continuous-time correlated random walks model movement persistence and irregular observations [1]; switching state-space models represent latent regimes [2]. Step-selection models link movement to environmental selection [3, 4]. Learned trajectory predictors [5, 6], neural SDE generators [7] and interpretable stochastic system identification [8] offer additional modelling approaches. These works provide context, not performance baselines executed here. Our empirical subject is future position in selected geographic recordings; the modelling object is a conditional distribution over those positions. A stochastic position model represents uncertainty through a fitted drift, residual diffusion and latent components. The study evaluates an existing, limited collection of these fitted predictors.

Proper scoring rules evaluate predictive distributions against realized targets [9, 10]; they do not certify coverage of a particular prediction region. We therefore ask how distributional score, error of the predictive mean and marginal disk coverage change across four horizons. The empirical

contribution has three parts: a short/long-horizon reversal hidden by the average score, an example of score improvement with worse mean-position error, and severe long-horizon undercoverage despite substantial region area.

This focus is an interpretation of existing results, not a newly registered hypothesis test. The original questions remain explicit: RQ1 asks whether the registered changes in mode structure, observation interval, fitting/adaptation and numerical prediction yield a practically useful distribution-score change relative to their named controls. RQ2 asks whether selected terrain groups add predictive value beyond a fitted history baseline, through overall, leave-one-group-out and leave-one-group-in comparisons. To interpret the existing results we use three diagnostic questions: does the Full/inertia ordering depend on horizon; do ES and mean-position error agree; and do nominal prediction disks achieve their stated coverage? The first two research questions retain their original inference rules; the three diagnostics are descriptive readings, not new registered tests. The following sections define the observed object, construct the forecasts, and answer each question within those bounds.

2 Data and predictive procedures

2.1 Research object and forecasting task

The empirical objects are timestamped geographic position recordings from a historical GPX collection. Each recording traces the movement of an unidentified recording carrier. The retained archive does not establish whether a carrier is walking, cycling or travelling by another means, or whether two recordings belong to the same person. We therefore study position changes in the selected recordings, rather than the behaviour of a defined participant population. A cleaned segment is a derived portion of a recording; a window splits that portion into an observed prefix and future targets. Multiple windows can belong to one recording-hash block. Only one primary origin per selected block enters the main evaluation.

For an origin with saved time t_0, let $\mathcal{H}_0$ contain the allowed observed position/time prefix, ending at the current position x_0. Its last three pairs initialize the retained motion-history buffer. The task is to predict future positions conditional on this visible information and on fixed development-fitted parameters $\widehat{\xi}$. The method's internal east/north frame is anchored at the first visible position of that whole prefix, which need not be the first of the last three buffered points. Terrain prediction and evaluation instead use a frame anchored at the last visible position, the forecast origin. Method outputs are converted through geographic coordinates into that common scoring frame; the distributions in Equation (1) are expressed there. The method also has the retained clock and its solar proxy; the terrain branch additionally queries fixed map fields at simulated positions. True future positions are reserved for selection and evaluation and do not update a forecast. With actual matched elapsed times $u_1, \ldots, u_4$, the reported predictive object is

$$\widehat{P}_j = \mathcal{L}_{\widehat{\xi}}(X_{t_0+u_j} \mid \mathcal{H}_0), \qquad \widehat{P}_{N,j} = \frac{1}{N}\sum_{p=1}^{N} \delta_{X_j^{(p)}}, \quad j = 1, \ldots, 4. \tag{1}$$

Here δ_x is a point mass at x, and each simulated particle p has a position at all four targets. The ensemble approximates a model-defined distribution; it is not a sample of people. Its mean gives a representative location, while its radial quantiles define prediction disks. These summaries answer different questions about the same forecast. No separate measurement-error model is fitted: the saved target position is used as the evaluation reference, although its physical accuracy is not independently verified.

Development windows supply parameter estimates; final-evaluation windows supply forecast origins and observed outcomes. The main analysis asks how well those fixed predictors describe these outcomes as the saved forecast horizon increases. It does not estimate a causal law of human movement or the probability of finding a person in an operational search.

2.2 Evaluation population and clock

The task is single-recording position forecasting from a visible trajectory prefix. The retained data pipeline identifies its source as a historical OpenStreetMap GPX harvest, not the separate GeoLife cohort used in earlier work. User identities, transport-mode labels and original GPX segment boundaries are not preserved; these records cannot be treated as verified pedestrian participants. R1-derived segments supply midpoint history/target windows. The frozen recording-hash assignment separates development and evaluation: qualified development contains 404 outer-training windows (328 training and 76 adaptation) and 81 validation windows. Terrain fitting uses one first-future transition per outer-training window. Exact identity intersections across splits are zero, which does not establish participant or near-route independence. The scientific supplement details this assignment.

The released final-evaluation split contains 12,370 windows from 1,094 recording-hash blocks. Temporal eligibility retains 127 windows/89 blocks, joint feature validity retains 106/73, and the fixed primary-origin rule selects 46/46 (Figure 1). A block identifies a saved recording, not a verified independent participant. The selected cohort spans 12 countries and 27 source labels; 33/46 blocks are from Great Britain, Belgium or Luxembourg. It is not a representative population sample.




Figure 1: Evaluation selection and population descriptors. Counts distinguish windows from recording blocks; duration and speed panels show saved medians and interquartile ranges. Eligibility includes map validity along the observed future route, which is unavailable at deployment from the prefix alone. Selection exclusions are not forecast failures.

Future observations are used to select this cohort and to score forecasts, but are not inputs to prediction: online map queries use simulated positions. This distinction prevents direct target leakage during rollout without removing the retrospective selection restriction. The evaluation data had previously been inspected; the full influence of that inspection on design choices is not recoverable. This is not a fresh blinded holdout.

Targets are the actual observations matched to nominal 1, 5, 15 and 30 minutes, with elapsed-

time ranges 49–67, 294–305, 889–910 and 1785–1814 saved seconds. No target is interpolated to an exact nominal time. Physical acquisition-time and UTC provenance are unverified. An absolute offset affects the calendar input; a relative-scale distortion also affects rates, diffusion units and the physical interpretation of the horizons. All minute labels below refer to the saved clock.

2.3 Reference stochastic predictor and comparators

In the descriptive reference comparisons, Full denotes Full01, the ordinary three-mode fitted reference, not the all-terrain configuration. In saved local east/north coordinates $X = (X_e, X_n)^\top$ (metres), it uses

$$dX_t = \Theta_{m_t}^\top \psi(X_t, t)\, dt + L_{m_t}\, dW_t, \quad \psi = (1, X_e, X_n, S)^\top, \quad L_m L_m^\top = Q_m = \tau R_m. \tag{2}$$

What the stochastic equation represents. $X_t \in \mathbb{R}^2$ is position and $m_t \in \{1, 2, 3\}$ is an unobserved model component. $\Theta_m \in \mathbb{R}^{4\times 2}$ maps the intercept, two position coordinates and solar proxy to an east/north drift $b_m = \Theta_m^\top \psi$. This drift is a conditional displacement rate, measured in metres per saved second. W_t is a two-dimensional standard Wiener process with independent increments of covariance ΔI_2 over duration Δ; L_m maps them to position noise. Thus $Q_m = L_m L_m^\top$ is a diffusion covariance in square metres per saved second, and L_m has units of metres per square root of a saved second. The deterministic and random parts respectively represent a fitted average trend and residual variation, not a separately observed intention or velocity process.

For frozen drift and diffusion, the local increment has the illustrative law

$$X_{t+\Delta} - X_t \mid (X_t, m_t = m) \;\approx\; \mathcal{N}(b_m \Delta, Q_m \Delta)\,. \tag{3}$$

The mean displacement grows with Δ, while its random standard deviation grows with $\sqrt{\Delta}$. A sampled component adds uncertainty about which fitted trend applies. For constant component drifts and a segment-fixed mode, the corresponding illustrative mixture covariance is $u \sum_m \pi_m Q_m + u^2 \operatorname{Cov}_\pi(b_m)$: within-component diffusion and between-component drift variation need not grow at the same rate. Full has position- and solar-dependent drift, so this expression explains the ingredients rather than gives its actual multi-step variance or proves calibrated coverage.

How the reference parameters are learned. Development segments are resampled at the registered observation interval τ; an incomplete terminal interval is removed from fitting, while scoring observations remain unchanged. At $\tau = 60$ saved seconds, the 328 training, 76 adaptation and 81 validation windows supply 26,009, 5,762 and 6,149 complete transitions, respectively. A transition target is the two-dimensional displacement rate $y_i = (x_{i+1} - x_i)/\tau$. Within each fitting batch, segments are sorted by their net heading and divided into three balanced groups; all transitions from a segment inherit its group. The groups identify regression components, not externally validated walking, turning or transport states. For group m, with feature matrix D_m and rate matrix Y_m, the base estimates are

$$\widehat{\Theta}_m = \operatorname*{argmin}_{\Theta} \left\{ \tfrac{1}{2} \|D_m \Theta - Y_m\|_F^2 + \tfrac{\rho}{2} \|\Theta\|_F^2 \right\}, \qquad \rho = 10^{-6}, \tag{4}$$

$$\widehat{R}_m = \frac{1}{n_m - 1} \sum_{i \in m} (e_i - \bar{e}_m)(e_i - \bar{e}_m)^\top + 10^{-8} I_2, \quad e_i = y_i - \widehat{\Theta}_m^\top \psi_i. \tag{5}$$

The Frobenius norm sums squared entries; all coefficients, including the intercept, are penalised. n_m is the number of transitions in group m, and its base mixture weight is $n_m / \sum_k n_k$. R_m measures

centred residual variation in squared displacement-rate units. The small diagonal term stabilises the covariance calculation; it does not calibrate prediction regions. These are regression estimates, not continuous-time path-likelihood estimates.

Full fits training and adaptation batches separately, then combines their coefficients, covariances and proportions with weights 0.35 and 0.65. Validation selects a covariance scale $s \in \{0.55, 0.75, 1, 1.3, 1.7\}$ with drift held fixed. The saved R_m already includes this s^2 scaling. Its continuous-time embedding uses $Q_m = \tau R_m$: at a constant-coefficient duration τ, $Q_m\tau = R_m\tau^2$ matches the displacement covariance implied by the rate regression. This explains the time factor; it does not establish equivalence to maximum-likelihood estimation of an SDE. The chosen parameters remain fixed across evaluation origins and seeds. Full's saved component probabilities are 0.3350, 0.3302 and 0.3348, rather than a posterior re-estimated from each new prefix. The first visible position anchors the coordinate frame; forecasting starts at the last permitted prefix position. S is a solar/calendar proxy computed from the retained absolute tick [15], not a validated daylight mechanism. R_m is a centred fitted displacement-rate residual covariance; its conversion uses observation interval τ, not integration step h. Each particle draws a mode from saved weights π at the origin and segment-fixed Full retains that mode. Given frame, position, clock and random stream, ordinary Full does not read initial speed or recent heading. For the 46 primary origins, the origin-state buffer retains the last three observed positions and their strictly increasing saved times. The origin is the last point. Inertia extrapolates $\widehat{x}^{\mathrm{I}}(u) = x_0 + u\widehat{v}_0$, where $\widehat{v}_0$ is the secant from the first to last of those three points, divided by their saved elapsed time; no future position or future speed is used. These spans range from 2 to 61 saved seconds (median 7). Ordinary Full and inertia thus receive the same permitted prefix but use its movement information differently.

The validation selection proxy is a one-step rate-residual Gaussian using the unweighted average mode covariance, whereas evaluation scores a multi-mode, multi-step position distribution. The GMM variant groups training residuals once by $\arg\max(r_e, r_n, -r_e - r_n)$ but uses segment-heading labels for validation components: these labels are not aligned. Independently heading-sorted batches are also mixed by mode index in the transfer procedures [12]. These are implementation limitations, not evidence that the corresponding modelling ideas fail.

Evaluation uses $N = 512$ particles, five fixed seeds and maximum integration step $h = 5$ saved seconds. Full uses conditional Gaussian moment propagation, evaluating nonlinear conditions at the conditional mean; this is not an exact solution of a globally nonlinear SDE [11]. Four targets come from a continuing joint path, not independently sampled marginals. Observation-interval variants change fitting data, $Q = \tau R$ and update cadence together while retaining the integration cap. In particular, dt300 and dt600 mean observation intervals of 300 and 600 saved seconds, not integration steps. Identical-parameter aliases and multi-factor changes make the 28 saved identities procedures rather than 28 independent algorithms. All 21 registered primary contrasts retain their named Full controls.

From the fitted model to a forecast. Each particle starts at its internal-frame origin position X_0 and samples a mode from π; Full keeps that mode throughout its path. On each event interval $\Delta \leq 5$ saved seconds, write the locally affine drift as $\mathsf{A}_m x + a_n$, freezing the non-position feature at the conditional mean. A_m is the transpose of the two position rows of Θ_m; a_n contains the intercept and frozen solar term. Full uses the registered split kernel: an exact affine half-drift, an additive

noise increment, then another half-drift. Its matrices and conditional-moment update are

$$H = e^{\Delta \mathsf{A}_m/2}, \qquad J_{1/2} = \int_0^{\Delta/2} e^{v\mathsf{A}_m}\, dv,$$

$$F = H^2, \qquad J = (I+H)J_{1/2}, \qquad \Omega = H(\Delta Q_m)H^\top, \tag{6}$$

$$\mu_{n+1} = F\mu_n + Ja_n, \qquad C_{n+1} = FC_nF^\top + \Omega, \tag{7}$$

$$X_{n+1}^{(p)} = FX_n^{(p)} + Ja_n + Gz_{n,p}, \quad GG^\top = \Omega, \quad z_{n,p} \sim \mathcal{N}(0, I_2). \tag{8}$$

Initially $\mu_0 = X_0$ and $C_0 = 0$. The same noisy path supplies its positions at all four actual target times. The drift substeps are exact for frozen affine coefficients; the split covariance generally differs from the affine-exact integral $\int_0^\Delta e^{v\mathsf{A}_m} Q_m e^{v\mathsf{A}_m^\top}\, dv$. Evaluating the solar feature at the conditional mean also approximates the nonlinear full model. In particular, "FP" in the saved procedure denotes conditional Gaussian moments, not a spatial Fokker–Planck PDE solution. Numerical variants use Euler–Maruyama or affine-exact reference kernels, and MC/CRN variants evaluate nonlinear conditions along individual paths. Their complete definitions are in the scientific supplement. This construction produces the empirical measures in Equation (1); no subsequent radius calibration is part of the evaluated forecast.

2.4 Terrain branch and scope of attribution

The separate terrain branch augments a fitted three-position history drift with independently fitted residual conditioners and diffusion for ten configurations. Its fitted mechanism is specified below.

The terrain model's inputs and fitted mechanism. Here X_t uses the forecast-origin-anchored east/north frame. Let Z_t be a buffer of up to three visible or generated positions and times. Its first-to-last secant gives v_t, its normalized direction gives h_t, and η_t indicates whether that direction is valid. The history design is $r(Z_t) = (1, v_{t,e}, v_{t,n}, \|v_t\|, h_{t,e}, h_{t,n}, \eta_t)^\top$. The terrain drift and position dynamics are

$$b_g(X_t, Z_t) = B_0^\top r(Z_t) + \theta_g^\top \widetilde{\phi}_g(X_t, Z_t), \qquad dX_t = b_g(X_t, Z_t)\, dt + L_g\, dW_t. \tag{9}$$

$\widetilde{\phi}_g$ appends an intercept to the selected map/history feature vector, which contains encoded numerical values and validity masks. The road and river inputs describe nearest-feature distances and directions; land cover uses grouped indicators; surface inputs describe slope/orientation; registered interactions combine selected map and history quantities. These are fixed engineering representations. Their coefficients are learned from displacement rates, not constrained by a law of road attraction or river avoidance.

The shared B_0 is fitted first by ridge regression with strength 10^{-6}. Each of the 404 outer-training windows contributes one actual first-future transition $(\Delta x_i, \Delta t_i)$; the elapsed times need not equal the method interval τ. Then configuration g fits the remaining rate $y_i = \Delta x_i/\Delta t_i - B_0^\top r(Z_i)$ by

$$\widehat{\theta}_g = \operatorname*{argmin}_\theta \left\{ \frac{1}{2n}\|D_g\theta - Y\|_F^2 + \frac{\lambda}{2}\|\theta\|_F^2 \right\}, \quad n = 404, \quad \lambda = 10^{-4}. \tag{10}$$

D_g stacks the augmented feature rows. This two-stage fit is linear in its parameters; map lookup and history feedback nevertheless make its position dynamics state dependent. Each configuration estimates its own diffusion from fitted displacement residuals, $Q_g = L_gL_g^\top = n^{-1}\sum_i(\Delta x_i - \widehat{b}_{g,i}\Delta t_i)(\Delta x_i - \widehat{b}_{g,i}\Delta t_i)^\top/\Delta t_i$. The diffusion estimate differs from the method branch's centred τR.

Euler–Maruyama advances each path with drift $b_g\Delta$ and noise $\sqrt{\Delta}L_g z$; every five saved seconds the history buffer appends its generated position and recomputes the secant. Maps are queried at the predicted position. Future observed positions never enter this buffer.

The base configuration retains the shared history drift and its fitted history-only residual conditioner; it queries no map layer. All-terrain adds all registered groups and interactions; LOO removes one group from that full configuration, and LIO adds one group to the history baseline. Thus RQ2 compares these forecasting configurations on matched targets. The shared baseline and separate residual fits make the intended increment explicit, while refitting and feature penalties limit attribution to an individual factor.

Map sources and attribution limits. Model-source declarations identify WorldCover 2021 v200 [16], Overture transportation 2026-08-19.0 [23, 17], HydroRIVERS v1.0 [18, 19], and retained Mapzen Skadi / Copernicus GLO-30 rules [20, 21, 22]. These product declarations do not certify tile/track contemporaneity. Road/river direction points to the nearest feature, not along its tangent or flow; the retained roads are not the complete pedestrian network.

Constant validity masks, penalised intercepts, unstandardised features and refitted diffusion make effective regularisation differ among configurations. Thus even a complete comparison would concern configuration increments, not isolated terrain information. The scientific supplement gives full equations, input geometry, excluded road classes, fitting rules and procedure-specific controls. The terrain research question remains separate from ordinary Full.

3 Evaluation measures and inference

For particles $x_1, \dots, x_N$ and target y at one matched horizon, the execution-bound finite-ensemble energy score is

$$\mathrm{ES}_N = \frac{1}{N}\sum_i \|x_i - y\| - \frac{1}{2N(N-1)}\sum_{i\neq j}\|x_i - x_j\|, \qquad N \geq 2. \tag{11}$$

The second sum uses ordered pairs. For the deterministic inertial reference, ES is directly $\|x - y\|$. ES is measured in metres and lower is better; weighted ES is the equally weighted average over the four targets. Mean-position error is $e_j = \|\bar{x}_j - y_j\|$, with $\mathrm{ADE} = \frac{1}{4}\sum_j e_j$ and $\mathrm{FDE} = e_4$. FDE is neither best-particle nor average-particle error, and ADE is not a dense-path integral. ES includes interparticle spread and cannot be converted into error of the predictive mean. The first ES term measures particle distances to the target; the second subtracts half the average interparticle distance. Their balance evaluates the distribution, rather than the distance from its centre alone. Consequently, changing spread or mixture geometry can change ES without moving the mean closer to the target. This motivates reporting both quantities rather than using ES as a proxy for localisation.

A nominal level-q disk is centred at the ensemble mean, with radius the $\lceil qN \rceil$-th ordered distance of particles from that mean. Coverage records whether the true target lies inside; area is πr^2. Mean area is $\mathrm{E}[\pi r^2]$, not $\pi \mathrm{E}[r]^2$. These are horizon-wise marginal regions, not joint path regions. Enclosing about 90% of simulated particles does not guarantee 90% coverage of real targets. For multimodal clouds, the mean may lie in a low-density gap and a disk may include inaccessible terrain; it is not a highest-density or route-constrained region. For $\widehat{P}_j$'s own disk $D_{q,j}$, nominal mass concerns $\widehat{P}_j(D_{q,j}) \approx q$. Empirical reliability instead concerns the observed frequency of $Y_j \in D_{q,j}$ across origins. The two coincide only under suitable calibration of the predictive distribution; the

particle construction itself supplies no such guarantee. Area records how much spatial information that coverage costs.

For each recording block, the five seed values are averaged first; blocks then receive equal weight. There are 46 targets per horizon, not 230 independent targets. Six known-velocity and six point-only cases reuse primary records, giving 58 origin–mode cases rather than 58 independent blocks. Original family-wise simultaneous bootstrap-t intervals use 2,000 draws per stream, with shared block indices within a family; Holm-adjusted zero tests have a separate role [13, 14]. Intervals condition on the saved fits and unverified block independence / exchangeability; seeds represent simulation variation, not training or participant replication.

Deltas are candidate minus named control. The registered practical margin is $\delta = 41.100259$ m, 5% of validation inertial ES, not a validated operational utility threshold. Practical benefit/harm requires the entire original interval beyond $-\delta/+\delta$ and the original statistical, seed and qualification gates. Crossing zero is not equivalence. Practical equivalence, inconclusive information and unavailable evidence are distinct. The scientific supplement retains precise endpoints and finite bootstrap-tail limitations. The descriptive horizon and coverage summaries below are not new superiority or calibration hypothesis tests.

4 Results

4.1 Average score conceals a horizon reversal

Full's weighted ES is 477.85 m versus 824.73 m for inertia; the saved difference is -346.87 m, computed before rounding. At 1 minute, Full/inertia ES is 50.57/31.17 m; at 5 minutes it is 243.23/237.93 m. Thus inertia has lower ES at 1 and 5 minutes and Full at 15 and 30 minutes (Figure 2). The same figure separates ES from mean-position error. Four discrete targets do not identify an exact crossover time. The deterministic reference is a valid point-mass distribution




Figure 2: Full versus deterministic inertia at four nominal saved-clock horizons: ES (left) and predictive-mean error (right). Each point averages five seeds within each of 46 primary blocks, then weights blocks equally; inertia is deterministic. Lower is better. Connecting lines guide reading, not interpolation or evidence for a precise crossover.

for ES. It does not establish that Full's stochastic complexity is necessary relative to a simple probabilistic constant-velocity or correlated-random-walk model.

Table 1: Saved predictive-mean errors (m), using the original matched targets. These are not ES or mean particle errors.

Model	1 min	5 min	15 min	30 min
Full	73.77	332.31	812.26	1326.09
GMM	71.92	323.79	792.99	1299.11
dt300	75.88	340.10	825.76	1335.73
Inertial	31.17	237.93	934.03	2095.78

4.2 A better distribution score need not locate the mean better

The 300 s observation-interval procedure illustrates the distinction. Relative to its named Full01 control, weighted ES changes by -23.40 m but FDE by $+9.64$ m. Its mean-position error is also larger at each of the four saved targets (Table 1). ES accounts both for distances to the target and for spread between particles; it cannot be converted into mean-position error. The observation-interval comparison changes training exposure and diffusion binding as well as cadence, so this contrast does not isolate a numerical integration effect.

The dt300 ES simultaneous interval is $[-48.27, -2.47]$ m. Excluding zero does not satisfy the registered practical-benefit criterion: the whole interval would have to be below -41.100259 m, with the other gates met. The point thus communicates a metric conflict, not a practically superior method. All differences are calculated from unrounded source values.

4.3 Long-horizon disks substantially undercover

Full's nominal 90% disk coverage is 95.65%, 37.83%, 35.22% and 46.52% at the four horizons. At one minute the nominal 50% disk covers 33.04%, while at thirty minutes even the nominal 95% disk covers only 52.17%. The departure therefore depends on horizon and probability level; no radius rescaling has been tested. Across all 28 identities, thirty-minute 90% coverage ranges from 21.74% to 79.13%. The dt300 disk is about twice Full's mean area, yet its inclusion rate remains below the

Table 2: Thirty-minute nominal 90% disks: three configurations and nine saved summary values. Each configuration uses 46 blocks and five ensembles per block. Radius and area are averaged separately; the radius of the mean area is not the mean radius.

Model	Coverage (%)	Mean area (km^2)	Mean radius (m)
Full01	46.52	4.534	1183.6
GMM	45.22	3.079	989.5
dt300	64.35	9.109	1692.8

nominal 90%. A larger region can include more targets while its centre is less accurate. Conversely, GMM's smaller region does not by itself establish better uncertainty prediction. Coverage averages the same real target across five simulated disks in each block; there are 46 targets per horizon, not 230 independent targets. At block level this five-seed inclusion proportion takes only 0, 20, 40, 60, 80 or 100%. The 552 points in Supplement Figure 16 can overlap at those six levels; neither a block median of 100% nor a dense-looking cluster is a conditional calibration certificate. These observations support reporting size and inclusion together while retaining the complete block summaries. The 600 s procedure also has a larger thirty-minute mean area (12.609 km^2) and higher




Figure 3: Distribution-score versus endpoint-error trade-offs. All 21 comparisons (42 displayed scalar differences) retain five registered families and named controls. Each point averages five simulation seeds within each of 46 paired blocks, then weights blocks equally; ES first averages four targets while FDE uses only the last. Negative is favourable on both axes; the upper-left quadrant improves ES but worsens FDE. Facets have different scales. These means have no joint confidence region and do not establish a Pareto or global algorithm ranking.




Figure 4: Calibration and size from the saved 46-block, five-seed summaries. Left: 16 Full coverage values (four probability levels at four horizons); closeness to the diagonal, not the highest curve, is the target. Right: 12 mean areas (three configurations at four horizons), to be read with coverage rather than ranked alone. Connecting lines guide reading; they do not add intermediate horizons or a calibration test.

coverage (79.13%) than Full, but its FDE is 1340.94 m. This is another descriptive size–coverage trade-off, not evidence of calibrated uncertainty or improved mean location. No radius or parameter is retuned after inspection of this cohort.

4.4 Limits of component and terrain conclusions

Of the 21 registered primary method contrasts, 14 have practically equivalent and seven inconclusive finite-algorithm states; no original simultaneous interval lies wholly beyond the practical-benefit or harm boundary. All resolution-invariant states remain inconclusive. These states qualify the implemented finite procedures, not general algorithms. For example, all five GMM seed effects are negative but its block interval is $[-21.22, 16.33]$ m. Seed-direction agreement does not resolve uncertainty across recording blocks. Complete effects, seeds and qualification reasons are retained in Supplement Table 25 and the accompanying qualification discussion.

Three required primary terrain forecasts failed: all-terrain once and LIO-river/LIO-road once each. LOO means leave-one-group-out and LIO means leave-one-group-in. The overall/LOO grid has 1379/1380 successes and LIO has 1148/1150, with no pending forecasts; the grids share baselines and are not additive. Under the original complete-family rule, both whole families are unavailable. Successful subsets or individual map cases cannot replace that rule. This is not a positive or null terrain result. Two further failures affect auxiliary known-velocity method families, not the complete primary method families. Failure identities and logging limits are in the audit record; missing exception stacks prevent attributing a specific technical cause.

5 Discussion

5.1 What the model description lets us interpret

The task definition connects the horizon result to the two reference constructions. Inertia extends a short observed secant as a single straight trajectory; it can remain accurate when that local rate persists, but any rate error is multiplied by elapsed time. Full instead assigns probability to several fitted drift/diffusion components, without updating their weights from the new prefix's heading. This gives different uses of recent movement and different uncertainty representations. The observed short/long ordering is consistent with a changing forecasting trade-off, but does not identify which mechanism causes it: the comparison changes both mean dynamics and distributional spread, and no task-matched probabilistic inertia control isolates these effects. The constant-drift covariance explanation above is not a fitted explanation of the observed reversal.

The other two diagnostics follow from the forecast summaries. ES evaluates both distances to the realized target and distances within the ensemble, whereas FDE evaluates only the final mean. An ES decrease with FDE increase therefore establishes a disagreement between criteria, not a mathematical contradiction. A radial particle quantile describes the model's own spread; coverage compares that region with actual target positions. The observed undercoverage rejects the interpretation of Full's nominal disks as empirically reliable regions for this cohort, even when average ES is favourable. It does not identify whether drift error, insufficient spread, mixture geometry or selection is responsible.

5.2 Implications and remaining limits

The three findings address different decisions. The horizon reversal warns against choosing a predictor from its average score alone. The dt300 contrast shows that a distribution-score gain does not imply a more accurate representative location. The coverage results show that a model-internal particle quantile is not an empirical reliability guarantee. Together they support a reporting rule: pair horizon-specific distribution scores with mean-position errors and the coverage and size of any proposed region. This is a diagnostic implication, not a validated operational policy.

The calibration proxy differs from the evaluated predictive distribution, and GMM training/validation component labels differ. Both are plausible contributors to poor uncertainty behaviour, but these results do not identify causes or validate a correction. Likewise, wider regions can increase inclusion while retaining an inaccurate centre; coverage alone would reward sufficiently uninformative regions. The metric conflict is therefore substantive, not a reason to discard an unfavourable metric.

The scope is deliberately conditional. Retrospective future-route selection, previous evaluation exposure, uncertain recording independence and unverified clock provenance limit population and physical-time interpretation. Fixed fits and seeds omit training uncertainty. Parameter aliases, unmatched mode labels, joint procedure changes and representation-dependent penalties limit component attribution. The missing matched probabilistic baseline precludes a claim that complex stochastic structure is necessary. None of these limitations is removed by reproducing saved numerical summaries.

Future work should compare a task-matched probabilistic motion baseline, align mode labels before transfer and calibration, and evaluate calibration of the actual multi-step distribution. Terrain comparisons require complete families and matched effective penalties. Independent route grouping and clock-source verification are needed for broader inference. These are unexecuted studies, not results or editorial repairs supplied here.

6 Conclusion

This study evaluates conditional future-position distributions from selected GPX recordings, using fixed development-fitted stochastic position models and a visible-prefix inertial reference. Its diagnostic questions have concrete answers within the 46-recording cohort: inertia has lower ES at 1 and 5 saved minutes and Full at 15 and 30; dt300 lowers average ES by 23.40 m while increasing final mean-position error by 9.64 m; and Full's nominal 90% disk covers only 46.52% of 30-minute targets. Thus favourable average distribution scores do not establish uniformly better localisation or reliable prediction regions.

For the original questions, RQ1 yields 14 practically equivalent and seven inconclusive finite-procedure contrasts, with all resolution-invariant judgements inconclusive. RQ2 remains unavailable because required terrain forecasts failed under the registered complete-family rule. The learned terrain mechanism is specified, but its specification is not evidence of terrain efficacy. These results justify reporting horizon-specific ES, mean-position error, coverage and area together; they do not establish general algorithm superiority, the necessity of stochastic complexity or an operational search guarantee. Unverified carrier identity and clock provenance, retrospective selection and prior evaluation exposure bound the population and physical-time interpretation of this conclusion.

Data and reproducibility. The accompanying scientific supplement contains full specifications and all 50 tables; the separate audit record contains execution and revision history. The public repository provides saved aggregate evidence and reproduction code; private trajectories and local case maps are not redistributed. Repository: `https://github.com/HuAzurestar/learnable-sde-paper-to-predict-movement`. The public argument does not rely on those private case illustrations. Trajectory research-use permissions and the applicable ethics-review / consent basis require author verification; public availability of a source or map product is not taken as evidence of that authorization. This manuscript revision changes no fit, forecast, score or inference rule. Author identities, affiliations and contribution/conflict declarations require owner confirmation before submission and are not inferred from repository accounts.

References

[1] D. S. Johnson, J. M. London, M.-A. Lea and J. W. Durban (2008). Continuous-time correlated random walk model for animal telemetry data. *Ecology*, 89(5), 1208–1215. `https://doi.org/10.1890/07-1032.1`.

[2] Z. Ghahramani and G. E. Hinton (2000). Variational learning for switching state-space models. *Neural Computation*, 12(4), 831–864. `https://doi.org/10.1162/089976600300015619`.

[3] T. Avgar, J. R. Potts, M. A. Lewis and M. S. Boyce (2016). Integrated step selection analysis: bridging the gap between resource selection and animal movement. *Methods in Ecology and Evolution*, 7(5), 619–630. `https://doi.org/10.1111/2041-210X.12528`.

[4] (2017). Corrigendum. [Correction to Avgar et al. (2016)]. *Methods in Ecology and Evolution*, 8(9), 1168. `https://doi.org/10.1111/2041-210X.12725`.

[5] A. Gupta, J. Johnson, L. Fei-Fei, S. Savarese and A. Alahi (2018). Social GAN: Socially Acceptable Trajectories with Generative Adversarial Networks. *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition*, 2255–2264. `https://openaccess.thecvf.com/content_cvpr_2018/html/Gupta_Social_GAN_Socially_CVPR_2018_paper.html`.

[6] T. Salzmann, B. Ivanovic, P. Chakravarty and M. Pavone (2020). Trajectron++: Dynamically-Feasible Trajectory Forecasting with Heterogeneous Data. In *Computer Vision – ECCV 2020*, Lecture Notes in Computer Science, 12363, 683–700. Springer. https://doi.org/10.1007/978-3-030-58523-5_40.

[7] P. Kidger, J. Foster, X. Li and T. J. Lyons (2021). Neural SDEs as Infinite-Dimensional GANs. *Proceedings of the 38th International Conference on Machine Learning*, Proceedings of Machine Learning Research, 139, 5453–5463. https://proceedings.mlr.press/v139/kidger21b.html.

[8] T.-T. Gao, B. Barzel and G. Yan (2024). Learning interpretable dynamics of stochastic complex systems from experimental data. *Nature Communications*, 15, 6029. https://doi.org/10.1038/s41467-024-50378-x.

[9] T. Gneiting and A. E. Raftery (2007). Strictly proper scoring rules, prediction, and estimation. *Journal of the American Statistical Association*, 102(477), 359–378. https://doi.org/10.1198/016214506000001437.

[10] T. Gneiting, L. I. Stanberry, E. P. Grimit, L. Held and N. A. Johnson (2008). Assessing probabilistic forecasts of multivariate quantities, with an application to ensemble predictions of surface winds. *TEST*, 17, 211–235. https://doi.org/10.1007/s11749-008-0114-x.

[11] P. E. Kloeden and E. Platen (1992). *Numerical Solution of Stochastic Differential Equations*. Springer. https://doi.org/10.1007/978-3-662-12616-5.

[12] A. Nichol, J. Achiam and J. Schulman (2018). On First-Order Meta-Learning Algorithms. [Preprint], *arXiv:1803.02999v3*. https://arxiv.org/abs/1803.02999v3.

[13] S. Holm (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6(2), 65–70. https://www.jstor.org/stable/4615733.

[14] A. C. Davison and D. V. Hinkley (1997). *Bootstrap Methods and their Application*. Cambridge University Press. https://doi.org/10.1017/CBO9780511802843.

[15] NOAA Global Monitoring Division (n.d.). *General Solar Position Calculations*. [Online documentation]. Accessed 7 October 2026. https://gml.noaa.gov/grad/solcalc/solareqns.PDF.

[16] D. Zanaga et al. (2022). *ESA WorldCover 10 m 2021 v200*. [Dataset], version v200. https://doi.org/10.5281/zenodo.7254221.

[17] Overture Maps Foundation (n.d.). *Attribution and Licensing*. [Online documentation]. Accessed 6 October 2026. https://docs.overturemaps.org/attribution/.

[18] B. Lehner and G. Grill (2013). Global river hydrography and network routing: baseline data and new approaches to study the world's large river systems. *Hydrological Processes*, 27(15), 2171–2186. https://doi.org/10.1002/hyp.9740.

[19] HydroSHEDS (n.d.). *HydroRIVERS product documentation and licence*. [Online documentation], version v1.0. Accessed 6 October 2026. https://www.hydrosheds.org/products/hydrorivers.

[20] Mapzen / Tilezen (2017). *Terrain Tiles: Types of Terrain Tiles and Skadi format*. [Dataset distribution documentation]. Document revision e3d4351; accessed 7 October 2026. `https://registry.opendata.aws/terrain-tiles/`. `https://github.com/tilezen/joerd/blob/e3d4351ee3e5be333e23f47cb500e9a7c310656a/docs/formats.md`.

[21] Mapzen / Tilezen (2017). *Terrain Tiles: Attribution*. [Online documentation]. Document revision d8f587b; accessed 7 October 2026. `https://github.com/tilezen/joerd/blob/d8f587b73d26e0a0c42cdccd9ae8c55de4197763/docs/attribution.md`.

[22] Copernicus Programme / Sinergise (2021). *Copernicus Digital Elevation Model: GLO-30 Public*. [Dataset distribution documentation], 2021 release; accessed 7 October 2026. `https://registry.opendata.aws/copernicus-dem/`.

[23] Overture Maps Foundation (2026). *2026-08-19 release notes*. [Dataset release documentation]. Data version 2026-08-19.0; accessed 7 October 2026. `https://docs.overturemaps.org/blog/2026/08/19/release-notes/`.