



# Anglo-Chinese Junior College

JC2 Preliminary Examination  
Higher 2



A Methodist Institution  
(Founded 1866)

CANDIDATE  
NAME

**SUGGESTED SOLUTIONS**

FORM  
CLASS

TUTORIAL  
CLASS

INDEX  
NUMBER

## CHEMISTRY

Paper 3 Free Response

**9729/03**

28 August 2023  
2 hours

Candidates answer on the Question Paper.  
Additional Materials: Data Booklet

### READ THESE INSTRUCTIONS FIRST

Write your index number and name on all the work you hand in.  
Write in dark blue or black pen.  
You may use an HB pencil for any diagrams or graphs.  
Do not use staples, paper clips, glue or correction fluid.

Answer **all** questions in the spaces provided on the Question Paper. If additional space is required, you should use the pages at the end of this booklet. The question number must be clearly shown.

#### Section A

Answer **all** questions.

#### Section B

Answer **one** question.

Circle the number of the question you have attempted.

A Data Booklet is provided.

The use of an approved scientific calculator is expected, where appropriate.

At the end of the examination, fasten all your work securely together.


The number of marks is given in brackets [ ] at the end of each question or part question.

| For Examiners' use only |      |
|-------------------------|------|
| Section A               |      |
| 1                       | / 17 |
| 2                       | / 18 |
| 3                       | / 25 |
| Section B               |      |
| 4 / 5                   | / 20 |
| <b>Total</b>            | / 80 |


## Section A

Answer **all** the questions in this section.

- 1 Ibuprofen belongs to a class of drugs called non-steroidal anti-inflammatory drugs (NSAIDs) used to reduce inflammation, pain and fever. Pain, fever, and inflammation are caused by the release in the body of chemicals called prostaglandins. Ibuprofen blocks the enzyme that makes prostaglandins resulting in lower levels of prostaglandins.



The following diagram shows the first step in the synthesis route to form ibuprofen from compound **A**.




- (a) A series of experiments were carried out at 25 °C to find the effect of initial concentration of hydrogen cyanide, cyanide ions and **A** on the initial rate of formation of **B**.

Table 1.1

| experiment | initial concentration<br>/ mol dm <sup>-3</sup> |       |                    | initial rate of<br>formation of <b>B</b><br>/ mol dm <sup>-3</sup> s <sup>-1</sup> |
|------------|---|-------|--------------------|--|
|            | [ <b>A</b> ]                                    | [HCN] | [CN <sup>-</sup> ] |  |
| 1          | 0.020   | 0.060 | 0.060              | 1.00   |
| 2          | 0.020   | 0.050 | 0.050              | 0.833  |
| 3          | 0.020   | 0.050 | 0.060              | 1.00   |
| 4          | 0.025   | 0.050 | 0.050              | 1.042  |

- (i) Write a balanced equation for the reaction that produces compound **B**. [1]



- (ii) Use the data in Table 1.1 to deduce the order of reaction with respect to each of the reactants. [3]

Comparing Expt 1 and 3: **[A]** and **[CN<sup>-</sup>]** were unchanged. Initial rate of formation of **B** remained unchanged when **[HCN]** was increased by 1.2 times. The reaction is zero order with respect to HCN.

Comparing Expt 2 and 4: When **[A]** increased by 1.25 times while **[HCN]** and **[CN<sup>-</sup>]** were unchanged, initial rate of formation of **B** increased by 1.25 times. The reaction is first order with respect to **A**.

Comparing Expt 2 and 3: When **[CN<sup>-</sup>]** increased by 1.2 times while **[HCN]** and **[A]** were unchanged, initial rate of formation of **B** increased by 1.2 times. The reaction is first order with respect to **CN<sup>-</sup>**.

- (iii) Write the rate equation for this reaction. Hence, calculate the value of the rate constant using the data in experiment 2. Include its units. [2]


$$\text{Rate} = k[\mathbf{A}][\text{CN}^-]$$

Substituting values from expt. 2

$$0.833 = k(0.02)(0.05)$$


$$k = 833 \text{ mol}^{-1} \text{ dm}^3 \text{ s}^{-1}$$

- (iv) Describe a mechanism that is consistent with the rate equation deduced in (a)(iii) and indicate which step in the mechanism is the rate determining step. Explain your reasoning. [3]




Since HCN does not appear in rate equation, it is not involved in slow/rate determining step OR Only 1  $\text{CN}^-$  and 1 A is involved in the rate determining step/slow step as indicated by their order of reaction in the rate equation.

- (b) Explain, with the aid of a labelled Boltzmann distribution diagram, the effect on rate when the temperature is increased. [3]



- When temperature increases, there is an **increase in the number of particles with kinetic energy  $\geq E_a$**  (larger shaded area under the curve at  $T_2$  than  $T_1$ ).
- **This leads to an increase in the frequency of effective collisions** between reactant particles.
- Hence the **rate of reaction increases**.

- (c) Propose a 3-step synthesis of ibuprofen starting from the compound 1-isobutyl-4-vinylbenzene. Suggest the reagents and conditions, and draw the structure of the organic intermediate product in each step. [5]




1-isobutyl-4-vinylbenzene

**Step 1:** HBr(g) or HCl(g)

**Step 2:** alcoholic KCN or alcoholic NaCN, heat under reflux

**Step 3:** dilute H<sub>2</sub>SO<sub>4</sub>, heat under reflux



[Total: 17]

- 2 (a) Describe and explain the general trend in the first ionisation energy across Period 3.

[3]

- **There is an increasing nuclear charge** due to increase in number of protons.
- **There is an approximately constant shielding effect** as additional electrons are added to the same valence shell and provides negligible shielding effect.
- **Effective nuclear charge increases. OR Valence electrons thus experiences stronger nuclear attraction.**
- Therefore, **more energy** required to remove the valence electron. Hence, **first I.E. increases** across the period.

- (b) (i) Describe and explain the reactions, if any, of the chlorides  $\text{NaCl}$ ,  $\text{AlCl}_3$  and  $\text{PCl}_5$  with water. Write equations for all reactions that occur and suggest the pH of the resulting solutions. [4]

$\text{NaCl}$  - giant ionic structure. Solid readily soluble.

**Hydration:**  $\text{NaCl}(\text{s}) + \text{aq} \rightarrow \text{Na}^+(\text{aq}) + \text{Cl}^-(\text{aq})$  (**100% ionic  $\therefore$  no hydrolysis**).

$\text{Na}^+(\text{aq})$  ion **does not undergo hydrolysis** as it has a **low charge density** (low polarising power), thus the solution is neutral, **pH=7**.

$\text{AlCl}_3$  – Readily soluble with **substantial** hydrolysis

**Hydration:**



**Substantial hydrolysis:**




Substantial hydrolysis of  $\text{AlCl}_3$ , is due to the **high polarising power of the small, highly charged  $\text{Al}^{3+}$**  in solution, **distorts** the electron cloud of its surrounding water molecules and **weakens the O–H bond** in the complexing water molecules. It forms an acidic solution of **pH $\approx$ 3**.


$\text{PCl}_5$  is a simple covalent compound. It undergoes **complete** hydrolysis to give a strongly acidic solution, **pH=2**. **P has low-lying, vacant 3d-orbitals available for dative bonding with water molecules.**



- (ii) At certain temperatures, aluminium chloride exists as  $\text{Al}_2\text{Cl}_6$  molecules. Draw the structure of  $\text{Al}_2\text{Cl}_6$ . Label the co-ordinate bonds on your structure. [1]




- (c)  $\text{PCl}_5$  reacts with ketones in a 1 : 1 molar ratio to give gem-dichlorides. A gem-dichloride contains two chlorine atoms bonded to the same carbon atom.




- (i) Heating the above gem-dichloride, 3,3-dichloropentane, with ethanolic KOH produces a compound **S**,  $\text{C}_5\text{H}_8$ . Treating **S** with hydrogen over a nickel catalyst produces pentane.

Draw the displayed formula of **S** and on it, suggest values of **all** C–C–C bond angles. [2]




OR



- (ii) If 1,1-dichlorocyclohexane is similarly heated with an excess of ethanolic KOH, a product **T**,  $\text{C}_6\text{H}_9\text{Cl}$ , is obtained and no further loss of chlorine occurs. **T** decolourises aqueous bromine.

Suggest the structure of **T** and suggest why the expected loss of both chlorine atoms does not occur. [2]



Loss of both  $\text{Cl}$  would mean that  $\text{C}$  becomes  $\text{sp}$  hybridised and hence ring structure will not be possible.

- (d) Group 2 nitrates such as magnesium nitrate and barium nitrate decompose on heating.

- (i) Magnesium nitrate has a decomposition temperature of  $330^\circ\text{C}$  while barium nitrate has a decomposition temperature of  $592^\circ\text{C}$ .

Explain the difference in the decomposition temperatures. [2]

$\text{Mg}^{2+}$  has a **smaller ionic radius** compared to  $\text{Ba}^{2+}$ . Thus  $\text{Mg}^{2+}$  has a **higher charge density** and is able to **polarise the large  $\text{NO}_3^-$  anion to a larger extent** and the **N–O bonds are weakened more**. Hence magnesium nitrate decomposes at a lower temperature.



- (ii) Compare the decomposition temperature of aluminium nitrate to that of magnesium nitrate.

Explain your answer using data from the *Data Booklet*.

[2]

Aluminium nitrate has a lower decomposition temperature.

$\text{Al}^{3+}$  has a smaller ionic radius 0.050nm while  $\text{Mg}^{2+}$  0.065nm and a larger charge.

Thus  $\text{Al}^{3+}$  has a higher charge density and greater polarising power.

- (e) The value of the solubility product of barium sulfate is  $1.08 \times 10^{-10}$  at 25 °C.

Given that 100 cm<sup>3</sup> of 0.001 mol dm<sup>-3</sup> barium nitrate and 100 cm<sup>3</sup> of 0.001 mol dm<sup>-3</sup> sulfuric acid was mixed, calculate the mass of precipitate formed.

[2]

|                                    | $\text{Ba}^{2+}$ | $\text{SO}_4^{2-}$ |
|------------------------------------|------------------|--------------------|
| Initial / mol dm <sup>-3</sup>     | 0.0005           | 0.0005             |
| Change / mol dm <sup>-3</sup>      | -x               | -x                 |
| Equilibrium / mol dm <sup>-3</sup> | $0.0005 - x$     | $0.0005 - x$       |

$$K_{\text{sp}} = [\text{Ba}^{2+}(\text{aq})][\text{SO}_4^{2-}(\text{aq})]$$

$$1.08 \times 10^{-10} = (0.0005 - x)^2$$

$$x = 4.896 \times 10^{-4} \text{ mol dm}^{-3}$$

$$\begin{aligned} \text{mass of BaSO}_4 \text{ formed} &= 0.2 \times 4.896 \times 10^{-4} \times (137.3 + 32.1 + (16 \times 4)) \\ &= 0.0229 \text{ g} \end{aligned}$$


[Total: 18]

- 3 (a) (i) State and explain the conditions needed for a gas to behave ideally. [2]


High temperature so that that molecules have sufficient energy to break free from each other and experience negligible forces of interaction.

Low pressure so that the volume occupied by particles is negligible compared to the volume of the gas.

- (ii) Sketch and label a graph showing how the volume for a given mass of an ideal gas varies with temperature in Kelvin at constant pressure. [1]



- (b) Globally, ammonia is the second most produced chemical. The equation for the Haber process for manufacturing ammonia is shown.



A mixture of  $\text{H}_2$  and  $\text{N}_2$  in the molar ratio 3:1 is added to a sealed vessel and heated to  $450^\circ\text{C}$  with an iron catalyst. At equilibrium, 40% of the nitrogen had reacted. The total pressure in the vessel is  $2.80 \times 10^4 \text{ kPa}$  at equilibrium.

- (i) Write the expression for the equilibrium constant,  $K_p$ , for this reaction. Use your expression to calculate the value of  $K_p$  for this reaction. Include its units. [4]

|             | $3\text{H}_2(\text{g})$ | + | $\text{N}_2(\text{g})$ | $\rightleftharpoons$ | $2\text{NH}_3(\text{g})$ |
|-------------|-------------------------|---|------------------------|----------------------|--------------------------|
| Initial/kPa | 3x                      |   | x                      |                      | 0                        |
| Change/kPa  | -1.2x                   |   | -0.4x                  |                      | +0.8x                    |
| Eqm/kPa     | 1.8x                    |   | 0.6x                   |                      | 0.8x                     |

$$3.2x = 2.80 \times 10^4 \text{ kPa} \Rightarrow x = 8.75 \times 10^3 \text{ kPa}$$

$$P_{\text{NH}_3} = 7.00 \times 10^3 \text{ kPa}$$

$$P_{\text{H}_2} = 1.575 \times 10^4 \text{ kPa}$$

$$P_{\text{N}_2} = 5.25 \times 10^3 \text{ kPa}$$

$$\begin{aligned}
 K_p &= \frac{(P_{\text{NH}_3})^2}{(P_{\text{H}_2})^3 (P_{\text{N}_2})} \\
 &= \frac{(7.00 \times 10^3)^2}{(1.575 \times 10^4)^3 (5.25 \times 10^3)} \\
 &= 2.39 \times 10^{-9} \text{ kPa}^{-2}
 \end{aligned}$$

- (ii) Given that the Haber process is typically carried out at 450 °C and 200 atm, explain the rationale behind the conditions used in the Haber process for manufacturing ammonia. [2]


High temperature favours an increase in rate of reaction. However, the forward process is exothermic and increase in temperature will favour the backward endothermic process to absorb excess heat. Hence a compromise of 450 °C was chosen.

High pressure favours the forward process. However, too high a pressure leads to increase cost and quite good conversion/yield is already seen at moderate pressures. Hence 200 atm is chosen.

- (c) 80% of ammonia produced by the industry is used in agriculture as fertiliser, commonly sold in the form of ammonium nitrate,  $\text{NH}_4\text{NO}_3$ .

On 4<sup>th</sup> August 2020, in Beirut, Lebanon, approximately 2 750 tonnes of ammonium nitrate which had been improperly stored in a warehouse ignited and this led to significant loss of life, injuries, and destruction. This was the largest non-nuclear explosion in history, leaving a blast radius,  $r$ , of 62 m.

The decomposition of solid ammonium nitrate is shown.



1 tonne = 1 000 kg


- (i) Assuming that the explosion expands outwards in a hemisphere where volume =  $\frac{2}{3}\pi r^3$  and reaches 15 atm pressure at the edge of the blast radius, calculate the temperature of the gases within the blast radius. [2]

$$pV = nRT$$

$$T = \frac{pV}{nR} = \frac{15 \times 101325 \times \frac{2}{3} \pi (62^3)}{\frac{7}{2} \times \frac{2750000000}{(28.0+4.0+48.0)} \times 8.31} = 759 \text{ K}$$


$$V = 499 \times 10^3 \text{ m}^3 \text{ \& } n = 120 \times 10^6 \text{ mol}$$

- (ii) During the decomposition of solid ammonium nitrate, a side reaction occurs as shown.




Draw the 'dot-and-cross' diagram for  $\text{N}_2\text{O}$ .

[1]



- (d) Ammonium nitrate can also be found in instant cool packs for treating sports injuries. A typical instant cool pack contains 150 cm<sup>3</sup> of water which is separated from the solid ammonium nitrate by a thin film. When the film is broken, the ammonium nitrate dissolves, and the water reaches a temperature of 0 °C.

The equation for the dissolution of ammonium nitrate is shown.



- (i) Define the term *enthalpy change of solution*. [1]

Enthalpy change of solution is the energy change when one mole of substance (in its standard state) is completely dissolved in infinite dilution in an aqueous solution.

- (ii) Calculate the mass of ammonium nitrate needed to bring an instant cool pack from 25 °C to 0 °C. [2]

$$\begin{aligned} q &= mc\Delta T \\ &= 150 \times 4.18 \times 25 \\ &= 15\,675 \text{ J} \end{aligned}$$

$$\begin{aligned} \Delta H &= \frac{q}{n} = \frac{q}{m/M_r} \\ m &= \frac{q \times M_r}{\Delta H} \\ &= \frac{15\,675 \times 80.0}{25\,700} \\ &= 48.8 \text{ g} \end{aligned}$$

- (e) Lebanon exports perfumes. Geraniol, **E**,  $C_{10}H_{18}O$ , is commonly found in essential oils used in perfumes.

**E** does not react with 2,4-dinitrophenylhydrazine but reacts with hydrogen in the presence of heated nickel catalyst to form **F**,  $C_{10}H_{22}O$ . **E** reacts with warm acidified  $K_2Cr_2O_7$  to form **G**, which when purified, forms a gas **H** with aqueous sodium carbonate.

**E** reacts with hot acidified  $KMnO_4$  to form **J**,  $C_3H_6O$ , and **K**,  $C_5H_8O_3$  as the only organic products. **J** and **K** both give a yellow precipitate with alkaline aqueous iodine. **K** forms **H** with aqueous sodium carbonate and does not rotate plane polarised light.

Deduce the structures of **E** to **K**, identify the reactions involved and explain your reasoning. [10]


| Statement   | Type of reaction / Reason  |
|---|--|
| <b>E</b> , $C_{10}H_{18}O$  | <b>E</b> has two degrees of unsaturation [✓]   |
| <b>E</b> does not react with 2,4-dinitrophenylhydrazine but reacts with hydrogen in the presence of heated nickel catalyst to form <b>F</b> , $C_{10}H_{22}O$ . | <b>E</b> does not contain carbonyl group [✓]<br><b>E</b> underwent reduction [✓]<br>Two alkenes in <b>E</b> [✓]<br><b>E</b> and <b>F</b> are alcohols [✓]  |
| <b>E</b> reacts with warm acidified $K_2Cr_2O_7$ to form <b>G</b> , which when purified, forms a gas <b>H</b> with aqueous sodium carbonate.                    | Oxidation of <b>E</b> [✓]<br><b>H</b> is $CO_2$ (awarded as structure)<br>Hence <b>G</b> is carboxylic acid [✓]<br>Therefore <b>E</b> contains primary alcohol [✓]                                   |
| <b>E</b> reacts with hot acidified $KMnO_4$ to form <b>J</b> , $C_3H_6O$ , and <b>K</b> , $C_5H_8O_3$ as the only organic products.                             | Oxidation of <b>E</b> [✓]<br>Loss of 2 Cs implies two $CO_2$ loss via ethanedioic acid (not enough $C=C$ for 2x terminal alkenes) [✓]<br>$1^\circ$ alcohol on C-1 and alkene (H/R groups) on C-2 [✓] |
| <b>J</b> and <b>K</b> both give a yellow precipitate with alkaline aqueous iodine.  | <b>J</b> and <b>K</b> contain $CH_3CO-$ group [✓]  |
| <b>K</b> forms <b>H</b> with aqueous sodium carbonate and does not rotate plane polarised light.  | <b>K</b> is a carboxylic acid [✓]<br><b>K</b> is non-chiral [✓]  |

**H** is  $CO_2$


**J** is  $CH_3COCH_3$

**K** is  $CH_3COCH_2CH_2CO_2H$


E




or




or




F




or




or




G



or



or



[Total: 25]

## Section B


Answer **one** question from this section.

4 Copper, nickel, manganese and iron are transition elements.

- (a) When a particular copper ore was reduced, an alloy was produced which was composed mainly of copper, but with nickel and silver as minor impurities. It contained no other metal.

In order to purify it, this alloy was made the anode of an electrolytic cell, with a pure copper cathode and aqueous  $\text{CuSO}_4$  as electrolyte.

- (i) Explain, with reference to relevant  $E^\ominus$  values, what happens to the nickel and silver impurities during the purification procedure. [2]



When  $\text{Cu(s)}$  is oxidised at the anode,  $\text{Ni(s)}$  which has a negative reduction potential is also oxidised. At the cathode,  $\text{Cu}^{2+}(\text{aq})$  is preferentially reduced over  $\text{Ni}^{2+}$  due to its higher reduction potential.  $\text{Cu(s)}$  is deposited at the cathode while  $\text{Ni}^{2+}$  ions remain in the solution.

$\text{Ag(s)}$  is not oxidised as its reduction potential is more positive. It merely dropped off to the bottom of the electrolytic cell as anode sludge.

A current of 3.50 A was passed through the cell for 11 minutes 47 seconds, and the electrodes were removed, washed, dried and finally weighed. It was found that anode had lost 0.947 g.

After filtering it off and drying it, the deposit under the anode weighed 0.0497 g.

On adding an excess of dimethylglyoxime to the electrolyte, the highly insoluble red complex with formula  $\text{Ni}(\text{C}_4\text{H}_7\text{N}_2\text{O}_2)_2$ , molar mass  $288.7 \text{ g mol}^{-1}$ , was precipitated. Its mass was 0.587 g.

- (ii) Calculate the maximum increase in mass of the cathode. [1]

$$\begin{aligned} Q &= 3.50 \times 707 = 2474.5 \text{ C} \\ n(\text{e}^-) &= 2474.5/96500 = 2.564 \times 10^{-2} \\ \text{At the cathode: } \text{Cu}^{2+} + 2\text{e}^- &\rightarrow \text{Cu} \\ n(\text{Cu}) &= 1.28 \times 10^{-2} \text{ mol} \\ \text{maximum increase in mass at cathode} &= 0.814 \text{ g} \end{aligned}$$

- (iii) Calculate the actual masses of copper, nickel and silver removed from the alloy. [3]

Mass of silver = 0.0497 g


$n[\text{Ni}(\text{C}_4\text{H}_7\text{N}_2\text{O}_2)_2] = 0.587 / 288.7 = 2.033 \times 10^{-3}$

mass of Ni removed from alloy =  $2.033 \times 10^{-3} \times 58.7 = 0.119 \text{ g}$

mass of Cu removed from alloy =  $0.947 - 0.119 - 0.0497 = 0.778 \text{ g}$

In 1905, L. A. Chugaev suggested the use of dimethylglyoxime as a specific reagent for the qualitative and quantitative determination of nickel, earning dimethylglyoxime the nickname of “Chugaev’s reagent”. It is a bidentate ligand.

The structure of Chugaev’s reagent is given below.



- (iv) Suggest a reagent that reacts with butanedione to form Chugaev’s reagent. [1]

$\text{HONH}_2$

- (v) State the coordination number of Ni in the  $\text{Ni}(\text{C}_4\text{H}_7\text{N}_2\text{O}_2)_2$  complex. [1]  
4



- (b) Potassium manganate(VII) plays an important role in analytical chemistry.

The figures below show the standard electrode potentials,  $E^\ominus$ , of various redox half reactions. For example, in Fig. 4.1, the line shows that the  $E^\ominus$  for  $\text{MnO}_4^-/\text{Mn}^{2+}$  half-cell is +1.52 V.




Fig. 4.1






Fig. 4.2

- (i) Use Fig. 4.2 to predict what sulfur-containing species might be formed from sodium sulfite,  $\text{Na}_2\text{SO}_3$ , when it is titrated with acidified potassium manganate(VII) solution. Give a reason for your prediction. [2]



- (ii) Given that the two sulfur atoms in  $\text{S}_2\text{O}_6^{2-}$  are indistinguishable, draw the structure of  $\text{S}_2\text{O}_6^{2-}$ . [1]



- (c) In another experiment, a solution of iron(II) sulfate was titrated against  $0.020 \text{ mol dm}^{-3}$  acidified potassium manganate(VII), maintaining a constant pH throughout the titration. The electrode potential was measured against a standard hydrogen electrode and the following data was obtained.

| volume of $\text{KMnO}_4(\text{aq})$ added / $\text{cm}^3$ | $E^\ominus$ at $25^\circ\text{C}$ / V |
|--|---------------------------------------|
| 20.0   | +0.734                                |
| 40.0   | +0.760                                |
| 60.0   | +0.780                                |
| 80.0   | +0.806                                |
| 120.0  | +1.512                                |
| 140.0  | +1.515                                |

The electrode potential of the mixture at  $25^\circ\text{C}$ ,  $E^\ominus$ , is given by the expression

$$E = E^\ominus (\text{Fe}^{3+}/\text{Fe}^{2+}) + 0.0257 \ln \frac{[\text{Fe}^{3+}]}{[\text{Fe}^{2+}]}$$

- (i) Using the given expression and relevant information from the *Data Booklet*, determine the value of  $E$  when  $[\text{Fe}^{2+}]$  and  $[\text{Fe}^{3+}]$  are equal. [1]

When  $[\text{Fe}^{2+}] = [\text{Fe}^{3+}]$ ,  $\ln \frac{[\text{Fe}^{3+}]}{[\text{Fe}^{2+}]} = 0$ .

Hence,  $E = E^\ominus (\text{Fe}^{3+}/\text{Fe}^{2+}) = +0.77 \text{ V}$ .

- (ii) Estimate the volume of  $\text{KMnO}_4$  added which corresponds to your value of  $E$  in (c)(i). [1]

$50.0 \text{ cm}^3$

- (iii) Explain why there is a rapid increase in the value of  $E$  between 80.0 cm<sup>3</sup> and 120.0 cm<sup>3</sup> of KMnO<sub>4</sub>. [1]


The equivalence point is reached between 80 and 120 cm<sup>3</sup> (precisely at 100 cm<sup>3</sup>). This means that the  $E$  value switches from a calculation based on Fe<sup>3+</sup>/Fe<sup>2+</sup> to one based on MnO<sub>4</sub><sup>-</sup>/Mn<sup>2+</sup>.

- (d) Lanthanum is a Period 6 transition element. It exhibits an oxidation state of +3 in all of its compounds. Lanthanum iodate(V) is stable at room temperature, but when heated to 560 °C it decomposes, as shown in reaction 1.

Reaction 1      lanthanum iodate(V) → lanthanum periodate(VII) + oxygen + iodine

There are three common periodate(VII) ions that occur in compounds: IO<sub>4</sub><sup>-</sup>, IO<sub>5</sub><sup>3-</sup> and IO<sub>6</sub><sup>5-</sup>. The ion contained in the lanthanum periodate(VII) product of reaction 1 can be deduced by determining the stoichiometry of the reaction in the following way.

- A 1.50 g sample of lanthanum iodate(V) was heated at 560 °C in a closed vessel until no further reaction takes place.
- The iodine produced was dissolved in hexane and titrated against 0.300 mol dm<sup>-3</sup> sodium thiosulfate solution.



The end-point occurs when 18.10 cm<sup>3</sup> of 0.300 mol dm<sup>-3</sup> sodium thiosulfate was added.

- (i) State the shape of IO<sub>4</sub><sup>-</sup>. [1]

Tetrahedral

- (ii) State the colour of iodine dissolved in hexane. [1]

Violet/purple

- (iii) Use the titration results to show that the IO<sub>3</sub><sup>-</sup> : I<sub>2</sub> molar ratio in this reaction is 5:2. Hence write a balanced ionic equation showing the decomposition of IO<sub>3</sub><sup>-</sup> to I<sub>2</sub>, O<sub>2</sub> and the appropriate periodate(VII) ion. [3]


$$\text{Amt of thiosulfate} = 0.300 \times \frac{18.10}{1000} = 0.00543 \text{ mol}$$

$$\text{Amount of iodine} = 0.00543/2 = 0.002715 \text{ mol}$$


$$\text{Amount of lanthanum iodate, La(IO}_3)_3 = \frac{1.50}{3(126.9+16 \times 3)+138.9} = 0.0022604 \text{ mol}$$

$$\begin{aligned} \text{Amount of iodate ions} \\ = 0.0022604 \times 3 = 0.00678 \text{ mol} \end{aligned}$$

$$\text{Mole ratio of IO}_3^- : \text{I}_2 = 0.00678 : 0.002715 = 5 : 2 \text{ (shown)}$$



- (iv) Construct the full chemical equation for the decomposition of lanthanum iodate(V). [1]



[Total: 20]

5 Cobalt and chromium are transition elements.

- (a) A cobalt-chrome alloy is a metal alloy of cobalt and chromium. A sample of this alloy was dissolved in dilute HCl to form a solution containing  $1.55 \times 10^{-2} \text{ mol dm}^{-3}$  of  $\text{Cr}^{3+}(\text{aq})$  and  $5.77 \times 10^{-3} \text{ mol dm}^{-3}$  of  $\text{Co}^{2+}(\text{aq})$ . Aqueous NaOH was then added slowly to  $25 \text{ cm}^3$  of the solution.

Using the  $K_{\text{sp}}$  information below, show that the addition of NaOH will not allow for the separation of the  $\text{Cr}^{3+}$  and  $\text{Co}^{2+}$  metal cation ions in the above solution.

| compound                 | $K_{\text{sp}}$        |
|--------------------------|------------------------|
| $\text{Cr}(\text{OH})_3$ | $1.60 \times 10^{-20}$ |
| $\text{Co}(\text{OH})_2$ | $5.92 \times 10^{-15}$ |

[3]

For  $\text{Cr}^{3+}$ :

$$1.60 \times 10^{-20} = 1.55 \times 10^{-2} [\text{OH}^-]^3$$

$$[\text{OH}^-] = 1.0106 \times 10^{-6} \text{ mol dm}^{-3}$$

For  $\text{Co}^{2+}$ :

$$5.92 \times 10^{-15} = 5.77 \times 10^{-3} [\text{OH}^-]^2$$

$$[\text{OH}^-] = 1.0129 \times 10^{-6} \text{ mol dm}^{-3}$$

Since the  $[\text{OH}^-]$  required for both precipitates to form (i.e.  $\text{IP} = K_{\text{sp}}$ ) is similar, it does not allow for the separation of the two metal ions.

- (b) Given that



Explain why the reducing power of  $\text{Co}(\text{II})$  increases in the presence of excess aqueous ammonia. [1]



As the standard reduction potential of  $\text{Co}^{3+}$  to  $\text{Co}^{2+}$  is more positive than that of  $\text{Co}(\text{NH}_3)_6^{3+}$  to  $\text{Co}(\text{NH}_3)_6^{2+}$  there is a higher tendency for  $\text{Co}(\text{NH}_3)_6^{2+}$  to be oxidised.

- (c) There exists two cobalt-containing compounds, **X** and **Y**, which have the same relative formula mass of 259.3. Both contain ammonia and two different halides associated with the cobalt(II) ion.

When excess silver nitrate solution was added to equal masses of each of them, **X** produced 1.31 g of precipitate while **Y** produced 1.00 g of another precipitate.


By considering the ratio of the  $M_r$  of AgBr to the  $M_r$  of AgCl, deduce the molecular formula of the cobalt-containing compound and hence, determine the formula of the cation in **Y**. [3]

$1.31 / 1.00 = 1.31$  which is the ratio between the molar masses of AgBr and AgCl  
 $\Rightarrow$  the bromide is not datively bonded to the cobalt.

MF is  $\text{CoN}_5\text{H}_{15}\text{BrCl}$

Formula of the cation in **Y** –  $\text{Co}(\text{NH}_3)_5\text{Br}^+$

- (d) The simplest aldehyde, methanal,  $\text{HCHO}$ , can readily be oxidised to methanoic acid by acidified potassium manganate(VII).



The reaction proceeds very slowly at room temperature. A small amount of vanadium(III) ions,  $\text{V}^{3+}$ , are added to the mixture to speed up the reaction.

The half-equation for the reduction of methanoic acid is shown.



Use data from the *Data Booklet* and the half-equation given to construct equations and justify the two-step mechanism of the reaction. [3]


Step 1:



$$E_{\text{cell}}^\ominus = +1.52 - (+0.34) = \underline{\underline{+1.18\text{V}}}$$

Since  $E_{\text{cell}}^\ominus > 0$ , reaction is **feasible**.

Step 2:




$$E_{\text{cell}}^\ominus = +0.34 - (-0.03) = \underline{\underline{+0.37\text{V}}}$$

Since  $E_{\text{cell}}^\ominus > 0$ , reaction is **feasible**.

(e) Chromium is used in certain nutritional supplements.

- (i) Chromium(III) picolate is sold as a nutritional supplement to treat type 2 diabetes and promote weight loss.




chromium(III) picolate

It exists as two stereoisomers. Specify the type of stereoisomerism and draw either stereoisomer. You may represent each ligand as shown below.



[2]


### Enantiomerism



- (ii) By quoting and using relevant  $E^\ominus$  values from the *Data Booklet*, explain why the blue solution, containing  $\text{Cr}^{2+}(\text{aq})$ , obtained by dissolving chromium metal in dilute sulfuric acid slowly turns green even in the absence of air. [2]

Reaction is slow because of high activation energy due to electrostatic forces of repulsion between 2 cations.

$E^\ominus / \text{V}$



Calculation of  $E^\ominus_{\text{cell}}$ , and deduction that the redox reaction is feasible.

The  $\text{H}^+$  acts as oxidising agent, oxidising the blue  $\text{Cr}^{2+}$  to the green  $\text{Cr}^{3+}$  (or eqn).

- (f) Copper is also used in certain nutritional supplements.


Copper(II) glycinate is used as a dietary supplement for copper in animal feeds.

Copper(II) glycinate is formed when the  $\alpha$ -amino acid glycine ( $R = H$ ) is added to an aqueous solution of copper(II) ethanoate.


A blue precipitate of the square pyramidal copper(II) glycinate ( $C_4H_{10}CuN_2O_5$ ) is formed which contains one coordinated water molecule.

When heated, this precipitate undergoes isomerisation between three different stereoisomers. However, throughout the isomerisation, the water molecule remains in the axial position.

Draw all three stereoisomers and specify which is the pair of cis-trans isomers. You may represent each glycine ligand as shown below.




[3]





1<sup>st</sup> and 2<sup>nd</sup> – cis-trans isomers    1<sup>st</sup> and 3<sup>rd</sup> – cis-trans isomers

- (g) The Kharasch–Sosnovsky reaction is the radical oxidation of an allylic alkene to an allylic alcohol using a copper catalyst and a peroxide.




The mechanism is believed to involve radical intermediates and copper in its +1, +2 and +3 oxidation states, via four steps.

The first two steps of the mechanism are provided below.




- (i) Draw step 2 of the mechanism. Show all relevant curly arrows. [1]



- (ii)  $\text{CH}_2=\text{CHCHHR}-\text{Cu(III)}-\text{OH}$  is an intermediate formed in step 3, and reacts in step 4.

Write the equations for steps 3 and 4. [2]



[Total: 20]