


圖書，科學與數學，數學

Look inside ↗



查看所有3圖像

Linear Algebra, 4th Edition (English)

Hardcover – 十一月 21, 2002

Stephen H. Friedberg ▾ (Author), Arnold J. Insel (Author), & 1 更多

★★★★☆☆ ▾ 70 消費者評論

ISBN 13 碼: 978-0130084514 | ISBN 10 碼: 0130084514 | 版本: 4th

已使用

定價: US\$61.66

租金

定價: USD47.25

22 已使用 , 從 US\$62.32 | 2 租賃 從 US\$46.19

› 請參閱所有 6 格式和版本

電子教科書

US\$122.38

Hardcover

來自於US\$55.85

閱讀使用我們的 免費應用程式

出租 US\$46.19

22 二手 起 US\$55.85

:



Linear Algebra, Books a la Carte edition (5th Edition)

US\$109.99

現貨。

prime student


College student? Get FREE shipping and exclusive deals

[LEARN MORE](#)

For courses in Advanced Linear Algebra.

This top-selling, theorem-proof text presents a careful treatment of the principal topics of linear algebra, and illustrates the power of the subject through a variety of applications. It emphasizes the symbiotic relationship between linear transformations and

[閱讀更多](#)




東區, 新竹市天氣預報和天氣狀況—The...

iCloud 會備份哪些內容？ - Apple 支援

Mathematical Analysis - Tom M. A...

最近班表

Mathematical Analysis



Tom M. Apostol

Addison-Wesley, 1974 - 492 頁



4 書評

It provides a transition from elementary calculus to a

搜尋書籍內容

搜尋

過網
享受開

讀者評論 - 撰寫評論

我們找不到任何評論。

相關書籍

This book is printed on acid-free paper.

WALTER RUDIN
*Professor of Mathematics
 University of Wisconsin—Madison*

Library of Congress Cataloging in Publication Data

Rudin, Walter, date
Principles of mathematical analysis.
 (International series in pure and applied mathematics)
 Bibliography p.
 Includes index.
 I. Mathematical analysis. I. Title.
 QA300.R8 1976 515 75-17903
 ISBN 0-07-054235-X

Principles of Mathematical Analysis

THIRD EDITION

PRINCIPLES OF MATHEMATICAL ANALYSIS

Copyright © 1964, 1976 by McGraw-Hill, Inc. All rights reserved.
 Copyright 1953 by McGraw-Hill, Inc. All rights reserved.
 Printed in the United States of America. No part of this publication
 may be reproduced, stored in a retrieval system, or transmitted, in any
 form or by any means, electronic, mechanical, photocopying, recording, or
 otherwise, without the prior written permission of the publisher.

28 29 30 DOC/DOC 0 9 8 7 6 5 4 3 2 1 0

4 / 354



Rudin, Principles Of Mathematical Analysis



Publication date	1976-01-01
Topics	mathematics, analysis, calculus
Collection	opensource
Language	English

Principles of Mathematical Analysis (3rd edition, 1976) by Walter Rudin

Optimized PDF (5.0 MB) and DJVU (1.8 MB)

ISBN 10: 007054235X
 ISBN 13: 9780070542358

First published in 1953 (revised 1964, 1976), "Principles of Mathematical Analysis" has become a classic introduction to the field. Generations of students have become acquainted with this slim, blue-covered volume (fondly referred to as "Baby Rudin"), which continues to be used for honours courses at the top mathematics departments in the US. "Principles of Mathematical Analysis" has also been influential internationally through translated editions in Russian, German, French, and Chinese, among others.

Walter B. Rudin (1921-2010) was an Austrian-American mathematician and Professor of Mathematics at the University of Wisconsin, Madison (1959-1991). He was known for his contributions to complex and harmonic analysis and authorship of several widely adopted textbooks, including "Principles of Mathematical Analysis", "Real and Complex Analysis", and "Functional Analysis".

Identifier	RudinW.PrinciplesOfMathematicalAnalysis3e-MGH19769780070542358353S
Identifier-ark	ark:/13960/t6vx5986m
Ocr	ABBY FineReader 11.0
Ppi	300
Scanner	Internet Archive HTML5 Uploader 1.6.2

11,524 Views

11 Favorites

3 Reviews

DOWNLOAD OPTIONS

ABYY GZ	1 file
DAISY	1 file
For print-disabled users	
DJVU	1 file
EPUB	1 file
FULL TEXT	1 file
ITEM TILE	1 file
KINDLE	1 file
PDF	1 file
SINGLE PAGE PROCESSED JP2 ZIP	1 file
TORRENT	1 file
SHOW ALL	13 files
	7 Original

Mathematical Background

(Appendix
A)

1. Norm

Definition: A function $f: \mathbb{R}^n \rightarrow \mathbb{R}$

with $\underline{\text{dom } f} = \mathbb{R}^n$ is called a
norm if \uparrow domain

① $f(x) \geq 0 \quad \forall x \in \mathbb{R}^n$ (nonnegativity)

② $f(x) = 0$ only if $x = 0$.
 \uparrow (f is definite)

③ $f(tx) = |t| f(x) \quad \forall x \in \mathbb{R}^n, t \in \mathbb{R}$

④ $f(x+y) \leq f(x) + f(y)$ (homogeneity)
(triangle inequality)

→ Denote $f(x) = \|x\|$

which can be considered as a generalization
of the absolute value on \mathbb{R} .

Example: ℓ_p -norm $p \geq 1$ for $x \in \mathbb{R}^n$ $x = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$

$$\|x\|_p \triangleq \left(|x_1|^p + |x_2|^p + \cdots + |x_n|^p \right)^{1/p}.$$

$$p=1 : \|x\|_1 = |x_1| + |x_2| + \cdots + |x_n|$$

$$p=2 : \|x\|_2 = \left(|x_1|^2 + |x_2|^2 + \cdots + |x_n|^2 \right)^{1/2}$$

$$= (x_1^2 + x_2^2 + \cdots + x_n^2)^{1/2}$$

(Euclidean norm).

$p=\infty$: (maximum norm)
infinity norm

$$\|x\|_\infty = \lim_{p \rightarrow \infty} \|x\|_p = \max\{|x_1|, |x_2|, \dots, |x_n|\}$$

Remark: every norm is a convex function.

Quadratic norms

For $P \in \mathcal{S}_{++}^n$, $x \in \mathbb{R}^n$

$$\|x\|_P = (x^T P x)^{1/2}$$

$$= \|P^{1/2} x\|_2$$

$\mathbb{R}^{m \times n}$: matrices

\mathcal{S}^n : $n \times n$ symmetric matrices over \mathbb{R} .

\mathcal{S}_+^n : $n \times n$ positive semi-definite matrices

\mathcal{S}_{++}^n : $n \times n$ positive definite matrices.

If $P \in S_+$, $x_i^T x_j = \delta_{ji}$

P has a spectral decomposition

$$P = \sum_i \lambda_i x_i x_i^T \quad \text{with } \lambda_i \geq 0.$$

$$P^{1/2} \triangleq \sum_i \sqrt{\lambda_i} x_i x_i^T.$$

$$\begin{aligned} P^{1/2} \cdot P^{1/2} &= \left(\sum_i \sqrt{\lambda_i} x_i x_i^T \right) \left(\sum_j \sqrt{\lambda_j} x_j x_j^T \right) \\ &= \sum_i \sum_j \sqrt{\lambda_i} \sqrt{\lambda_j} x_i x_i^T \underbrace{x_j x_j^T}_{\delta_{ij}} \\ &= P \end{aligned}$$

A matrix M is normal if $M^T M = M M^T$.

For a normal matrix $M \in \mathbb{R}^{n \times n}$, $x \in \mathbb{R}^n$

if $Mx = \lambda x$ for some $\lambda \in \mathbb{R}$.

we say that x is an eigenvector of M .
 λ eigenvalue.


If M is normal, there exist λ_i, x_i such that

$$M = \sum_i \lambda_i x_i x_i^T.$$

Ex. $M = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, $MM^T = M^TM$

$$M \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \sqrt{\sum} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \sqrt{2} \left(\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right)$$

$$M = \sqrt{2} \left(\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right) \left(\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \end{pmatrix} \right)$$



For a symmetric & normal matrix M ,
if all the eigenvalues of M are $\geq 0, (> 0)$
 M is called positive semidefinite.
(positive definite)