


Parallel Coupled Joint Mechanism

The PCJM element allows translational as well as rotational movement by differential displacement of the two prismatic joints. It provides higher stiffness against mechanical disturbances than a serial configuration. Figure 2.5 shows one PCJM element. The angular offset θ is calculated as follows:


$$\theta = \arctan \left(\frac{L_2 - L_1}{d} \right) \quad (2.1)$$

where L_1 and L_2 are the linear displacements of each joint and $d = 17mm$ the distance from each other.

Screen clipping taken: 2017/06/02 12:49



Screen clipping taken: 2017/06/02 12:50



Parallel Coupled Joint Mechanism

The PCJM element allows translational as well as rotational movement by differential displacement of the two prismatic joints. It provides higher stiffness against mechanical disturbances than a serial configuration. Figure 2.5 shows one PCJM element. The angular offset θ is calculated as follows:

$$\theta = \arctan \left(\frac{L_2 - L_1}{d} \right) \quad (2.1)$$

where L_1 and L_2 are the linear displacements of each joint and $d = 17mm$ the distance from each other.

$$\theta = \arctan \left(\frac{L_2 - L_1}{d} \right)$$

$$\tan \theta = \frac{L_2 - L_1}{d}$$

Furthermore, the mapping from L_i and θ_i of the PCJMs and the configuration of the prismatic and revolute joints from the serial robot is defined as

$$L_2 := q_0, \quad \theta_1 := q_1 \quad (2.9)$$

$$L_4 := q_2, \quad \theta_2 := q_3 \quad (2.10)$$


and

$$L_1 := L_2 - d * \tan(\theta_1) \quad (2.11)$$

$$L_3 := L_4 + d * \tan(\theta_2) \quad (2.12)$$

where the signs differ because of the flipped coordinate frames at the respective revolute joints.

Screen clipping taken: 2017/06/02 12:50



Screen clipping taken: 2017/06/02 13:09



Table 2.1.: The Denavit-Hartenberg parameters for the serial robot.

Link	ϕ_i	d_i	a_i	α_i
1	0	$d_1 + q_0$	0	$-\frac{\pi}{2}$
2	$q_1 - \frac{\pi}{2}$	d_2	l_1	0
3	$\frac{\pi}{2}$	$d_3 + q_2$	l_2	$\frac{\pi}{2}$
4	$q_3 + \frac{\pi}{2}$	0	l_3	$-\frac{\pi}{2}$
5	0	$d_4 + q_4$	0	0
6	q_5	0	0	0

Offsets	mm
l_1	32.5
l_2	1.33
l_3	40.5
d_1	22
d_2	20
d_3	22
d_4	55.35

given as

$$T_0^6 = \begin{pmatrix} -s_1 s_5 - c_1 c_5 s_3 & c_1 s_3 s_5 - c_5 s_1 & -c_1 c_3 & l_2 c_1 + l_1 s_1 - c_1 c_3 (d_4 + q_4) - l_3 c_1 s_3 \\ c_3 c_5 & -c_3 s_5 & -s_3 & d_2 + d_3 + q_2 - s_3 (d_4 + q_4) + l_3 c_3 \\ c_5 s_1 s_3 - c_1 s_5 & -c_1 c_5 - s_1 s_3 s_5 & c_3 s_1 & d_1 + q_0 + l_1 c_1 - l_2 s_1 + c_3 s_1 (d_4 + q_4) + l_3 s_1 s_3 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (2.3)$$

with $s_i := \sin(q_i)$, $c_i := \cos(q_i)$.

Screen clipping taken: 2017/06/02 13:38

vrep_setq([0.00 0.0 0.005 0.5 0.00 0.00],clientID, blocking)

Screen clipping taken: 2017/06/02 13:10