

# harary graph $H_{k,n}$ is $k$ -connected

171240513 黄秉焜

南京大学匡亚明学院

2018 年 11 月 19 日



## 1 Harary Graph

- Definition
- Origin

## 2 Construction Method

## 3 Proof

- Idea of Proof
- Classified Discussion



## 1 Harary Graph

### ■ Definition

### ■ Origin

## 2 Construction Method

## 3 Proof

### ■ Idea of Proof

### ■ Classified Discussion



## Definition (Harary Graph)

The Harary graph  $H_{k,n}$  is a particular example of a  $k$ -connected graph with  $n$  graph vertices having the smallest possible number of edges.



## 1 Harary Graph

- Definition

- Origin

## 2 Construction Method

## 3 Proof

- Idea of Proof

- Classified Discussion



*In the second book on graph theory ever written, Berge' lists 14 unsolved problems, one of which is the following: "11. Quelle est la connexité maximum d'un graphe de  $n$  sommets et de  $m$  arêtes? L'intérêt de ce problème est analogue à celui de trouver le diamètre minimum d'un graphe." The purpose of this note is to solve the problem.*

TO FIND THE MAXIMUM CONNECTIVITY OF A GRAPH



## Theorem

*Among all  $p, q$  graphs, the maximum connectivity is 0 when  $q < p - 1$  and is  $\lceil 2q/p \rceil$  when  $q > p - 1$ .*



## Theorem

*Among all  $p, q$  graphs, the maximum connectivity is 0 when  $q < p - 1$  and is  $\lceil 2q/p \rceil$  when  $q > p - 1$ .*

- The connectivity of a  $p, q$  graph cannot exceed  $\lceil 2q/p \rceil$ .





## Theorem

*Among all  $p, q$  graphs, the maximum connectivity is 0 when  $q < p - 1$  and is  $\lceil 2q/p \rceil$  when  $q > p - 1$ .*

- The connectivity of a  $p, q$  graph cannot exceed  $\lceil 2q/p \rceil$ .  
(Let some vertex be isolated.)



## Theorem

*Among all  $p, q$  graphs, the maximum connectivity is 0 when  $q < p - 1$  and is  $\lceil 2q/p \rceil$  when  $q > p - 1$ .*

- The connectivity of a  $p, q$  graph cannot exceed  $\lceil 2q/p \rceil$ . (Let some vertex be isolated.)
- There exists a  $p, q$  graph  $H$  whose connectivity is  $\lceil 2q/p \rceil$ .



## Theorem

*Among all  $p, q$  graphs, the maximum connectivity is 0 when  $q < p - 1$  and is  $\lceil 2q/p \rceil$  when  $q > p - 1$ .*

- The connectivity of a  $p, q$  graph cannot exceed  $\lceil 2q/p \rceil$ .  
(Let some vertex be isolated.)
- There exists a  $p, q$  graph  $H$  whose connectivity is  $\lceil 2q/p \rceil$ .  
(Construct some graphs that satisfy the conditions——**the Harary Graph**)



## 1 Harary Graph


- Definition
- Origin

## 2 Construction Method

## 3 Proof

- Idea of Proof
- Classified Discussion


- two vertices  $v_i$  and  $v_j$  are linked if and only if  $i - r \leq j \leq i + r$ ;



$k = 2r + 1$  is odd, and  $n$  is even




- $H_{k,n}$  is obtained by joining  $v_i$  and  $v_{i+\frac{n}{2}}$  in  $H_{2r,n}$  for every  $i \in [0, \frac{n}{2} - 1]$ ;



$k = 2r + 1$  and  $n$  are both odd



- $H_{k,n}$  is obtained from  $H_{2r,n}$  by first linking  $v_0$  to both  $v_{\lfloor \frac{n}{2} \rfloor}$  and  $v_{\lceil \frac{n}{2} \rceil}$ , and then each vertex  $v_i$  to  $v_{i+\lceil \frac{n}{2} \rceil}$  for every  $i \in [1, \lfloor \frac{n}{2} \rfloor - 1]$ ;





## 1 Harary Graph

- Definition
- Origin

## 2 Construction Method

## 3 Proof

- Idea of Proof
- Classified Discussion



## 1 Harary Graph

- Definition
- Origin

## 2 Construction Method

## 3 Proof

- Idea of Proof
- Classified Discussion



To prove  $\kappa(H_{k,n}) = k$ . (CZ: Theorem 5.15)

- $\kappa(H_{k,n}) \leq k$
- $\kappa(H_{k,n}) \geq k$

## 1 Harary Graph

- Definition
- Origin

## 2 Construction Method

## 3 Proof

- Idea of Proof
- Classified Discussion



reduction to absurdity:

Suppose that  $V$  is a vertex cut with  $|V| < 2r$ :

- 1 Choose vertices  $i$  and  $j$  in different components of  $H_{2r,n} - V$ .  
Divide the vertices into the following two parts:
  - ▶  $S = \{i, i+1, \dots, j-1, j\}$
  - ▶  $T = \{j, j+1, \dots, i-1, i\}$

reduction to absurdity:

Suppose that  $V$  is a vertex cut with  $|V| < 2r$ :

- 1 Choose vertices  $i$  and  $j$  in different components of  $H_{2r,n} - V$ .  
Divide the vertices into the following two parts:
  - ▶  $S = \{i, i+1, \dots, j-1, j\}$
  - ▶  $T = \{j, j+1, \dots, i-1, i\}$
- 2 Since  $|V| < 2r$ , assume that  $|V \cap S| < r$ .  
Find that in  $S \setminus V$ , the difference between any two consecutive terms is at most  $r$ .



reduction to absurdity:

Suppose that  $V$  is a vertex cut with  $|V| < 2r$ :

- 1 Choose vertices  $i$  and  $j$  in different components of  $H_{2r,n} - V$ .  
Divide the vertices into the following two parts:
  - ▶  $S = \{i, i+1, \dots, j-1, j\}$
  - ▶  $T = \{j, j+1, \dots, i-1, i\}$
- 2 Since  $|V| < 2r$ , assume that  $|V \cap S| < r$ .  
Find that in  $S \setminus V$ , the difference between any two consecutive terms is at most  $r$ .
- 3 Since each vertex of  $H_{2r,n}$  is adjacent with the near  $2r$  vertices, any two consecutive terms in  $S \setminus V$  are always adjacent.

reduction to absurdity:

Suppose that  $V$  is a vertex cut with  $|V| < 2r$ :

- 1 Choose vertices  $i$  and  $j$  in different components of  $H_{2r,n} - V$ .  
Divide the vertices into the following two parts:
  - ▶  $S = \{i, i+1, \dots, j-1, j\}$
  - ▶  $T = \{j, j+1, \dots, i-1, i\}$
- 2 Since  $|V| < 2r$ , assume that  $|V \cap S| < r$ .  
Find that in  $S \setminus V$ , the difference between any two consecutive terms is at most  $r$ .
- 3 Since each vertex of  $H_{2r,n}$  is adjacent with the near  $2r$  vertices, any two consecutive terms in  $S \setminus V$  are always adjacent.
- 4 Thus, we have a path from  $i$  to  $j$ , which is a contradiction.  
i.e.  $\kappa(H_{k,n}) \geq 2r = k$



review the proof just now, we find that:

### Lemma

*In the case of  $H_{2r,n}$ , it is necessary (and sufficient) to remove two separate subsets of  $r$  consecutive vertices each, along the circumference of the polygon.*



$k = 2r + 1$  is odd and  $n$  is even



1 Similar to the case  $k = 2r$ , we have that  $\kappa(H_{k,n}) \geq 2r$ .



- 1 Similar to the case  $k = 2r$ , we have that  $\kappa(H_{k,n}) \geq 2r$ .
- 2 When  $|V| = 2r$ , we must remove  $r$  consecutive vertices from  $S$  according to the above lemma.



- 1 Similar to the case  $k = 2r$ , we have that  $\kappa(H_{k,n}) \geq 2r$ .
- 2 When  $|V| = 2r$ , we must remove  $r$  consecutive vertices from  $S$  according to the above lemma.
- 3 Since we also join pairs of vertices which are diametrically opposite, at least one more vertex must also be removed to break the diametric connection.  
i.e.  $\kappa(H_{k,n}) \geq 2r + 1 = k$

$k = 2r + 1$  and  $n$  are both odd



- 1 The proof is similar to the case that  $k = 2r + 1$  is odd and  $n$  is even.



- 1 The proof is similar to the case that  $k = 2r + 1$  is odd and  $n$  is even.
- 2 In the last case, we need to remove one extra vertex to break the diametric connection.  
While in this case, we need to remove one extra vertex to break the connection between  $V_i$  and  $V_{i+\lfloor \frac{n}{2} \rfloor}$ .
- 3 So, we can also get that  $\kappa(H_{k,n}) \geq 2r + 1 = k$

$k = 2r + 1$  is odd



From the proof, we sense that the construction method can be more universe.

### Lemma

*To construct an  $H_{2r+1,n}$ , we only need to adjoin some random edges in one  $H_{2r,n}$ , so that each vertex in the graph is adjacent some vertex that is not its near  $2r$  vertices.*



- Frank Harary. The maximum connectivity of a graph.
- Gary Chartrand, Ping Zhang. First course in graph theory.
- Olivier Baudon, Julien Bensmail, Eric Sopena. Partitioning Harary graphs into connected subgraphs containing prescribed vertices.
- J. A. Bondy, U. S. R. Murty. Graph theory with application.