


计算机问题求解 — 论题3-14

- 矩阵计算

2016年12月14日

矩阵的逆与线性方程组的解

$$\begin{aligned}a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &= b_1 \\a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &= b_2 \\&\vdots \\a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nn}x_n &= b_n\end{aligned}$$


$$\begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}$$

or, equivalently, letting $A = (a_{ij})$, $x = (x_i)$, and $b = (b_i)$, as


$$Ax = b.$$

If A is nonsingular, it possesses an inverse A^{-1} , and



逆矩阵存在的条件

A square matrix has full rank if and only if it is nonsingular.

A matrix A has full column rank if and only if it does not have a null vector.

A square matrix A is singular if and only if it has a null vector.

An $n \times n$ matrix A is singular if and only if $\det(A) = 0$.

这是什么意思？

问题2:

如何计算非奇异矩阵的逆?

- 1: 矩阵 A 的逆= A 的伴随矩阵/行列式 A 的值
- 2: 矩阵 A 的逆: 对 $(A|E)$ 进行行初等变换得到 $(E|A^{-1})$

例：求3阶方阵 $A = \begin{pmatrix} 2 & 2 & 1 \\ 3 & 1 & 5 \\ 3 & 2 & 3 \end{pmatrix}$ 的逆矩阵.

解： $|A| = 1$, $M_{11} = -7$, $M_{12} = -6$, $M_{13} = 3$,
 $M_{21} = 4$, $M_{22} = 3$, $M_{23} = -2$,
 $M_{31} = 9$, $M_{32} = 7$, $M_{33} = -4$,

则

$$\begin{aligned} A^{-1} &= \frac{1}{|A|} A^* = A^* = \begin{pmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{23} & A_{33} \end{pmatrix} \\ &= \begin{pmatrix} M_{11} & -M_{21} & M_{31} \\ -M_{12} & M_{22} & -M_{32} \\ M_{13} & -M_{23} & M_{33} \end{pmatrix} = \begin{pmatrix} -7 & -4 & 9 \\ 6 & 3 & -7 \\ 3 & 2 & -4 \end{pmatrix} \end{aligned}$$

例 1 设 $A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 2 & 1 \\ 3 & 4 & 3 \end{pmatrix}$, 求 A^{-1} .

解 $(A | E) = \begin{pmatrix} 1 & 2 & 3 & | & 1 & 0 & 0 \\ 2 & 2 & 1 & | & 0 & 1 & 0 \\ 3 & 4 & 3 & | & 0 & 0 & 1 \end{pmatrix}$

$$\begin{array}{l} \underbrace{r_2 - 2r_1} \\ \underbrace{r_3 - 3r_1} \end{array} \begin{pmatrix} 1 & 2 & 3 & | & 1 & 0 & 0 \\ 0 & -2 & -5 & | & -2 & 1 & 0 \\ 0 & -2 & -6 & | & -3 & 0 & 1 \end{pmatrix} \begin{array}{l} \\ \underbrace{r_1 + r_2} \\ \underbrace{r_3 - r_2} \end{array}$$

问题3:

为什么通常不直接用求逆矩阵的办法来解线性方程组？

高斯消元法
过程中可能
出现的现象！

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 4 & 1 \end{bmatrix} X = \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 & -2 \\ 0 & 1 & 4 \\ 0 & 0 & 3 \end{bmatrix} X = \begin{bmatrix} 1 \\ 3 \\ 6 \end{bmatrix}$$

问题4:

三角阵会给解
线性方程组带
来什么便利？

问题5:


三角阵确实会极大简化方程求解，但是多数情况下，我们不会遇到三角阵。

$$Ax = b$$

怎么办？

问题6:

你能否借助右边的图解释一下用LUP分解方法解线性方程组的基本思想? 这个方法的关键在哪里?



$$Ly = Pb$$

$$\begin{aligned} y_1 &= b_{\pi[1]}, \\ l_{21}y_1 + y_2 &= b_{\pi[2]}, \\ l_{31}y_1 + l_{32}y_2 + y_3 &= b_{\pi[3]}, \\ &\vdots \\ l_{n1}y_1 + l_{n2}y_2 + l_{n3}y_3 + \cdots + y_n &= b_{\pi[n]}. \end{aligned}$$

问题7:
 π 数组是什么?

So:
$$y_i = b_{\pi[i]} - \sum_{j=1}^{i-1} l_{ij} y_j .$$

$$Ux = y$$

$$u_{11}x_1 + u_{12}x_2 + \cdots + u_{1,n-2}x_{n-2} + u_{1,n-1}x_{n-1} + u_{1n}x_n = y_1 ,$$

$$u_{22}x_2 + \cdots + u_{2,n-2}x_{n-2} + u_{2,n-1}x_{n-1} + u_{2n}x_n = y_2 ,$$

$$\vdots$$

$$u_{n-2,n-2}x_{n-2} + u_{n-2,n-1}x_{n-1} + u_{n-2,n}x_n = y_{n-2} ,$$

$$u_{n-1,n-1}x_{n-1} + u_{n-1,n}x_n = y_{n-1} ,$$

$$u_{n,n}x_n = y_n .$$

$$x_i = \left(y_i - \sum_{j=i+1}^n u_{ij}x_j \right) / u_{ii} .$$

If we have LUP, we can solve the equations in $\Theta(n^2)$

LUP-SOLVE(L, U, π, b)

```
1   $n = L.rows$ 
2  let  $x$  be a new vector of length  $n$ 
3  for  $i = 1$  to  $n$ 
4       $y_i = b_{\pi[i]} - \sum_{j=1}^{i-1} l_{ij} y_j$ 
5  for  $i = n$  downto 1
6       $x_i = (y_i - \sum_{j=i+1}^n u_{ij} x_j) / u_{ii}$ 
7  return  $x$ 
```

But, how can we get LUP?

问题8：从以下的例子中，我们能观察到什么现象？我们能如何猜想？

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ c/a & 1 \end{bmatrix} \times \begin{bmatrix} a & b \\ 0 & d - cb/a \end{bmatrix}$$

$$\begin{bmatrix} a_{1,1} & a_{1,2} & a_{1,3} & a_{1,4} \\ a_{2,1} & a_{2,2} & a_{2,3} & a_{2,4} \\ a_{3,1} & a_{3,2} & a_{3,3} & a_{3,4} \\ a_{4,1} & a_{4,2} & a_{4,3} & a_{4,4} \end{bmatrix} = \begin{bmatrix} a & B \\ C & D \end{bmatrix} = ?$$

$$\begin{bmatrix} a & B \\ C & D \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ C/a & I \end{bmatrix} \times \begin{bmatrix} a & B \\ 0 & D - CB/a \end{bmatrix}$$

假如可以不考虑 P

$$A = \left(\begin{array}{c|ccc} a_{11} & a_{12} & \cdots & a_{1n} \\ \hline a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{array} \right)$$
$$= \begin{pmatrix} a_{11} & w^T \\ v & A' \end{pmatrix},$$

问题9:

为什么说这是采用了“高斯消去法”？

$$A = \begin{pmatrix} a_{11} & w^T \\ v & A' \end{pmatrix}$$
$$= \begin{pmatrix} 1 & 0 \\ v/a_{11} & I_{n-1} \end{pmatrix} \begin{pmatrix} a_{11} & w^T \\ \textcircled{0} & A' - vw^T/a_{11} \end{pmatrix}$$

问题10:

我们对 如何处理？

问题11:

我们确定能对 $A' - vw^T/a_{11}$ 进行递归处理吗？

We claim that if A is nonsingular, then the Schur complement is nonsingular, too. Why? Suppose that the Schur complement, which is $(n - 1) \times (n - 1)$, is singular. Then by Theorem D.1, it has row rank strictly less than $n - 1$. Because the bottom $n - 1$ entries in the first column of the matrix

$$\begin{pmatrix} a_{11} & w^T \\ 0 & A' - vw^T/a_{11} \end{pmatrix}$$

are all 0, the bottom $n - 1$ rows of this matrix must have row rank strictly less than $n - 1$. The row rank of the entire matrix, therefore, is strictly less than n . Applying Exercise D.2-8 to equation (28.8), A has rank strictly less than n , and from Theorem D.1 we derive the contradiction that A is singular.

递归显然是可行的：

$$\begin{aligned} A &= \begin{pmatrix} 1 & 0 \\ v/a_{11} & I_{n-1} \end{pmatrix} \begin{pmatrix} a_{11} & w^T \\ 0 & A' - vw^T/a_{11} \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ v/a_{11} & I_{n-1} \end{pmatrix} \begin{pmatrix} a_{11} & w^T \\ 0 & L'U' \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ v/a_{11} & L' \end{pmatrix} \begin{pmatrix} a_{11} & w^T \\ 0 & U' \end{pmatrix} \\ &= LU, \end{aligned}$$

$$\begin{pmatrix} 2 & 3 & 1 & 5 \\ 6 & 13 & 5 & 19 \\ 2 & 19 & 10 & 23 \\ 4 & 10 & 11 & 31 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 3 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 2 & 0 & 0 & 1 \end{pmatrix} \times \begin{pmatrix} 2 & 3 & 1 & 5 \\ 0 & 4 & 2 & 4 \\ 0 & 16 & 9 & 18 \\ 0 & 4 & 9 & 21 \end{pmatrix}$$

A

$$\begin{pmatrix} 4 & 2 & 4 \\ 16 & 9 & 18 \\ 4 & 9 & 21 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 4 & 1 & 0 \\ 1 & 7 & 1 \end{pmatrix} \times \begin{pmatrix} 4 & 2 & 4 \\ 0 & 1 & 2 \\ 0 & 0 & 3 \end{pmatrix}$$

$$\begin{pmatrix} 2 & 3 & 1 & 5 \\ 6 & 13 & 5 & 19 \\ 2 & 19 & 10 & 23 \\ 4 & 10 & 11 & 31 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 3 & 1 & 0 & 0 \\ 1 & 4 & 1 & 0 \\ 2 & 1 & 7 & 1 \end{pmatrix} \begin{pmatrix} 2 & 3 & 1 & 5 \\ 0 & 4 & 2 & 4 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 3 \end{pmatrix}$$

A L U

LU-DECOMPOSITION(A)

```
1   $n = A.rows$ 
2  let  $L$  and  $U$  be new  $n \times n$  matrices
3  initialize  $U$  with 0s below the diagonal
4  initialize  $L$  with 1s on the diagonal and 0s above the diagonal
5  for  $k = 1$  to  $n$ 
6       $u_{kk} = a_{kk}$ 
7      for  $i = k + 1$  to  $n$ 
8           $l_{ik} = a_{ik}/u_{kk}$            //  $l_{ik}$  holds  $v_i$ 
9           $u_{ki} = a_{ki}$                //  $u_{ki}$  holds  $w_i^T$ 
10     for  $i = k + 1$  to  $n$ 
11         for  $j = k + 1$  to  $n$ 
12              $a_{ij} = a_{ij} - l_{ik}u_{kj}$ 
13 return  $L$  and  $U$ 
```

问题12:

为什么算法中并没有用递归?

2	3	1	5
6	13	5	19
2	19	10	23
4	10	11	31

(a)

2	3	1	5
3	4	2	4
1	16	9	18
2	4	9	21

(b)

2	3	1	5
3	4	2	4
1	4	1	2
2	1	7	17

(c)

2	3	1	5
3	4	2	4
1	4	1	2
2	1	7	3

(d)

$$\begin{pmatrix} 2 & 3 & 1 & 5 \\ 6 & 13 & 5 & 19 \\ 2 & 19 & 10 & 23 \\ 4 & 10 & 11 & 31 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 3 & 1 & 0 & 0 \\ 1 & 4 & 1 & 0 \\ 2 & 1 & 7 & 1 \end{pmatrix} \begin{pmatrix} 2 & 3 & 1 & 5 \\ 0 & 4 & 2 & 4 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 3 \end{pmatrix}$$

A
 L
 U

(e)

问题13:

下例我们遇到了什么困难?
你有解决思路吗?

$$\begin{bmatrix} 2 & 3 & 1 & 5 \\ 6 & 9 & 5 & 19 \\ 2 & 19 & 10 & 23 \\ 4 & 10 & 11 & 31 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 3 & 1 & 5 \\ 3 & 0 & 2 & 4 \\ 1 & 16 & 9 & 18 \\ 2 & 4 & 9 & 21 \end{bmatrix}$$

问题14:

为什么需要置换矩阵？为什么一定能够找到可置换的行？

用置换矩阵进行主元选择（行初等变换：交换最大元到第一行）

某次递归过程中，如果舒尔补第一列全为0，该矩阵一定奇异，递归前的矩阵一定奇异，进而原矩阵一定奇异

$$QA = \begin{pmatrix} a_{k1} & w^T \\ v & A' \end{pmatrix} \bullet \bullet \bullet$$

初等行变换，你能
写出Q吗？

$$\begin{aligned} QA &= \begin{pmatrix} a_{k1} & w^T \\ v & A' \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ v/a_{k1} & I_{n-1} \end{pmatrix} \begin{pmatrix} a_{k1} & w^T \\ 0 & A' - vw^T/a_{k1} \end{pmatrix} \end{aligned}$$

递归




$$P'(A' - vw^T/a_{k1}) = L'U'$$

$$\begin{aligned} &\begin{pmatrix} 1 & 0 \\ 0 & P' \end{pmatrix} QA \\ &= \begin{pmatrix} 1 & 0 \\ 0 & P' \end{pmatrix} \begin{pmatrix} 1 & 0 \\ v/a_{k1} & I_{n-1} \end{pmatrix} \begin{pmatrix} a_{k1} & w^T \\ 0 & A' - vw^T/a_{k1} \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ P'v/a_{k1} & P' \end{pmatrix} \begin{pmatrix} a_{k1} & w^T \\ 0 & A' - vw^T/a_{k1} \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ P'v/a_{k1} & I_{n-1} \end{pmatrix} \begin{pmatrix} a_{k1} & w^T \\ 0 & P'(A' - vw^T/a_{k1}) \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ P'v/a_{k1} & I_{n-1} \end{pmatrix} \begin{pmatrix} a_{k1} & w^T \\ 0 & L'U' \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ P'v/a_{k1} & L' \end{pmatrix} \begin{pmatrix} a_{k1} & w^T \\ 0 & U' \end{pmatrix} \\ &= LU, \end{aligned}$$

Define

$$P = \begin{pmatrix} 1 & 0 \\ 0 & P' \end{pmatrix} Q$$



PA=LU而且无需担
心除0或者不稳定！

行置换的处理

```
3  for  $i = 1$  to  $n$ 
4       $\pi[i] = i$ 
5  for  $k = 1$  to  $n$ 
6       $p = 0$ 
7      for  $i = k$  to  $n$ 
8          if  $|a_{ik}| > p$ 
9               $p = |a_{ik}|$ 
10              $k' = i$ 
11      if  $p == 0$ 
12          error “singular matrix”
13      exchange  $\pi[k]$  with  $\pi[k']$ 
```

问题15:

如何理解数组pi?

问题16:

算法中并没有出现两个三角矩阵和置换矩阵P，这些矩阵的值是如何体现的？

```
5  for  $k = 1$  to  $n$ 
```

```
    .....
```

```
13  exchange  $\pi[k]$  with  $\pi[k']$ 
```

```
14  for  $i = 1$  to  $n$ 
```

```
15      exchange  $a_{ki}$  with  $a_{k'i}$ 
```

```
16  for  $i = k + 1$  to  $n$ 
```

```
17       $a_{ik} = a_{ik} / a_{kk}$ 
```

```
18      for  $j = k + 1$  to  $n$ 
```

```
19           $a_{ij} = a_{ij} - a_{ik}a_{kj}$ 
```

1	2	0	2	0.6
2	3	3	4	-2
3	5	5	4	2
4	-1	-2	3.4	-1

(a)

3	5	5	4	2
2	3	3	4	-2
1	2	0	2	0.6
4	-1	-2	3.4	-1

(b)

3	5	5	4	2
2	0.6	0	1.6	-3.2
1	0.4	-2	0.4	-0.2
4	-0.2	-1	4.2	-0.6

(c)

3	5	5	4	2
2	0.6	0	1.6	-3.2
1	0.4	-2	0.4	-0.2
4	-0.2	-1	4.2	-0.6

(d)

3	5	5	4	2
1	0.4	-2	0.4	-0.2
2	0.6	0	1.6	-3.2
4	-0.2	-1	4.2	-0.6

(e)

3	5	5	4	2
1	0.4	-2	0.4	-0.2
2	0.6	0	1.6	-3.2
4	-0.2	0.5	4	-0.5

(f)

3	5	5	4	2
1	0.4	-2	0.4	-0.2
2	0.6	0	1.6	-3.2
4	-0.2	0.5	4	-0.5

(g)

3	5	5	4	2
1	0.4	-2	0.4	-0.2
4	-0.2	0.5	4	-0.5
2	0.6	0	1.6	-3.2

(h)

3	5	5	4	2
1	0.4	-2	0.4	-0.2
4	-0.2	0.5	4	-0.5
2	0.6	0	0.4	-3

(i)

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 2 & 0 & 2 & 0.6 \\ 3 & 3 & 4 & -2 \\ 5 & 5 & 4 & 2 \\ -1 & -2 & 3.4 & -1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0.4 & 1 & 0 & 0 \\ -0.2 & 0.5 & 1 & 0 \\ 0.6 & 0 & 0.4 & 1 \end{pmatrix} \begin{pmatrix} 5 & 5 & 4 & 2 \\ 0 & -2 & 0.4 & -0.2 \\ 0 & 0 & 4 & -0.5 \\ 0 & 0 & 0 & -3 \end{pmatrix}$$

P
 A
 L
 U

问题17: 置换矩阵如何获得?

问题18:

你能否解释一下，为什么可以利用**LUP**分解来计算逆矩阵？

In general, once we have the LUP decomposition of A , we can solve, in time $\Theta(k n^2)$, k versions of the equation $Ax = b$ that differ only in b .

We can think of the equation

$$AX = I_n, \tag{28.10}$$

which defines the matrix X , the inverse of A , as a set of n distinct equations of the form $Ax = b$. To be precise, let X_i denote the i th column of X , and recall that the unit vector e_i is the i th column of I_n . We can then solve equation (28.10) for X by using the LUP decomposition for A to solve each equation

$$AX_i = e_i$$

separately for X_i .

问题19:

这有多复杂?

课外作业

- TC Ex.28.1: 2, 3, 6, 7
 - TC Ex.28.2: 1, 2, 3
 - TC Ex.28.3: 1, 3
 - TC Prob.28.1
-

Open topics

- 自学Hill_n加密方法，并向同学们介绍
- 如何用PLU分解求矩阵的行列式？