

MFE 409 LECTURE 2A

MEASURING VALUE-AT-RISK

Valentin Haddad

Spring 2019



LECTURE OBJECTIVES

Measuring Value-at-Risk:

- How to judge validity of a VaR estimate?
- Historical approach
- Model-building approach
- How to get a measure for a given approach but also how to choose an appropriate approach

OUTLINE


1 BACK-TESTING

2 HISTORICAL SIMULATION


BACK-TESTING

- **Back-testing:** How well a current procedure would have performed if applied in the past
 - ▶ Investment strategy
 - ▶ Risk measure
- Our context: How would a method to compute Value-at-Risk would have performed in the past?


S&P500 INDEX, 2012-2013



S&P500 DAILY RETURNS, 2012-2013




A 99% VaR MEASURE



EXCEPTIONS

ANOTHER 99% VaR MEASURE



NUMBER OF EXCEPTIONS

- Say we measure the daily VaR with confidence c
- On a given day:
 - ▶ Probability of exception:
 - ▶ Probability of no exception:

NUMBER OF EXCEPTIONS

- Say we measure the daily VaR with confidence c
- On a given day:
 - ▶ Probability of exception:
 - ▶ Probability of no exception:
- For 99% VaR, a 2-year sample should have on average ? exceptions

NUMBER OF EXCEPTIONS

- Say we measure the daily VaR with confidence c
- On a given day:
 - ▶ Probability of exception:
 - ▶ Probability of no exception:
- For 99% VaR, a 2-year sample should have on average exceptions

DISTRIBUTION OF NUMBER OF EXCEPTIONS

- What if we see 6 exceptions? 11 exceptions?

DISTRIBUTION OF NUMBER OF EXCEPTIONS

- What if we see 6 exceptions? 11 exceptions?
- Probability of observing k exceptions:

$$\frac{n!}{k!(n-k)!} (1-c)^k c^{n-k}$$

DISTRIBUTION OF NUMBER OF EXCEPTIONS

- What if we see 6 exceptions? 11 exceptions?

- Probability of observing k exceptions:

$$\frac{n!}{k!(n-k)!} (1-c)^k c^{n-k}$$

- Probability of observing more (or equal) than m exceptions:

$$\sum_{k=m}^n \frac{n!}{k!(n-k)!} (1-c)^k c^{n-k}$$

DISTRIBUTION OF NUMBER OF EXCEPTIONS

- What if we see 6 exceptions? 11 exceptions?

- Probability of observing k exceptions:

$$\frac{n!}{k!(n-k)!} (1-c)^k c^{n-k}$$

- Probability of observing more (or equal) than m exceptions:

$$\sum_{k=m}^n \frac{n!}{k!(n-k)!} (1-c)^k c^{n-k}$$

- Binomial distribution: $P(\# \text{ exceptions} \geq m) = 1 - F(m-1|n, 1-c)$
 - $F(\cdot|n, p)$ c.d.f. of a binomial with n trials and success probability p


APPLICATION

- 99% - daily VaR
- 2 years: 502 daily returns
- Probability of 6 or more exceptions:
- Probability of 11 or more exceptions:

APPLICATION

- 99% - daily VaR
- 2 years: 502 daily returns
- Probability of 6 or more exceptions:
- Probability of 11 or more exceptions:

ANOTHER VAR MEASURE



OTHER TESTS

- Probability of observing less (or equal) than m exceptions:

$$\sum_{k=0}^m \frac{n!}{k!(n-k)!} (1-c)^k c^{n-k}$$
$$= F(m|n, 1-c)$$

OTHER TESTS

- Probability of observing less (or equal) than m exceptions:


$$\sum_{k=0}^m \frac{n!}{k!(n-k)!} (1-c)^k c^{n-k} \\ = F(m|n, 1-c)$$

- Two sided test (for large n):

$$-2 \ln [c^{n-m} (1-c)^m] + 2 \ln [(1-m/n)^{n-m} (m/n)^m] \sim \chi^2(1)$$

- ▶ Chi-squared 5% threshold: 3.84

BUNCHING



BUNCHING

OUTLINE

1 BACK-TESTING

2 HISTORICAL SIMULATION

HISTORICAL SIMULATION

- Assume the future will be drawn from the same distribution as the past
- Past data reveals the future distribution

HISTORICAL SIMULATION


- Assume the future will be drawn from the same distribution as the past
- Past data reveals the future distribution
- Formally, to compute the daily value-at-risk at confidence level c :
 - ▶ You have n past observations of daily **returns**
 - ▶ Assume the next return will be any of these draws with probability $1/n$

HISTORICAL SIMULATION

- Assume the future will be drawn from the same distribution as the past
- Past data reveals the future distribution
- Formally, to compute the daily value-at-risk at confidence level c :
 - ▶ You have n past observations of daily **returns**
 - ▶ Assume the next return will be any of these draws with probability $1/n$
 - ▶ The VaR corresponds to the loss in the $[(1 - c) \times n]$ -th worst past realization
 - ★ if not integer, round up

EXAMPLE

- Assume we are 04/11/2017
- You have \$4m invested in S&P500, \$5m in NASDAQ Composite, \$1m in DJIA
- You know the value of the indices for the last 3 years (file *indices.xls*)



- What is your 1-day 99% VaR?

CONSTRUCTING THE VAR

CONSTRUCTING THE VAR

CONSTRUCTING THE VAR

EXPECTED SHORTFALL

- Can use the same method to compute expected shortfall

EXPECTED SHORTFALL

- Can use the same method to compute expected shortfall
- Average of the $[(1 - c) \times n]$ worst realizations
 - ▶ Still round up

STRESSED VaR

- Stressed VaR (or ES): VaR (or ES) for the worst consecutive 251-day period in the historical sample

STRESSED VaR

- Stressed VaR (or ES): VaR (or ES) for the worst consecutive 251-day period in the historical sample
- Introduced by regulators to capture the idea that some periods are worse than others
- (Stressed VaR) \geq VaR ?

ACCURACY OF VaR

- If you backtest historical VaR, you find exactly ? deviations

ACCURACY OF VaR

- If you backtest historical VaR, you find exactly c deviations
- But if you had the true VaR, you would sometimes find more, sometimes find less: historical VaR is not perfectly accurate
- Standard error of the estimate:

$$\frac{1}{f(x)} \sqrt{\frac{c(1-c)}{n}}$$

- ▶ $f(x)$: p.d.f. at quantile c
- ▶ Need to know distribution!

EXAMPLE: ACCURACY OF VAR

- Back to portfolio example
- Historical VaR: \$249,000

EXAMPLE: ACCURACY OF VaR

- Back to portfolio example
- Historical VaR: \$249,000
- Approximate by a normal (in \$000s): mean 4, standard deviation 87

$$x = \mu - \sigma \Phi^{-1}(0.01) = -198.4$$

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right) = 3.06 \times 10^{-4}$$

$$\text{StdDev}(\text{VaR}) = \frac{1}{f(x)} \sqrt{\frac{0.99 \times 0.01}{753}} = 12$$

- 95% confidence interval for the VaR is between \$229,000 and \$269,000 → not that precise

BOOTSTRAP

- **Bootstrap:**

BOOTSTRAP

- **Bootstrap:** Draw samples from historical data to understand behavior of statistics

BOOTSTRAP

- **Bootstrap:** Draw samples from historical data to understand behavior of statistics
- Suppose there are 500 daily changes and you want to calculate a 95% confidence interval for VaR
 - ① Sample 500,000 times with replacement from daily changes to obtain 1000 sets of changes over 500 days
 - ② Calculate VaR for each set
 - ③ Calculate a confidence interval by taking the range between the 2.5% lowest and 97.5% largest value

HOW MUCH HISTORICAL DATA?

- Portfolio example used 3 years of data
- How much data would you like to use?

HOW MUCH HISTORICAL DATA?

- Portfolio example used 3 years of data
- How much data would you like to use?
- More data, more precise estimates

HOW MUCH HISTORICAL DATA?

- Portfolio example used 3 years of data
- How much data would you like to use?
- More data, more precise estimates
- But “future same as past” less likely to be true

WEIGHTING OF OBSERVATIONS

- Use as much data as possible, but put more weight on recent data
- Weight observations with an exponential decay as you go back in time.
- Observation i receives weight:

$$\lambda^{n-i} \frac{1 - \lambda}{1 - \lambda^n}$$

- Sort observations, VaR is the scenario just over $1 - c$ cumulative weight

PORTFOLIO EXAMPLE WITH WEIGHTING

- $\lambda = 0.995$

PORTFOLIO EXAMPLE WITH WEIGHTING

- $\lambda = 0.995$

ESTIMATING THE TAIL

- Extreme tail estimated imprecisely with historical method: 99.9% would need multiple thousands of observations
- To get more precise estimates, make assumptions about the shape of the distribution
- Model the whole distribution, e.g. normal distribution
 - ▶ VaR depends of σ
 - ▶ Every observation helps estimate σ

ESTIMATING THE TAIL

- Extreme tail estimated imprecisely with historical method: 99.9% would need multiple thousands of observations
- To get more precise estimates, make assumptions about the shape of the distribution
- Model the whole distribution, e.g. normal distribution
 - ▶ VaR depends of σ
 - ▶ Every observation helps estimate σ
- Model the left tail of the distribution, e.g. using a power law
 - ▶ VaR depends of the shape of the left tail of the distribution
 - ▶ Every tail observation helps estimate the shape of the left tail

ESTIMATING THE TAIL

- Extreme tail estimated imprecisely with historical method: 99.9% would need multiple thousands of observations
- To get more precise estimates, make assumptions about the shape of the distribution
- Model the whole distribution, e.g. normal distribution
 - ▶ VaR depends of σ
 - ▶ Every observation helps estimate σ
- Model the left tail of the distribution, e.g. using a power law
 - ▶ VaR depends of the shape of the left tail of the distribution
 - ▶ Every tail observation helps estimate the shape of the left tail
- **Extreme value theory**: this approach is valid for many distributions

POWER LAW

- **Power law:** X follows a power law, with

$$\text{Prob}(X > x) = Kx^{-1/\xi}$$

- ▶ Also called Pareto distribution
- ▶ $\xi < 1$ controls thickness of tail: low ξ , thin tail


POWER LAW

- **Power law:** X follows a power law, with

$$\text{Prob}(X > x) = Kx^{-1/\xi}$$

- ▶ Also called Pareto distribution
 - ▶ $\xi < 1$ controls thickness of tail: low ξ , thin tail
-
- Regress $\log[\text{Prob}(X > x)]$ on $\log(x)$: slope $-1/\xi$
 - ▶ In historical distribution: $\text{Prob}(X > x_i) = \text{rank}(x_i)/n$

LOG-LOG PLOT FOR PORTFOLIO LOSS



■ Slope: -3 , $\xi = 1/3$

EXTREME VALUE THEORY

- Key result: a wide range of probability distributions have common properties in the tail

EXTREME VALUE THEORY

- Key result: a wide range of probability distributions have common properties in the tail

- Tail distribution:

$$F_u(y) = \frac{F(u+y) - F(u)}{1 - F(u)}$$

- Result: as u becomes large, $F_u(y)$ converges to a generalized Pareto distribution:

$$G_{\xi,\beta}(y) = 1 - \left[1 + \xi \frac{y}{\beta}\right]^{-1/\xi}$$

EXTREME VALUE THEORY

- Key result: a wide range of probability distributions have common properties in the tail

- Tail distribution:

$$F_u(y) = \frac{F(u+y) - F(u)}{1 - F(u)}$$

- Result: as u becomes large, $F_u(y)$ converges to a generalized Pareto distribution:

$$G_{\xi,\beta}(y) = 1 - \left[1 + \xi \frac{y}{\beta}\right]^{-1/\xi}$$

- **Model of right tail!** Remember to find the c -th quantile of losses

ESTIMATING THE POWER LAW

- Partial distribution function:

$$g_{\xi,\beta}(y) = \frac{1}{\beta} \left(1 + \frac{\xi y}{\beta} \right)^{-1/\xi-1}$$

- Choose u : typically 95th percentile of historical distribution

ESTIMATING THE POWER LAW

- Partial distribution function:

$$g_{\xi,\beta}(y) = \frac{1}{\beta} \left(1 + \frac{\xi y}{\beta} \right)^{-1/\xi-1}$$

- Choose u : typically 95th percentile of historical distribution
- Maximize log likelihood:

$$\max_{\xi,\beta} \sum_{i \in tail} \ln [g_{\xi,\beta}(v_i - u)]$$

VAR AND ES FOR A POWER LAW

- Probability distribution:

$$\text{Prob}(\text{Loss} > V) = \underbrace{[1 - F(u)]}_{n_u/n} [1 - G_{\xi, \beta}(V - u)]$$

VaR AND ES FOR A POWER LAW

- Probability distribution:

$$\text{Prob}(\text{Loss} > V) = \underbrace{[1 - F(u)]}_{n_u/n} [1 - G_{\xi, \beta}(V - u)]$$

- V is VaR if this is $1 - c$

$$\text{VaR} = u + \frac{\beta}{\xi} \left(\left[\frac{n}{n_u} c \right]^{-\xi} - 1 \right)$$

VaR AND ES FOR A POWER LAW

- Probability distribution:

$$\text{Prob}(\text{Loss} > V) = \underbrace{[1 - F(u)]}_{n_u/n} [1 - G_{\xi, \beta}(V - u)]$$

- V is VaR if this is $1 - c$

$$\text{VaR} = u + \frac{\beta}{\xi} \left(\left[\frac{n}{n_u} c \right]^{-\xi} - 1 \right)$$

- Can also obtain ES:

$$\text{ES} = \frac{\text{VaR} + \beta - \xi u}{1 - \xi}$$