

# Lecture 3

## Autocorrelation

Lars A. Lochstoer

UCLA Anderson School of Management

Winter 2019

# Overview of Lecture 3

## Autocorrelation

### 1 Introduction to autocorrelations

- ▶ The autocorrelation function

### 2 The Ljung-Box Q-test

# Correlation

## Definition

The **correlation** between two random variables  $X$  and  $Y$  is defined as

$$\rho_{x,y} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

- $\rho_{x,y}$  is known as Pearson's correlation
- measures linear dependence
- bounded between  $-1$  and  $1$
- two variables are uncorrelated if  $\rho_{x,y} = 0$ , perfectly (negatively) correlated if  $\rho_{x,y} = 1$  ( $\rho_{x,y} = -1$ )
- if  $X$  and  $Y$  are random normal variables, then  $\rho_{x,y} = 0$  if and only if  $X$  and  $Y$  are independent

# Sample Correlation

## Definition

The **sample correlation** between two random variables  $X$  and  $Y$  is:

$$\hat{\rho}_{x,y} = \frac{\sum_{t=1}^T (x_t - \bar{x})(y_t - \bar{y})}{\sqrt{\sum_{t=1}^T (x_t - \bar{x})^2 \sum_{t=1}^T (y_t - \bar{y})^2}}$$

where  $\bar{x}$  and  $\bar{y}$  are the sample means.

- this is **not** a regression coefficient
- $\hat{\rho}_{x,y}$  consistently estimates  $\rho_{x,y}$
- $\hat{\rho}_{x,y}$  is built from **method of moments** estimators

# Autocorrelation

## Definition

The **autocorrelation** for a series  $\{r_t\}$  is defined as:

$$\rho_j = \frac{\text{Cov}(r_t, r_{t-j})}{\sqrt{\text{Var}(r_t) \text{Var}(r_{t-j})}} = \frac{\text{Cov}(r_t, r_{t-j})}{\text{Var}(r_t)} = \frac{\gamma_j}{\gamma_0}.$$

- a covariance-stationary series  $r_t$  is not serially correlated if  $\rho_j = 0$  for all  $j$
- autocorrelations are a key signature of the dynamics of the time series you're interested in modeling

# 1st Order Autocorrelation

## Definition

The **sample autocorrelation** for a series  $\{x_t\}$  is:


$$\hat{\rho}_1 = \frac{\sum_{t=2}^T (x_t - \bar{x})(x_{t-1} - \bar{x})}{\sum_{t=1}^T (x_t - \bar{x})^2}, 0 \leq j \leq T - 1$$

where  $\bar{x}$  are the sample means.

- under some conditions,  $\hat{\rho}_1$  is a consistent estimator of  $\rho_1$
- $\hat{\rho}_1$  is asymptotically normal with mean zero and variance  $(1/T)$  if  $\{x_t\}$  are independently and identically distributed over time.
- to test  $H_0 : \rho_1 = 0$ , use t-stat :


$$t = \sqrt{T} \hat{\rho}_1$$

# Autocorrelation in stock returns?



Scatter plot of monthly log returns (VW-CRSP) 1925-2013.

# Autocorrelation in Real Estate Returns?



Scatter plot for Monthly log House Price Changes in AZ. Case-Shiller Index.  
1987.1-2013.10



# Higher-order Autocorrelations

## Definition

The **sample autocorrelation** for a series  $\{x_t\}$  at lag  $j$  is:

$$\hat{\rho}_j = \frac{\sum_{t=j+1}^T (x_t - \bar{x})(x_{t-j} - \bar{x})}{\sum_{t=1}^T (x_t - \bar{x})^2}, 0 \leq j \leq T-1$$

where  $\bar{x}$  are the sample means.

# Autocorrelation

- ① financial time series: e.g. stock returns and housing returns
  - ▶ financial returns tend to be only very weakly autocorrelated [if markets are fairly efficient and liquid]
  - ▶ strong autocorrelations in returns would create huge profit opportunities!
- ② macroeconomic time series: e.g. GDP growth rates
  - ▶ macroeconomic time series have growth rates that are highly autocorrelated
  - ▶ macroeconomic shocks tend to have very persistent effects (e.g. think about the effect of the subprime crisis on GDP growth rates)


# The Autocorrelation Function (ACF)

## Definition

The **sample autocorrelation function** (ACF) of a time series is defined as


$$\hat{\rho}_1, \hat{\rho}_2, \dots, \hat{\rho}_k, \dots,$$

# Autocorrelation in Quarterly GDP Growth




Autocorrelation Function for Quarterly U.S. GDP growth. Two standard error bands around zero. 1947.I-2012.IV

# Autocorrelation in Monthly Inflation




Autocorrelation Function for Monthly U.S. Inflation. 1950-2007.  $\hat{\rho}_1 = 0.52$ .

# Autocorrelation of Annual Log Stock Returns




Autocorrelation Function for Annual log Returns on VW-CRSP Index. Two standard error bands around zero. 1926-2012 .  $\hat{\rho}_1 = 0.05$ .

# Autocorrelation of Monthly Log Stock Returns



Autocorrelation Function for Monthly log Returns on VW-CRSP Index. Two standard error bands around zero. 1926-2012 .  $\hat{\rho}_1 = 0.088$ .


# Autocorrelation of Monthly Log Stock Returns



Autocorrelation Function for Monthly log Returns on EW-CRSP Index (equal weighted). Two standard error bands around zero. 1926-2012 .  $\hat{\rho}_1 = 0.12$ .




# Autocorrelation of Monthly Log Stock Returns



Autocorrelation Function for Monthly log Returns on VW-CRSP Index. Two standard error bands around zero. 1990-2012.  $\hat{\rho}_1 = 0.09$ .

# Autocorrelation of Monthly Log Stock Returns




Autocorrelation Function for Monthly log Returns on EW-CRSP Index (equal weighted). Two standard error bands around zero. 1990-2012.  $\hat{\rho}_1 = 0.17$ .

# Monthly ACF for the Stock Market


- some positive autocorrelation in stock returns (for the market) at the one-month horizon
- autocorrelation is stronger for small stocks (see EW-CRSP)
  - ▶  $\hat{\rho}_1$  varies between 0.09 (VW) and 0.17 (EW) on the short sample
  - ▶  $\hat{\rho}_1$  varies between 0.08 (VW) and 0.12 (EW) on the long sample

# Autocorrelation of Daily Stock Returns




Autocorrelation Function for Daily log Returns on VW-CRSP Index (value-weighted). Two standard error bands around zero. 1926-2012.  $\hat{\rho}_1 = 0.07$ .

# Autocorrelation of Daily Stock Returns




Autocorrelation Function for Daily log Returns on EW-CRSP Index (equal-weighted). Two standard error bands around zero. 1926-2012.  $\hat{\rho}_1 = 0.21$ .

# Autocorrelation of Daily Stock Returns



Autocorrelation Function for Daily log Returns on VW-CRSP Index. Two standard error bands around zero. 1990-2012 .  $\hat{\rho}_1 = 0.01$ .

# Autocorrelation of Daily Stock Returns



Autocorrelation Function for Daily log Returns on EW-CRSP Index. Two standard error bands around zero. 1990-2012 .  $\hat{\rho}_1 = 0.13$ .

# Daily ACF for the Stock Market


- some positive autocorrelation in stock returns (for the market) at the one-day horizon, but only for small stocks
  - ▶  $\hat{\rho}_1$  varies between  $-0.01$  (VW) and  $0.13$  (EW) on the short sample
  - ▶  $\hat{\rho}_1$  varies between  $0.07$  (VW) and  $0.21$  (EW) on the long sample



# Benchmark Model of Portfolio Theory

- In the benchmark model of portfolio theory, returns are assumed to be independently and identically distributed (i.i.d.) over time.
  - ▶ If returns are i.i.d., the variance grows linearly in the investment horizon
  - ▶ The investor's horizon turns out to be irrelevant for optimal portfolio allocation.
- Not quite true in the data

# Autocorrelation of Monthly Log House Price Changes



Autocorrelation Function for Monthly log House Price Changes. Two standard error bands around zero. 1987-2013

# Testing for autocorrelation: Ljung and Box (1978)

## Definition



The **Ljung-Box** statistic tests the null that  $H_0 : \rho_1 = \dots = \rho_m = 0$

$$Q(m) = T(T+2) \sum_{i=1}^m \frac{\hat{\rho}_i^2}{T-i}$$

$Q(m)$  is asymptotically  $\chi^2$  with  $m$  degrees of freedom.

- reject the null if  $Q(m) > \chi^2(\alpha)$  where  $\chi^2(\alpha)$  denotes the  $(1 - \alpha) \times 100$ -th percentile

# Q-test on Monthly Returns



Q-test for Monthly log Returns on VW-CRSP Index. 1926-2012

# Monthly Stock Returns

- choice of  $m$  matters: rule of thumb  $m = \ln(T)$
- if we use this rule of thumb,  $m = 6$  and  $Q(6) = 26$  and p-value is  $1.9e - 4$
- in any case, we reject the null that there is no autocorrelation in monthly U.S. stock returns for all holding periods considered..
- even though these autocorrelations are small, they're measured rather precisely, allowing us to reject the null.

# White Noise

## Definition

A time series  $\varepsilon_t$  is said to be **white noise** if  $\{\varepsilon_t\}$  is a sequence of independent and identically distributed random variables.

Notation:  $\varepsilon_t \sim \text{WN}(0, \sigma_\varepsilon^2)$

If  $\varepsilon_t$  is white noise + normally distributed with mean zero and variance  $\sigma^2$ , then it is called **Gaussian white noise**.

Notation:  $\varepsilon_t \sim \text{GWN}(0, \sigma_\varepsilon^2)$

- There is no autocorrelation. ACF's are all zero