

Week 8 - Linear Algebra

A YouTube playlist with all the videos for this lab sheet can be found [here](#).

1. Use Sage to solve the following system of equations:

$$\begin{cases} 10x + 2y = 0 \\ 2x - y = 154 \end{cases}$$

2. **TICKABLE** Note that the above system of equations is equivalent to the following systems of equations:

$$\begin{cases} 10a + 2b = 0 \\ 2a - b = 154 \end{cases}$$

$$\begin{cases} 10m + 2n = 0 \\ 2m - n = 154 \end{cases}$$

In essence the only thing that defines the system of equations is the coefficients:

$$\begin{pmatrix} 10, 2, 0 \\ 2, -1, 154 \end{pmatrix}$$

We can of course separate the right hand side of our equation and perhaps include those elements in a vector. Our system can now be represented as:

$$\begin{pmatrix} 10, 2 \\ 2, -1 \end{pmatrix} \begin{pmatrix} 0 \\ 154 \end{pmatrix}$$

Let us attempt to represent the above system in Sage.

The following defines b as a vector:

```
b = vector(0,154)
```

The representation of coefficients is a well defined mathematical object called a matrix. The following code defines A as a matrix:

```
A = matrix([[10, 2], [2, -1]])
```

If we define a vector X as a vector of the symbolic variables:

```
X = vector([x, y])
```

We can **multiply** A by X:

```
A * X
```

Verify that $X = (x_0, y_0)$ where (x_0, y_0) is the solution to our system of equations (obtained in Q1).

[Video hint](#)

3. **TICKABLE** In linear algebra (you will study this next semester) a matrix equation is an equation of the form:

$$AX = b$$

or

$$XA = b$$

If we define A and b as in question 2 we can solve this equation quite simply using the `solve_right` or `solve_left` methods. The following obtains a solution to the equation $AX = b$:

`A.solve_right(b)`

Note that `A\b` is shorthand for `A.solve_right`

Use the above to solve the following system of equations using matrix notation:

$$\begin{cases} 4x - 2y + 3z = 10 \\ -x - 5y - 8z = 9 \\ x + y + z = 1 \end{cases}$$

[Video hint](#)

4. For reasons that will become clear, the following definition of matrix multiplication is required:

$$(AB)_{ij} = \sum_{j'} \sum_{i'} A_{ij'} B_{i'j}$$

For 2×2 matrices this is equivalent to:

$$AB = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} = \begin{pmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{pmatrix}$$

As an example create the following two matrices in Sage:

```
A = matrix([[1, 2], [3, 4]])
B = matrix([[7, 8], [9, 10]])
```

Attempt to multiply these matrices by hand and carry out their multiplication in Sage:

`A * B`

Repeat the exercise by multiplying the following pairs of matrices:

1. $A = \begin{pmatrix} -1 & 1 \\ -1 & -1 \end{pmatrix}, B = \begin{pmatrix} -1 & 4 \\ 1 & 1 \end{pmatrix}$
2. $A = \begin{pmatrix} 0 & 144 \\ -2 & 1 \end{pmatrix}, B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$
3. $A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, B = \begin{pmatrix} -2 & 0 \\ -1 & -17 \end{pmatrix}$
4. $A = \begin{pmatrix} 0 & -1 \\ 3 & 1 \end{pmatrix}, B = \begin{pmatrix} 1/3 & 1/3 \\ -1 & 0 \end{pmatrix}$

[Video hint](#)

5. **TICKABLE** The previous exercise shows that when considering matrix multiplication there exists a matrix which does not have a multiplicative affect: “the identity matrix”.

The identity matrix of size $n \times n$ is denoted by \mathbb{I}_n . The following Sage code gives \mathbb{I}_n :

```
identity_matrix(4)
```

Note also, that the previous exercise showed that we can sometimes find a matrix B such that $AB = \mathbb{I}_n$. Finding such a matrix is referred to as ‘inverting’ A and if certain properties hold (you will see this in further details next semester) this matrix is denoted A^{-1} .

If we recall the matrix equation $AX = b$ and if we assume that A^{-1} exists then multiplying both sides by A^{-1} gives:

$$A^{-1}AX = A^{-1}b \Rightarrow \mathbb{I}_n X = A^{-1}b \Rightarrow X = A^{-1}b$$

In Sage we can obtain A^{-1} (if it exists) with the following code:

```
A.inverse()
```

Thus another approach to solving $AX = b$ is:

`A.inverse() * b`

Use this approach to solve the systems of equations we have considered so far.

[Video hint](#)

6. **TICKABLE** Recalling your basic python knowledge. Create a function that returns the inverse of a matrix if it exists (return False if it does not). Do this using the try and except statements that you will need to research.

Using this obtain a list of the inverses of the following matrices:

$$\begin{pmatrix} \frac{1}{2} & 0 & 0 & -1 & 1 \\ -1 & -1 & 1 & -\frac{1}{2} & 2 \\ 0 & -1 & 0 & -2 & 0 \\ 0 & 0 & \frac{1}{2} & -1 & 0 \\ -1 & 0 & -2 & 2 & 0 \end{pmatrix}$$
$$\begin{pmatrix} -1 & -1 & 0 & 0 & -1 \\ 2 & 1 & 0 & 1 & 1 \\ -2 & 0 & 1 & 2 & 2 \\ -\frac{1}{2} & 0 & -\frac{1}{2} & 0 & \frac{1}{2} \\ 0 & 0 & 0 & \frac{1}{2} & -1 \end{pmatrix}$$
$$\begin{pmatrix} -\frac{1}{2} & -\frac{1}{2} \\ -2 & -1 \end{pmatrix}$$
$$\begin{pmatrix} 2 & -2 & 1 \\ 6 & -1 & 1 \\ 12 & -2 & 2 \end{pmatrix}$$
$$\begin{pmatrix} 1 & 2 \\ 2 & 0 \end{pmatrix}$$

For all the above matrices, find the output of the `.det` method. This gives the **determinant** of the matrices. It is a very important quantity that will be explained next semester.

[Video hint](#)

7. **TICKABLE** The `random_matrix` command can be used to obtain a random matrix:

`random_matrix(ZZ, 5)` # Gives a random square matrix of size 5 in the Integers.
`random_matrix(QQ, 5)` # Gives a random square matrix of size 5 in the Rationals.

Using this, attempt to conjecture a connection between the determinant of a matrix and it's inverse (and the determinant of it's inverse).

[Video hint](#)

8. **TICKABLE** The file [W08_D01.csv](#) contains 4 columns of data:

a, b, c, d, f, g

For each row of data, obtain the solution to the system of equations:

$$\begin{cases} ax + by = c \\ dx + fy = g \end{cases}$$

Write to file a new data set containing the following columns:

A, B, C, D

Where A is the number of the original data set, B and C are the solutions to the system of equation in question: $B = x, C = y$. D is the 'norm of the solution vector': $D = \sqrt{C^2 + B^2}$.

If there is no solution to the system of equations set $B=C=D=False$. The data set is a randomly sampled set of problems, how often does a solution exist?

9. The file [W08_D02.csv](#) contains a large number of columns and rows. Investigate the dimensions and plot methods on this matrix.