

Brief Report

LYMPHOID NEOPLASIA

Recurrent *CDKN1B* (p27) mutations in hairy cell leukemia

Sascha Dietrich,¹⁻³ Jennifer Hüllein,³ Stanley Chun-Wei Lee,⁴ Barbara Hutter,⁵ David Gonzalez,⁶ Sandrine Jayne,⁷ Martin J. S. Dyer,⁷ Małgorzata Oleś,² Monica Else,⁸ Xiyang Liu,³ Mikołaj Słabicki,³ Bian Wu,³ Xavier Troussard,⁹ Jan Dürig,¹⁰ Mindaugas Andriulis,¹¹ Claire Dearden,¹² Christof von Kalle,^{3,13} Martin Granzow,¹⁴ Anna Jauch,¹⁴ Stefan Fröhling,³ Wolfgang Huber,² Manja Meggendorfer,¹⁵ Torsten Haferlach,¹⁵ Anthony D. Ho,¹ Daniela Richter,³ Benedikt Brors,⁵ Hanno Glimm,³ Estella Matutes,¹² Omar Abdel Wahab,⁴ and Thorsten Zenz^{1,3}

¹Department of Medicine V, University Hospital Heidelberg, ²Genome Biology Unit, European Molecular Biology Laboratory, and ³Department of Translational Oncology, National Center for Tumor Diseases and German Cancer Research Center (DKFZ), Heidelberg, Germany; ⁴Human Oncology and Pathogenesis Program, Leukemia Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY; ⁵Division Applied Bioinformatics, German Cancer Research Center (DKFZ), Heidelberg, Germany; ⁶Centre for Molecular Pathology, Royal Marsden NHS Foundation Trust, London, United Kingdom; ⁷Ernest and Helen Scott Hematological Research Institute, University of Leicester, Leicester, United Kingdom; ⁸Division of Molecular Pathology, The Institute of Cancer Research, London, United Kingdom; ⁹Laboratoire d'Hématologie, CHU Côte de Nacre, Caen, France; ¹⁰Department of Hematology, University Hospital, Essen, Germany; ¹¹Department of General Pathology, Institute of Pathology, University of Heidelberg, Heidelberg, Germany; ¹²Royal Marsden Hospital, London, United Kingdom; ¹³Heidelberg Center for Personalized Oncology (DKFZ-HIPO), German Cancer Research Center (DKFZ), Heidelberg, Germany; ¹⁴Institute of Human Genetics, University Hospital Heidelberg, Heidelberg, Germany; and ¹⁵Munich Leukemia Laboratory, Munich, Germany

Key Points

- Somatic *CDKN1B* (p27) mutations were identified in 16% (13/81) of HCL patients and coexist with *BRAFV600E* mutations.
- *CDKN1B* is the second most common mutated gene in HCL implicating altered cell cycle regulation and/or senescence in HCL.

Hairy cell leukemia (HCL) is marked by near 100% mutational frequency of *BRAFV600E* mutations. Recurrent cooperating genetic events that may contribute to HCL pathogenesis or affect the clinical course of HCL are currently not described. Therefore, we performed whole exome sequencing to explore the mutational landscape of purine analog refractory HCL. In addition to the disease-defining *BRAFV600E* mutations, we identified mutations in *EZH2*, *ARID1A*, and recurrent inactivating mutations of the cell cycle inhibitor *CDKN1B* (p27). Targeted deep sequencing of *CDKN1B* in a larger cohort of HCL patients identify deleterious *CDKN1B* mutations in 16% of patients with HCL (n = 13 of 81). In 11 of 13 patients the *CDKN1B* mutation was clonal, implying an early role of *CDKN1B* mutations in the pathogenesis of HCL. *CDKN1B* mutations were not found to impact clinical characteristics or outcome in this cohort. These data identify HCL as having the highest frequency of *CDKN1B* mutations among cancers and identify *CDKN1B* as the second most common mutated gene in HCL. Moreover, given the known function of *CDKN1B*, these data suggest a novel role for alterations in regulation of cell cycle and senescence in HCL with *CDKN1B* mutations. (*Blood*. 2015;126(8):1005-1008)

Introduction

Hairy-cell leukemia (HCL) is a rare, mature B-cell malignancy presenting with slow progressing pancytopenia and splenomegaly. Classical HCL is successfully treated with chemotherapy, but eradication of minimal residual disease is rarely achieved.¹ Standard treatment fails in a minority of patients, with a potentially fatal outcome.

Gain-of-function mutations of the BRAF serine/threonine protein kinase (*BRAFV600E*) have been identified in nearly all cases of classical HCL, and mitogen-activated protein kinase signaling is considered the key oncogenic pathway in HCL.² Chung et al³ recently identified hematopoietic stem cells as the cell of origin of HCL by demonstrating that hematopoietic stem cells, and subsequently cells along the hematopoietic hierarchy, contain mutated *BRAF*. Currently, however, no other recurrently mutated genes are known to coexist with *BRAFV600E* mutations in HCL. It is unclear if *BRAFV600E* mutations

alone are sufficient to induce HCL. Moreover, it is also not known if additional mutations may be acquired in *BRAFV600E*-mutant HCL cells, resulting in acquired resistance to therapies commonly administered to patients with HCL such as purine analogs. Therefore, we performed whole-exome sequencing (WES) in 3 HCL patients who were refractory to purine analog treatment and received the BRAF inhibitor (BRAFi) vemurafenib followed by recurrence testing of novel mutations in a larger cohort of HCL patients.

Study design

Clinical samples were provided by S.D., A.D.H., and T.Z. (n = 10), D.G. and E.M. (n = 50), T.H. (n = 17), M.J.S.D. (n = 1), J.D. (n = 2), and X.T. (n = 6),

Submitted April 30, 2015; accepted June 2, 2015. Prepublished online as *Blood* First Edition paper, June 11, 2015; DOI 10.1182/blood-2015-04-643361.

The online version of this article contains a data supplement.

There is an Inside *Blood* Commentary on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2015 by The American Society of Hematology




Figure 1. Recurrent CDKN1B mutations in classical HCL. (A) Cancer consensus and COSMIC annotated gene mutations identified in HCL by whole exome sequencing. Patient 3 progressed after vemurafenib treatment and was sequenced before and after BRAFi (vemurafenib) treatment. Exome sequencing identified recurrent inactivating somatic mutations of *CDKN1B* (for complete list of somatic mutations see supplemental Table 1b). (B) Gene regions of *CDKN1B* and distribution of mutations in HCL. Binding regions of important interaction partners are shown below the gene diagram. There were 17 *CDKN1B* mutations identified in 13 patients. In cases in which normal material was not available for the remaining mutations with sufficient read coverage (mean = 1259 reads, supplemental Table 1) the allele frequencies indicated somatic origin (supplemental Figure 2). (C) Frequency of *CDKN1B* mutations across cancer entities. Mutation frequencies were obtained from <http://www.cbioportal.org/public-portal/>.¹⁵ HCL shows the highest *CDKN1B* mutations frequency in cancer. ACyC, adenoid cystic carcinoma; CCLL, Cancer Cell Line Encyclopedia; ccRCC, clear cell renal cell carcinoma; GBM, glioblastoma multiforme; MM, Multiple myeloma; pRCC, papillary renal cell carcinoma; RCC, chromophobe renal cell carcinoma.

and studies were approved by local ethics committees. Exome and targeted sequencing analysis was performed as previously described.^{4,5} For further details, please see supplemental Methods, available on the *Blood* Web site.

Results and discussion

We first performed copy number variation analysis based on WES data from these 3 initial patients. This revealed the loss of chromosome 21 in all 3 patients (see supplemental data). Although these findings were validated by Affymetrix Cytoscan HD Arrays, copy number variation analysis of 7 additional patients failed to reveal chromosome 21 abnormalities (supplemental Table 1). Of note, 5 of 63 patients harbored deletions of 7q involving *BRAF* (7q34) and thereby causing loss of heterozygosity of the mutant *BRAF*-allele.

In addition to the *BRAF*-V600E mutation, WES identified 15 to 37 somatic mutations per patient (supplemental Table 2). Each case had several mutations previously associated with cancer or mutations in genes annotated in the COSMIC database (Figure 1A). All mutations called by WES were visually inspected in the Integrated Genome

Viewer. Among these mutations, we identified a frame shift deletion of *EZH2* (NM_004456: p.K406KfsX17), which has been described in myelodysplastic syndromes⁶ and a missense mutation of *ARID1A* (NM_139135: p.K1515Q), a gene that has been previously seen to be mutated in HCL.^{3,7}

We were able to compare the mutational landscape before and after BRAFi treatment in a patient with aggressive relapse after vemurafenib.⁸ Six new mutations emerged with BRAFi treatment (supplemental Figure 1). No mutations directly or indirectly activating RAS signaling and thus explaining resistance to BRAFi⁹ were identified. A stop gain in *KDM6A* (NM_021140: p.Q333*) was identified at relapse. *KDM6A* encodes a histone demethylase that specifically demethylates histone H3, which has been shown to be recurrently mutated in bladder cancer¹⁰ and rarely in acute myeloid leukemia or head and neck cancer. Mutations in *ARID1A*, *EZH2*, and *KDM6A* suggest that the epigenetic regulation machinery is recurrently targeted in HCL.

Surprisingly, 2 of 3 patients with refractory HCL that underwent WES were found to harbor somatic, inactivating mutations of *CDKN1B* (Figure 1A, NM_004064: p.W60* and p.A167QfsX37).^{8,11} To determine whether *CDKN1B* mutations are recurrent in HCL, we developed a custom, targeted next-generation sequencing panel and

sequenced *CDKN1B* (exon 1 and 2), *BRAF* (exon 15), and *MAP2K1* (exon 2 and 3)⁷ in 81 patients with HCL and 5 patients with HCL-variant, diagnosed according to standard criteria.¹ All 81 patients with classical HCL were found to harbor *BRAFV600E* mutations. One of 5 patients with HCLv had an *MAP2K1* (NM_002755: p.K57T) mutation. We identified 17 mutations of *CDKN1B*, including a 4-splice site, 11 nonsense and 2 missense mutations affecting 13 of 81 (16%) patients with classical HCL (Table 1; Figure 1B; supplemental Tables 2 and 3). All but one sample harbored at least one *CDKN1B* nonsense or splice site variant, except P0811, in which a missense mutation was identified. Three patients had more than one mutation implying selective pressure to inactivate CDKN1B. Comparison of allele frequencies of *BRAF* and *CDKN1B* mutations revealed that the majority (11 of 13, including treatment naïve patients) of *CDKN1B* mutations had allele frequencies very similar to those of the *BRAF* mutant clone (supplemental Figure 2). This suggests that *CDKN1B* mutations are early lesions and contribute to HCL pathogenesis. *CDKN1B* mutations were not present in 5 *BRAFV600E*-negative HCLv patients.

CDKN1B (p27) is a critical element of cell-cycle control and a known tumor suppressor.¹² CDKN1B binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controls cell-cycle progression in the G1 phase. Germ-line mutations of *CDKN1B* cause a multiple endocrine neoplasia 1-like (MEN1) phenotype. Menin, the product of MEN1, enhances expression of CDKN1B and CDKN2C, which suggests a functional link between MEN1 and CDKN1B.¹³ Recently, 8% (14 of 180) neuroendocrine tumors of the small intestine were reported to harbor frameshift *CDKN1B* deletions and 14% (7 of 50) neuroendocrine tumors of the small intestine had hemizygous deletions encompassing *CDKN1B*.¹⁴ *CDKN1B* mutations occur at very low frequencies in other malignancies (<5%) and have not been shown to cooccur with *BRAF V600E*. Comparison of the frequency of *CDKN1B* mutations across different cancers revealed that HCL has the highest incidence of *CDKN1B* mutations across cancer (Figure 1C). *CDKN1B* alterations have not been reported in other B-lymphoid malignancies, including the Cancer Genome Atlas (TCGA) study of diffuse large B-cell lymphoma.¹⁵ WES of 113 patients with CLL did not reveal variants for *CDKN1B* (unpublished data). In T-prolymphocytic leukemia, only one *CDKN1B* mutation was identified, but almost 50% harbored hemizygous deletions of *CDKN1B*.¹⁶

To test the clinical impact of *CDKN1B* mutations we correlated *CDKN1B* mutations with prior treatment and response. Although 2 of 5 of the initial refractory patients had CDKN1B mutations, 2 of 8 pre-treated patients and 10 of 43 treatment naïve patients had *CDKN1B* mutations (supplemental Tables 4 and 6; Fisher's exact test, $P = .73$). An influence of *CDKN1B* mutation on response to standard treatment was not detected (supplemental Tables 5 and 6; Fisher's exact test, $P = 1.00$).

Across *BRAF* mutant cancers, upregulation of cell cycle inhibitors such as *CDKN2A* (INK4/ARF), *CDKN1A*, and *CDKN1B* leads to senescence and constitutes a tumor suppressor mechanism. In fact, these genes are recurrently inactivated by genetic mechanisms during pathogenesis of many tumors. For example, *BRAF*-induced senescence in premalignant naevi¹⁷ is circumvented by deletion or mutation of *CDKN2A* in invasive melanoma (supplemental Figure 2A). Thus, the identification of recurrent inactivating mutations of *CDKN1B* in a *BRAFV600E*-mutant cancer suggests that *CDKN1B* loss may serve as a mechanism to impair cell cycle control and/or circumvent oncogene-induced senescence.

In addition to affecting cell cycle regulation and senescence, CDKN1B is a known opponent of cyclin D1, a gene regulated in a *BRAFV600E*-dependent manner and highly expressed in HCL.¹⁸ Thus, identification of inactivating mutations on CDKN1B in HCL further indicate pathway convergence.¹⁹

Table 1. *CDKN1B* (p27) mutations in *BRAFV600E*-mutant classical HCL

Sample identifier	<i>BRAF</i> mutation	<i>CDKN1B</i> mutation
P0001	c.1799T>A; (p.V600E)	c.500delC, (p.A167QfsX37)
P0013	c.1799T>A; (p.V600E)	c.180G>A, (p.W60*) c.238G>C; (p.E80Q)
P0621	c.1799T>A; (p.V600E)	c.333-353del21; p.G111del6
P0774	c.1799T>A; (p.V600E)	c.475+1G>A; (splice acceptor variant)
P0806	c.1799T>A; (p.V600E)	c.475+1G>T; (splice acceptor variant); c.596A>C; (p.X199S)
P0811	c.1799T>A (p.V600E)	c.281C>T; (p.P94L)*
P0911	c.1799T>A; (p.V600E)	c.5C>G; p.S2* c.58C>T; p.Q20* c.475+1G>T; (splice acceptor variant)
P0912	c.1799T>A; (p.V600E)	c.87C>A; (p.C29*)
P0914	c.1799T>A; (p.V600E)	c.179G>A; (p.W60*)
P0919	c.1799T>A; (p.V600E)	c.475+1G>T; (splice acceptor variant)
P0941	c.1799T>A; (p.V600E)	c.227G>A; (p.W76*)
P0897	c.1799T>A; (p.V600E)	c.596A>C; (p.X199S)
P0902	c.1799T>A; (p.V600E)	c.228G>A; (p.W76*)

In summary, we demonstrate that *CDKN1B* is inactivated in 16% of patients with classical HCL and is the second most common mutated gene in HCL. These results implicate cell cycle deregulation in the pathogenesis of HCL and suggest that *CDKN1B* serves as an important tumor suppressor in this disease.

Acknowledgments

The authors thank the DKFZ Genomics and Proteomics Core Facility, Ron Schweibinger (Heidelberg University), Ivo Buchhalter (Division of Applied Bioinformatics German Cancer Research Center [DKFZ]), Michael Heinold (Division of Applied Bioinformatics, DKFZ), and Rolf Kabbe (Division of Applied Bioinformatics, DKFZ) for their technical support and expertise; Heidelberg Center for Personalized Oncology (DKFZ-HIPO) for continued support and funding through the HIPO projects H005/H021; Roland Eils, Peter Lichter, Stephan Wolf, Katja Beck, and Janna Kirchhof for infrastructure and program development within DKFZ-HIPO and National Center for Tumor Diseases Precision Oncology Program; and Tina Uhrig for technical assistance.

This work was supported by grants from the Heidelberg Research Centre for Molecular Medicine (S.D.), the Arbib Foundation (M.E.), the Hairy Cell Leukemia Foundation, and the "Deutsche Krebshilfe."

Authorship

Contribution: S.D., T.Z., and O.A.W. designed research; S.D., J.H., S.C.-W.L., X.L., M.S., M.G., A.J., M.M., T.H., D.R., O.A.W., and T.Z. performed research; D.G., S.J., M.J.S.D., X.T., J.D., C.D., T.H., A.D.H., and E.M. provided clinical samples; S.D., B.H., M.O., B.W., M.A., W.H., B.B., and T.Z. analyzed data; X.T. and C.D. provided follow-up of clinical samples; M.E. provided follow-up of patients; and S.D., C.v.K., S.F., W.H., H.G., O.A.W., and T.Z. wrote the paper.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Thorsten Zenz, Department of Medicine V, University Hospital Heidelberg, Heidelberg, Germany, Im Neuenheimer Feld 410, 69120 Heidelberg, Germany; or Department of Translational

Oncology, National Center for Tumor Diseases (NCT), German Cancer Research Center (DKFZ); e-mail: thorsten.zenz@nct-heidelberg.de.

References

- Grever MR. How I treat hairy cell leukemia. *Blood*. 2010;115(1):21-28.
- Tiacci E, Trifonov V, Schiavoni G, et al. BRAF mutations in hairy-cell leukemia. *N Engl J Med*. 2011;364(24):2305-2315.
- Chung SS, Kim E, Park JH, et al. Hematopoietic stem cell origin of BRAFV600E mutations in hairy cell leukemia. *Sci Transl Med*. 2014;6(238):238ra271.
- Jethwa A, Hüllein J, Stolz T, et al. Targeted resequencing for analysis of clonal composition of recurrent gene mutations in chronic lymphocytic leukaemia. *Br J Haematol*. 2013;163(4):496-500.
- Yaktapour N, Meiss F, Mastroianni J, et al. BRAF inhibitor-associated ERK activation drives development of chronic lymphocytic leukemia. *J Clin Invest*. 2014;124(11):5074-5084.
- Nikoloski G, Langemeijer SM, Kuiper RP, et al. Somatic mutations of the histone methyltransferase gene EZH2 in myelodysplastic syndromes. *Nat Genet*. 2010;42(8):665-667.
- Waterfall JJ, Arons E, Walker RL, et al. High prevalence of MAP2K1 mutations in variant and IGHV4-34-expressing hairy-cell leukemias. *Nat Genet*. 2014;46(1):8-10.
- Samuel J, Macip S, Dyer MJ. Efficacy of vemurafenib in hairy-cell leukemia. *N Engl J Med*. 2014;370(3):286-288.
- Nazarian R, Shi H, Wang Q, et al. Melanomas acquire resistance to B-RAF(V600E) inhibition by RTK or N-RAS upregulation. *Nature*. 2010;468(7326):973-977.
- Cancer Genome Atlas Research Network. Comprehensive molecular characterization of urothelial bladder carcinoma. *Nature*. 2014;507(7492):315-322.
- Dietrich S, Glimm H, Andrulis M, von Kalle C, Ho AD, Zenz T. BRAF inhibition in refractory hairy-cell leukemia. *N Engl J Med*. 2012;366(21):2038-2040.
- Chu IM, Hengst L, Slingerland JM. The Cdk inhibitor p27 in human cancer: prognostic potential and relevance to anticancer therapy. *Nat Rev Cancer*. 2008;8(4):253-267.
- Milne TA, Hughes CM, Lloyd R, et al. Menin and MLL cooperatively regulate expression of cyclin-dependent kinase inhibitors. *Proc Natl Acad Sci USA*. 2005;102(3):749-754.
- Francis JM, Kiezun A, Ramos AH, et al. Somatic mutation of CDKN1B in small intestine neuroendocrine tumors. *Nat Genet*. 2013;45(12):1483-1486.
- Cerami E, Gao J, Dogrusoz U, et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012;2(5):401-404.
- Le Toriell E, Despouy G, Pierron G, et al. Haploinsufficiency of CDKN1B contributes to leukemogenesis in T-cell prolymphocytic leukemia. *Blood*. 2008;111(4):2321-2328.
- Michaloglou C, Vredevelde LC, Soengas MS, et al. BRAFE600-associated senescence-like cell cycle arrest of human naevi. *Nature*. 2005;436(7051):720-724.
- Dietrich S, Hüllein J, Hundemer M, et al. Continued response off treatment after BRAF inhibition in refractory hairy cell leukemia. *J Clin Oncol*. 2013;31(19):e300-e303.
- Motti ML, De Marco C, Califano D, et al. Loss of p27 expression through RAS→BRAF→MAP kinase-dependent pathway in human thyroid carcinomas. *Cell Cycle*. 2007;6(22):2817-2825.



blood

2015 126: 1005-1008

doi:10.1182/blood-2015-04-643361 originally published
online June 11, 2015

Recurrent *CDKN1B* (p27) mutations in hairy cell leukemia

Sascha Dietrich, Jennifer Hülle, Stanley Chun-Wei Lee, Barbara Hutter, David Gonzalez, Sandrine Jayne, Martin J. S. Dyer, Malgorzata Oles, Monica Else, Xiyang Liu, Mikolaj Slabicki, Bian Wu, Xavier Troussard, Jan Dürig, Mindaugas Andrusis, Claire Dearden, Christof von Kalle, Martin Granzow, Anna Jauch, Stefan Fröhling, Wolfgang Huber, Manja Meggendorfer, Torsten Haferlach, Anthony D. Ho, Daniela Richter, Benedikt Brors, Hanno Glimm, Estella Matutes, Omar Abdel Wahab and Thorsten Zenz

Updated information and services can be found at:

<http://www.bloodjournal.org/content/126/8/1005.full.html>

Articles on similar topics can be found in the following Blood collections

[Brief Reports](#) (1840 articles)

[Lymphoid Neoplasia](#) (2097 articles)

Information about reproducing this article in parts or in its entirety may be found online at:

http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:

<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:

<http://www.bloodjournal.org/site/subscriptions/index.xhtml>