

ITYM 2021 - Problem 9: Wobbly Tables.

Team France

Composed by :

De Ridder Achille, Harter Louis-Max,
Fourcin Emile, Quille Maxime
Varnet Philémon, Leroux Hubert

Supervised by :

Lenoir Théo et Béreau Antoine

June 2021

Abstract

This document is dealing with the ninth problem of the ITYM 2021.
The problem consist of placing a table on the floor of a room. The floor is parametrised by a surface $z = f(x, y)$, where $f : [-1, 1]^2 \rightarrow \mathbb{R}$ is a continuous map.
For now, the document is dealing just with the first question. But a lot of elements of the first question help to do the next questions.

Contents

Introduction	3
Question 1	3
0.1 Square	3
0.1.1 The horizontal plane	3
0.1.2 The inclined plane	3
0.1.3 Horse Saddle	4
0.1.4 Sphere	4
0.1.5 Double periodic surface	6
0.2 Rectangle	6
0.2.1 The horizontal plane	6
0.2.2 The inclined plane	6
0.2.3 The horse saddle	8
0.2.4 The sphere	10
0.2.5 The double periodic surface	12
0.3 Rhombus	12
0.3.1 Horizontal plane	12
0.3.2 Inclined Plane	13
0.3.3 Horse Saddle	18
0.3.4 Sphere	19
0.4 Any convex quadrilateral	20
Question 2	20
0.5 Stabilised	22
0.6 Wobbly	22
Question 3	22
0.7 Inclined Plane	22

Introduction

Question 1

In this question, we will try to ground some tables on the different possible floors.

0.1 Square

The table is a square with sides of length $\sqrt{2}$.

0.1.1 The horizontal plane

The floor is the function

$$f_1 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto 0 \end{cases}$$

We can put each legs:

- $P_1 = (-1, 0, 0)$
- $P_2 = (1, 0, 0)$
- $P_3 = (0, -1, 0)$
- $P_4 = (0, 1, 0)$

$P_1P_2P_3P_4$ is a square with sides of length $\sqrt{2}$ and the table is *grounded*:

$$\forall i \in [1, 4], z_i = f(x_i, y_i)$$

0.1.2 The inclined plane

We will now generalise with an inclined plane defined as follow:

$$f_2 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto ax + by, \text{ with } a, b \in \mathbb{R} \end{cases}$$

The plane Π of the square $P_1P_2P_3P_4$ should be $ax + by - z = 0$

Firstly we put P_1 in the corner. The coordinates of P_1 are now : $(1, 1, a + b)$

Now, we will take care of P_2 We want:

- (1) $P_1P_2 = \sqrt{2}$
- (2) $y_2 = y_1 = 1$
- (3) $z_2 = f(x_2, y_2) = ax_2 + by_2$

From the first equality, we obtain:

$$\sqrt{(1 - x_2)^2 + (1 - y_2)^2 + (a + b - ax_2 - by_2)^2} = \sqrt{2}$$

so, $(1 - x_2)^2 + (1 - y_2)^2 + (a + b - ax_2 - by_2)^2 = 2$

but, with $y_2 = 1$, we can reduce:

$$(1 - x_2)^2 + (a - ax_2)^2 = 2$$

and we have: $(1 - x_2)^2 \times (a^2 + 1) = 2$

$$1 - x_2 = \frac{\sqrt{2}}{\sqrt{a^2 + 1}} \text{ car } a^2 + 1 \geq 0$$


we finally have $x_2 = 1 - \frac{\sqrt{2}}{\sqrt{a^2 + 1}}$

With the same process for P_4 , where:

- (1) $P_1P_4 = \sqrt{2}$
- (2) $x_4 = x_1 = 1$
- (3) $z_4 = f(x_4, y_4) = ax_4 + by_4$

We obtain: $P_4 = (1, 1 - \frac{\sqrt{2}}{\sqrt{b^2 + 1}}, a + b \times (1 - \frac{\sqrt{2}}{\sqrt{b^2 + 1}}))$

And the final point is $P_3 = (1 - \frac{\sqrt{2}}{\sqrt{a^2 + 1}}, 1 - \frac{\sqrt{2}}{\sqrt{b^2 + 1}}, a \times (1 - \frac{\sqrt{2}}{\sqrt{a^2 + 1}}) + b \times (1 - \frac{\sqrt{2}}{\sqrt{b^2 + 1}}))$



1: Square on an inclined plane

0.1.3 Horse Saddle

$$f_3 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto sxy, \text{ with } s \in \mathbb{R}^* \end{cases}$$

We can put the legs like this:

1. $P_1 = (0, 1, 0)$
2. $P_2 = (0, -1, 0)$
3. $P_3 = (-1, 0, 0)$
4. $P_4 = (1, 0, 0)$

And we have a *grounded* table because:


$$\forall i \in [1, 4], z_i = f(x_i, y_i)$$

0.1.4 Sphere


$$f_4 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto \sqrt{1 - \frac{x^2 + y^2}{R^2}}, \text{ with } R > \sqrt{2} \end{cases}$$

1. $P_1 = (0, 1, z)$
2. $P_2 = (0, -1, z)$
3. $P_3 = (-1, 0, z)$
4. $P_4 = (1, 0, z)$


With $z = \sqrt{1 - \frac{1}{R^2}} = \sqrt{\frac{(R+1)(R-1)}{R^2}}$, the table is grounded.



2: Square on a saddle



3: Square on a sphere



4: Square on a double periodic surface

0.1.5 Double periodic surface

$$f_5 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto \cos\left(\frac{2\pi x}{\omega}\right) \cos\left(\frac{2\pi y}{\omega}\right), \text{ with } \omega > \sqrt{2} \end{cases}$$

1. $P_1 = (0, 1, z)$
2. $P_2 = (0, -1, z)$
3. $P_3 = (-1, 0, z)$
4. $P_4 = (1, 0, z)$

We can see that $\forall X \in \mathbb{R}, \cos(-X) = \cos(X)$ and $\cos(0) = 1$. With these properties, $z = \cos\left(\frac{2\pi}{\omega}\right)$, the table is grounded.

0.2 Rectangle

Now we have a **rectangle** with sides r and s such that $r^2 + s^2 = 4$. This implicate that $r \leq 2$ and $s \leq 2$ which is logical because the surface is defined as a square area of dimensions 2×2 .

0.2.1 The horizontal plane

This surface is a sub-case of the next surface with $a = 0$ and $b = 0$.

0.2.2 The inclined plane

$$f_2 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto ax + by, \text{ with } a, b \in \mathbb{R} \end{cases}$$

First we set $P_1 = (1, 1, f_2(1, 1)) = (1, 1, a + b)$

We want to place P_2 according to these hypotheses:

1. $P_1P_2 = r$
2. $y_1 = y_2 = 1$
3. $z_2 = f(x_2, y_2) = ax_2 + by_2$

From the first equality, we deduce:

$$\sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2} = r$$

Using (1) and (2), we have, and by squaring each sides:

$$(x_1 - x_2)^2 + (a + b - ax_2 - b)^2 = r^2$$

i.e.

$$(1 - x_2)^2 + a^2(1 - x_2)^2 = r^2$$

so by factorising by $(1 - x_2)^2$, we obtain:

$$(1 - x_2)^2(1 + a^2) = r^2$$

We can divide each sides by $(1 + a^2) > 0$:

$$(1 - x_2)^2 = \frac{r^2}{1 + a^2}$$

$$x_2 = 1 - \sqrt{\frac{r^2}{1 + a^2}}$$

In conclusion, we have the point $P_2 = (1 - \sqrt{\frac{r^2}{1 + a^2}}, 1, a(1 - \sqrt{\frac{r^2}{1 + a^2}}) + b)$

We can apply the same process for P_3 with these hypotheses:

1. $P_1P_4 = s$
2. $x_1 = x_4 = 1$
3. $z_4 = f(x_4, y_4) = ax_4 + by_4$

From the first equality, we deduce:

$$\sqrt{(x_1 - x_4)^2 + (y_1 - y_4)^2 + (z_1 - z_4)^2} = s$$

Using (1) and (2), we have, and by squaring each sides:

$$(y_1 - y_4)^2 + (a + b - a - b \times y_4)^2 = s^2$$

i.e.

$$(1 - y_4)^2 + b^2(1 - y_4)^2 = s^2$$

so by factorising by $(1 - y_4)^2$, we obtain:

$$(1 - y_4)^2(1 + b^2) = s^2$$

We can divide each sides by $(1 + b^2) > 0$:

$$(1 - y_4)^2 = \frac{s^2}{1 + b^2}$$

$$y_4 = 1 - \sqrt{\frac{s^2}{1+b^2}}$$

In conclusion, we have the point $P_4 = (1, 1 - \sqrt{\frac{s^2}{1+b^2}}, a + b \times (1 - \sqrt{\frac{s^2}{1+b^2}})$

For P_3 , the coordinates are: $P_3 = (1 - \sqrt{\frac{r^2}{1+a^2}}, 1 - \sqrt{\frac{s^2}{1+b^2}}, a(1 - \sqrt{\frac{r^2}{1+a^2}}) + b \times (1 - \sqrt{\frac{s^2}{1+b^2}})$

We finally have our square. We can note $R = 1 - \sqrt{\frac{r^2}{1+a^2}}$ and $S = 1 - \sqrt{\frac{s^2}{1+b^2}}$ To sum up, the coordiantes of the points are:

- $P_1(1, 1, a + b)$
- $P_2(R, 1, aR + b)$
- $P_3(R, S, aR + bS)$
- $P_4(1, S, a + bS)$

0.2.3 The horse saddle

We search $(a, b) \in \mathbb{R}_+^2$ such that:

- $P_1 = (a, b, f_3(a, b))$
- $P_2 = (b, a, f_3(b, a))$
- $P_3 = (-a, -b, f_3(-a, -b))$
- $P_4 = (-b, -a, f_3(-b, -a))$

Because in this situation, we have:

$$f_3(a, b) = f_3(b, a) = f_3(-b, -a) = f_3(-a, -b) = sab$$

So the table is grounded and we have a rectangle.

Now we can search a and b depending on r and s

We know that:

1. $P_1P_2 = r$
2. $P_1P_4 = s$

This is equivalent to:

$$\begin{aligned} & \begin{cases} (x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2 = r^2 \\ (x_4 - x_1)^2 + (y_4 - y_1)^2 + (z_4 - z_1)^2 = s^2 \end{cases} \\ & \Leftrightarrow \begin{cases} (b - a)^2 + (a - b)^2 = r^2 \\ (-b - a)^2 + (-a - b)^2 = s^2 \end{cases} \\ & \Leftrightarrow \begin{cases} 2(a - b)^2 = r^2 \\ 2(a + b)^2 = s^2 \end{cases} \\ & \Leftrightarrow \begin{cases} 2(a^2 - 2ab + b^2) = r^2 \quad (L_1) \\ 2(a^2 + 2ab + b^2) = s^2 \quad (L_2) \end{cases} \end{aligned}$$

$$L_1 \leftarrow L_1 + L_2$$

$$L_2 \leftarrow L_2 - L_1$$

$$\Leftrightarrow \begin{cases} 4(a^2 + b^2) = r^2 + s^2 \\ 8ab = s^2 - r^2 \end{cases}$$

If $r^2 = s^2$, we return to the case of the square. Here, $r \neq s$, so $s^2 - r^2 \neq 0$. Then, $a \neq 0$ and $b \neq 0$ due to the second equality.

So $a = \frac{s^2 - r^2}{8b}$ (we have $b \neq 0$).

We can substitute a in the first equality, we obtain:

$$(E) : 4 \times \left[\left(\frac{s^2 - r^2}{8b} \right)^2 + b^2 \right] = r^2 + s^2$$

$$b \text{ is solution of (E)} \Leftrightarrow \frac{(s^2 - r^2)^2}{16 \times b^2} + 4 \times b^2 - (r^2 + s^2) = 0$$

$$\Leftrightarrow \frac{(s^2 - r^2)^2}{16 \times b^2} + 4 \times b^2 - (r^2 + s^2) = 0$$

$$\Leftrightarrow 4 \times b^4 - (r^2 + s^2) \times b^2 + \frac{(s^2 - r^2)^2}{16} = 0 \quad (b \neq 0)$$

Let's set $B = b^2$

$$b \text{ is solution of (E)} \Leftrightarrow B \text{ is solution of (E')}$$

With: $(E') : 4 \times B^2 - (r^2 + s^2) \times B + \frac{(s^2 - r^2)^2}{16} = 0$ Let Δ be the discriminant of this equation. We have:

$$\Delta = (r^2 + s^2)^2 - 4 \times 4 \times \frac{(s^2 - r^2)^2}{16}$$

$$\Delta = (r^2 + s^2)^2 - (s^2 - r^2)^2$$

$$\Delta = (r^2 + s^2 + s^2 - r^2) \times (r^2 + s^2 - s^2 + r^2)$$

$$\Delta = (2 \times s^2) \times (2 \times r^2)$$

$$\Delta = (2rs)^2$$

We have $\Delta \geq 0$ and $\sqrt{\Delta} = 2rs$

The solutions are:

$$B = \frac{r^2 + s^2 \pm (2rs)}{8}$$

$$\Leftrightarrow b^2 = \frac{(r \pm s)^2}{8}$$

$$\Leftrightarrow b = \frac{r \pm s}{2\sqrt{2}} \quad (b > 0)$$


Then, $a = \frac{1}{8b} \times (s^2 - r^2)$

$$a = \frac{2\sqrt{2}}{8} \times \frac{s^2 - r^2}{r \pm s}$$

$$a = \frac{1}{2\sqrt{2}} \times \frac{(s^2 - r^2) \times (r \mp s)}{(r \pm s) \times (r \mp s)}$$

$$a = \frac{s \pm r}{2\sqrt{2}}$$

In conclusion, we have: $(a, b) = \left(\frac{s+r}{2\sqrt{2}}, \frac{s-r}{2\sqrt{2}} \right)$ ou $(a, b) = \left(\frac{s-r}{2\sqrt{2}}, \frac{s+r}{2\sqrt{2}} \right)$



5: Rectangle on a saddle

0.2.4 The sphere

We have this surface:

$$f_4 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto \sqrt{1 - \frac{x^2 + y^2}{R^2}}, \text{ with } R > \sqrt{2} \end{cases}$$

We name Ω the center of the rectangle. Let $x_\Omega = y_\Omega = 0$. By applying Pythagoras theorem in $P_1 P_2 P_4$, we obtain:

$$P_1 P_2^2 + P_1 P_4^2 = P_2 P_4^2$$

$$\text{i.e. } s^2 + r^2 = (2P_2 \Omega)^2$$

$$\text{i.e. } 4 = 4P_2 \Omega^2$$

so, $P_2 \Omega = 1$

The rectangle is in the circle $C : x^2 + y^2 = 1$ and $z = z_\Omega$


Now we want to calculate z for each leg of the table.

We want the leg on the surface defined as :


$$f_4(x, y) = \sqrt{1 - \frac{x^2 + y^2}{R^2}}, \text{ with } R > \sqrt{2}$$

But we have $x^2 + y^2 = 1$

$$\text{Then } z_1 = z_2 = z_3 = z_4 = \sqrt{1 - \frac{1}{R^2}} = \frac{\sqrt{(R+1) \times (R-1)}}{R}$$



6: Rectangle



7: Rectangle on a Sphere

And we have our table that is *stabilised*.

0.2.5 The double periodic surface

$$f_5 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto \cos(\frac{2\pi x}{\omega}) \cos(\frac{2\pi y}{\omega}), \text{ with } \omega > \sqrt{2} \end{cases}$$

Using the fact that *cosinus* is an **even** function, we can take the points:

1. $P_1 = (-\frac{r}{2}, -\frac{s}{2}, z_1)$
2. $P_2 = (\frac{r}{2}, -\frac{s}{2}, z_2)$
3. $P_3 = (\frac{r}{2}, \frac{s}{2}, z_3)$
4. $P_4 = (-\frac{r}{2}, \frac{s}{2}, z_4)$

And we have $z_1 = z_2 = z_3 = z_4 = \cos(\frac{r\pi}{\omega}) \times \cos(\frac{s\pi}{\omega})$ So the table is stabilised.

0.3 Rhombus


We have a rhombus with an axis of length 2 and an axis of length l , where $0 < l < 2$. Let Ω be the center of the rhombus.

0.3.1 Horizontal plane

$$f_1 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto 0 \end{cases}$$

Then we place the points as follow:

1. $P_1 = (-1, 0, 0)$



8: Rectangle on a double periodic surface

2. $P_2 = (0, -\frac{l}{2}, 0)$
3. $P_1 = (1, 0, 0)$
4. $P_4 = (0, +\frac{l}{2}, 0)$

And we have a stabilised rhombus on the horizontal plane.

0.3.2 Inclined Plane

$$f_2 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto ax + by, \text{ with } a, b \in \mathbb{R}^2 \end{cases}$$

Let's name Ω the center of our rhombus.

We will place Ω on the (Oz) axis. Then $x_\Omega = y_\Omega = 0$.


Ω is on the floor, so $z_\Omega = f_2(x_\Omega, y_\Omega) = a \times 0 + b \times 0 = 0$

$\Omega = (0, 0, 0)$


We will search the coordinates of $P_1 = (x_1, y_1, z_1)$ and $P_2 = (x_2, y_2, z_2)$ that satisfies these propositions:

- $z_1 = f_2(x_1, y_1)$
- $z_2 = f_2(x_2, y_2)$
- $\Omega P_1 = 1$ because Ω is the middle of $[P_1 P_3]$ and $P_1 P_3 = 2$
- $\Omega P_2 = \frac{l}{2}$ because Ω is the middle of $[P_2 P_4]$ and $P_2 P_4 = l$
- $\overrightarrow{\Omega P_1} \cdot \overrightarrow{\Omega P_2} = \vec{0}$

We obtain the system (S) :



9: Rhombus on an horizontal plane



10: Rhombus

$$(S) : \begin{cases} z_1 = a \times x_1 + b \times y_1 \\ z_2 = a \times x_2 + b \times y_2 \\ (x_1 - x_\Omega)^2 + (y_1 - y_\Omega)^2 + (z_1 - z_\Omega)^2 = 1 \\ (x_2 - x_\Omega)^2 + (y_2 - y_\Omega)^2 + (z_2 - z_\Omega)^2 = \frac{l^2}{4} \\ (x_1 - x_\Omega) \times (x_2 - x_\Omega) + (y_1 - y_\Omega) \times (y_2 - y_\Omega) + (z_1 - z_\Omega) \times (z_2 - z_\Omega) = 0 \end{cases}$$

But $x_\Omega = y_\Omega = z_\Omega = 0$

$$(S) \Leftrightarrow \begin{cases} z_1 = a \times x_1 + b \times y_1 \\ z_2 = a \times x_2 + b \times y_2 \\ x_1^2 + y_1^2 + (ax_1 + by_1)^2 = 1 \\ x_2^2 + y_2^2 + (ax_2 + by_2)^2 = \frac{l^2}{4} \\ x_1 \times x_2 + y_1 \times y_2 + (ax_1 + by_1) \times (ax_2 + by_2) = 0 \end{cases}$$

$$\Leftrightarrow \begin{cases} z_1 = a \times x_1 + b \times y_1 \\ z_2 = a \times x_2 + b \times y_2 \\ x_1^2 \times (a^2 + 1) + y_1^2 \times (b^2 + 1) + 2abx_1y_1 = 1 \\ x_2^2 \times (a^2 + 1) + y_2^2 \times (b^2 + 1) + 2abx_2y_2 = \frac{l^2}{4} \\ x_1x_2 \times (a^2 + 1) + y_1y_2 \times (b^2 + 1) + ab \times (x_1y_2 + x_2y_1) = 0 \end{cases}$$

Now we will take arbitrarily $y_1 = 0$:

$$(S) \Leftrightarrow \begin{cases} z_1 = a \times x_1 & (1) \\ z_2 = a \times x_2 + b \times y_2 & (2) \\ x_1^2 \times (a^2 + 1) = 1 & (3) \\ x_2^2 \times (a^2 + 1) + y_2^2 \times (b^2 + 1) + 2abx_2y_2 = \frac{l^2}{4} & (4) \\ x_1x_2 \times (a^2 + 1) + ab \times (x_1y_2) = 0 & (5) \end{cases}$$

From (3), we can deduce that: $x_1 = \pm \sqrt{\frac{1}{a^2+1}}$. We take x_1 negativ so $x_1 = -\sqrt{\frac{1}{a^2+1}} < 0$. We can see that $x_1 \neq 0$

From (5), we deduce: $abx_1y_2 = -x_1x_2(a^2 + 1)$, so $y_2 = \frac{-x_2 \times (a^2+1)}{ab}$ (we suppose that $a \neq 0$ and $b \neq 0$ and we know that $x_1 \neq 0$)

x_2 remains to be found.

We will use (4): $x_2^2 \times (a^2 + 1) + y_2^2 \times (b^2 + 1) + 2abx_2y_2 = \frac{l^2}{4}$

By replacing the value of y_2 by $\frac{-x_2 \times (a^2+1)}{ab}$, we obtain:

$$\begin{aligned} & x_2^2 \times (a^2 + 1) + x_2^2 \times \frac{(a^2 + 1)^2}{(ab)^2} \times (b^2 + 1) - 2abx_2^2 \times \frac{a^2 + 1}{ab} = \frac{l^2}{4} \\ \Leftrightarrow & x_2^2 \times (a^2 + 1) + x_2^2 \times \frac{(a^2 + 1)^2}{(ab)^2} \times (b^2 + 1) - 2x_2^2 \times (a^2 + 1) = \frac{l^2}{4} \\ \Leftrightarrow & x_2^2 \times (a^2 + 1) \times \left[1 + \frac{(a^2 + 1)}{(ab)^2} \times (b^2 + 1) - 2 \right] = \frac{l^2}{4} \\ \Leftrightarrow & x_2^2 \times (a^2 + 1) \times \left[\frac{a^2 \times b^2 + a^2 + b^2 + 1 - (ab)^2}{(ab)^2} \right] = \frac{l^2}{4} \\ \Leftrightarrow & x_2^2 \times (a^2 + 1) \times \left[\frac{a^2 + b^2 + 1}{(ab)^2} \right] = \frac{l^2}{4} \\ \Leftrightarrow & x_2^2 = \frac{l^2 \times (ab)^2}{4 \times (a^2 + 1) \times (a^2 + b^2 + 1)} \end{aligned}$$

$$\Leftrightarrow x_2 = \pm \frac{l \times ab}{2 \times \sqrt{(a^2+1) \times (a^2+b^2+1)}}$$

We will take $x_2 < 0$.

$$\text{So } x_2 = -\frac{l \times ab}{2 \times \sqrt{(a^2+1) \times (a^2+b^2+1)}}$$

Then:

$$y_2 = +\frac{l \times ab}{2 \times \sqrt{(a^2+1) \times (a^2+b^2+1)}} \times \frac{(a^2+1)}{ab} = \frac{l \times \sqrt{(a^2+1)}}{2 \times \sqrt{(a^2+b^2+1)}}$$

and:

$$z_2 = ax_2 + by_2 \tag{1}$$

$$= \frac{-l \times (a^2b)}{2 \times \sqrt{(a^2+1) \times (a^2+b^2+1)}} + \frac{l \times b \times \sqrt{(a^2+1)}}{2 \times \sqrt{(a^2+b^2+1)}} \tag{2}$$

$$= \frac{l \times b}{2 \times \sqrt{(a^2+b^2+1)}} \times \left(\sqrt{a^2+1} - \frac{a^2}{\sqrt{a^2+1}} \right) \tag{3}$$

$$= \frac{l \times b}{2 \times \sqrt{(a^2+b^2+1)}} \times \left(\frac{a^2+1-a^2}{\sqrt{a^2+1}} \right) \tag{4}$$

$$= \frac{l \times b}{2 \times \sqrt{(a^2+b^2+1)}} \times \left(\frac{1}{\sqrt{a^2+1}} \right) \tag{5}$$

$$= \frac{l \times b}{2 \times \sqrt{(a^2+b^2+1)(a^2+1)}} \tag{6}$$

Now we have the coordinates of P_1 and P_2 :

1. $P_1 = \left(-\sqrt{\frac{1}{a+1}}, 0, -a \times \sqrt{\frac{1}{a+1}} \right)$
2. $P_2 = \left(-\frac{l \times ab}{2 \times \sqrt{(a^2+1) \times (a^2+b^2+1)}}, \frac{l \times \sqrt{(a^2+1)}}{2 \times \sqrt{(a^2+b^2+1)}}, \frac{l \times b}{2 \times \sqrt{(a^2+b^2+1)(a^2+1)}} \right)$

By symmetrie, we deduce:

$$\begin{cases} (x_3, y_3, z_3) = (-x_1, -y_1, -z_1) \\ (x_4, y_4, z_4) = (-x_2, -y_2, -z_2) \end{cases}$$

so:

1. $P_3 = \left(\sqrt{\frac{1}{a+1}}, 0, a \times \sqrt{\frac{1}{a+1}} \right)$
2. $P_4 = \left(+\frac{l \times ab}{2 \times \sqrt{(a^2+1) \times (a^2+b^2+1)}}, -\frac{l \times \sqrt{(a^2+1)}}{2 \times \sqrt{(a^2+b^2+1)}}, -\frac{l \times b}{2 \times \sqrt{(a^2+b^2+1)(a^2+1)}} \right)$


And we have our stabilised rhombus.

In our demonstration we supposed that neither a and b were equalled to 0. If $a = b = 0$, we have a horizontal plane, which had already been done in the previous section.

Let see what happens if $b = 0$ (the case where $a = 0$ is symmetrical).

We have

$$f'_2 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto bx, \text{ with } a \in \mathbb{R}^* \end{cases}$$



11: Rhombus on an inclined plane

We have $\Omega = (0, 0, 0)$. The system is now:

$$(S) : \begin{cases} z_1 = a \times x_1 \\ z_2 = a \times x_2 \\ (x_1 - x_\Omega)^2 + (y_1 - y_\Omega)^2 + (z_1 - z_\Omega)^2 = 1 \\ (x_2 - x_\Omega)^2 + (y_2 - y_\Omega)^2 + (z_2 - z_\Omega)^2 = \frac{l^2}{4} \\ (x_1 - x_\Omega) \times (x_2 - x_\Omega) + (y_1 - y_\Omega) \times (y_2 - y_\Omega) + (z_1 - z_\Omega) \times (z_2 - z_\Omega) = 0 \end{cases}$$


$$\Leftrightarrow \begin{cases} z_1 = a \times x_1 \\ z_2 = a \times x_2 \\ x_1^2 + y_1^2 + (ax_1)^2 = 1 \\ x_2^2 + y_2^2 + (ax_2)^2 = \frac{l^2}{4} \\ x_1 \times x_2 + y_1 \times y_2 + z_1 \times z_2 = 0 \end{cases}$$

We take $y_1 = 0$

$$(S) \Leftrightarrow \begin{cases} z_1 = a \times x_1 \\ z_2 = a \times x_2 \\ x_1^2 \times (1 + a^2) = 1 \\ x_2^2 + y_2^2 + (ax_2)^2 = \frac{l^2}{4} \\ x_1 \times x_2 + ax_1 \times ax_2 = 0 \end{cases}$$

$$\Leftrightarrow \begin{cases} z_1 = a \times x_1 \\ z_2 = a \times x_2 \\ x_1^2 = \frac{1}{1+a^2} \quad a^2 + 1 > 0 \\ x_2^2 + y_2^2 + (ax_2)^2 = \frac{l^2}{4} \\ x_1 \times x_2 \times (a^2 + 1) = 0 \end{cases}$$

$$\Leftrightarrow \begin{cases} z_1 = a \times x_1 \\ z_2 = a \times x_2 \\ x_1 s = \pm \sqrt{\frac{1}{1+a^2}} \quad a^2 + 1 > 0 \\ x_2^2 + y_2^2 + (ax_2)^2 = \frac{l^2}{4} \\ x_1 = 0 \text{ or } x_2 = 0 \end{cases}$$



12: Rhombus on an inclined plane bis

$$\Leftrightarrow \begin{cases} z_1 = a \times x_1 \\ z_2 = a \times x_2 \\ x_1 = \pm \sqrt{\frac{1}{1+a^2}} \quad \text{so } x_1 \neq 0 \\ x_2 = 0 \\ y_2^2 = \frac{l^2}{4} \end{cases}$$

$$\Leftrightarrow \begin{cases} z_1 = a \times x_1 \\ z_2 = a \times x_2 \\ x_1 = \pm \sqrt{\frac{1}{1+a^2}} \\ x_2 = 0 \\ y_2 = \pm \frac{l}{2} \end{cases}$$

We will take $x_1 = -\sqrt{\frac{1}{1+a^2}}$ and $y_2 = -\frac{l}{2}$

We have our points:

1. $P_1 = (-\sqrt{\frac{1}{1+a^2}}, 0, -a \times \sqrt{\frac{1}{1+a^2}})$
2. $P_2 = (0, -\frac{l}{2}, 0)$
3. $P_3 = (\sqrt{\frac{1}{1+a^2}}, 0, a \times \sqrt{\frac{1}{1+a^2}})$
4. $P_4 = (0, +\frac{l}{2}, 0)$

0.3.3 Horse Saddle


For the horse saddle it will much more simple.

$$f_3 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto sxy, \text{ with } s \in \mathbb{R}^* \end{cases}$$

If we place each leg on the (Ox) axis or the (Oy) axis, $f(x, y) = sxy = 0$

Then we place the points as follow:

1. $P_1 = (-1, 0, 0)$
2. $P_2 = (0, -\frac{l}{2}, 0)$



13: Rhombus on a saddle

3. $P_1 = (1, 0, 0)$

4. $P_4 = (0, +\frac{l}{2}, 0)$

And we have a stabilised rhombus on the saddle.

0.3.4 Sphere

$$f_4 : \begin{cases} [-1, 1] \times [-1, 1] \rightarrow \mathbb{R} \\ (x, y) \mapsto \sqrt{1 - \frac{x^2 + y^2}{R^2}}, \text{ with } R > \sqrt{2} \end{cases}$$

Now we will prove that it is impossible to have a stabilised table.

Let Ω be the center of our rhombus. Without losing generality, we take $\Omega \in (Oz)$. Thus $x_\Omega = y_\Omega = 0$. We place $P_2 = (-\frac{l}{2}, 0, f_4(-\frac{l}{2}, 0))$ and $P_4 = (\frac{l}{2}, 0, f_4(\frac{l}{2}, 0))$.

$$f_4(\frac{l}{2}, 0) = \sqrt{1 - \frac{l^2}{4R^2}} = \sqrt{1 - \frac{l^2}{4R^2}} = \frac{\sqrt{4R^2 - l^2}}{2R} = \frac{\sqrt{(2R-l)(2R+l)}}{2R}$$

$$\text{We have necessarily } z_\Omega = z_2 = z_4 = \frac{\sqrt{(2R-l)(2R+l)}}{2R}.$$

Now we will seek for P_1

We have the following hypothesis:

$$(S) \begin{cases} \overrightarrow{\Omega P_1} \cdot \overrightarrow{\Omega P_4} = \overrightarrow{0} & (1) \\ \Omega P_1 = 1 & (2) \\ z_1 = f_4(x_1, y_1) = \sqrt{1 - \frac{x_1^2 + y_1^2}{R^2}} & (3) \end{cases}$$

From the (1), we deduce:

$$(x_1 - x_\Omega) \times (x_4 - x_\Omega) + (y_1 - y_\Omega) \times (y_4 - y_\Omega) + (z_1 - z_\Omega) \times (z_4 - z_\Omega) = 0$$

But $x_\Omega = y_\Omega = 0$, $x_4 = \frac{l}{2}$, $y_4 = 0$ and $z_1 = z_\Omega$

Then,

$$x_1 \times \frac{l}{2} = 0$$

So, $x_1 = 0$

From (2), we deduce that:

$$(x_1 - x_\Omega)^2 + (y_1 - y_\Omega)^2 + (z_1 - z_\Omega)^2 = 1$$

But $x_1 = x_\Omega = 0$, $y_\Omega = 0$, $z_1 = \sqrt{1 - \frac{x_1^2 + y_1^2}{R^2}}$ cf. (3) and $z_\Omega = \frac{\sqrt{(2R-l)(2R+l)}}{2R}$ Then,

$$\begin{aligned} y_1^2 + \left(\sqrt{1 - \frac{y_1^2}{R^2}} - \frac{\sqrt{(2R-l)(2R+l)}}{2R} \right)^2 &= 1 \\ \Leftrightarrow y_1^2 + \left(\frac{2\sqrt{R^2 - y_1^2} - \sqrt{4R^2 - l^2}}{2R} \right)^2 &= 1 \\ \Leftrightarrow y_1^2 + \frac{4(R^2 - y_1^2) + (4R^2 - l^2) - 4\sqrt{(R^2 - y_1^2)(4R^2 - l^2)}}{4R^2} &= 1 \\ \Leftrightarrow y_1^2 \times 4R^2 + 4(R^2 - y_1^2) + (4R^2 - l^2) - 4\sqrt{(R^2 - y_1^2)(4R^2 - l^2)} - 4R^2 &= 0 \end{aligned}$$

If we pass the square root to the other side and then square each sides, we will get the equation (E):

$$\begin{aligned} (y_1^2 \times 4R^2 + 4(R^2 - y_1^2) + (4R^2 - l^2) - 4R^2)^2 &= \left(4\sqrt{(R^2 - y_1^2)(4R^2 - l^2)} \right)^2 \\ \Leftrightarrow y^4 \times (R^2 - 1)^2 + y^2 \times \left(4R^2 - l^2 - 2 \times \left(R^2 - \left(\frac{l}{2} \right)^2 \right) (R^2 - 1) \right) &+ \left(\left(R^2 - \left(\frac{l}{2} \right)^2 \right)^2 - 4R^4 + (Rl)^2 \right) = 0 \end{aligned}$$

By plugging the equation in wolfram, we get : Wolfram Solution of E

You can have a glance on the solutions on the figure.

N.B. Why do we have four solutions ? Use sage math

We will take :

$$y_1 = -\sqrt{\frac{l^2 \times (R^2 + 1) + 4R^2 \times (R^2 - 1)}{4(R^4 - 2R^2 + 1)}} - \frac{\sqrt{l^4 R^2 - 4l^2 R^6 - 4l^2 R^2 + 16R^8 - 16R^6 + 16R^4}}{2 \times (R^4 - 2R^2 + 1)}$$

$$\text{So } z_1 = \sqrt{1 - \frac{\frac{l^2 \times (R^2 + 1) + 4R^2 \times (R^2 - 1)}{4(R^4 - 2R^2 + 1)} - \frac{\sqrt{l^4 R^2 - 4l^2 R^6 - 4l^2 R^2 + 16R^8 - 16R^6 + 16R^4}}{2 \times (R^4 - 2R^2 + 1)}}{R^2}}$$

And we have $P_1(0, y_1, z_1)$

We will stop the calculations here because it is too heavy and we will just prove the impossibility to have a grounded table.

P_2 and P_3 are distant from Ω of $\frac{l}{2} < 1$ whereas P_1 and P_3 are distant from Ω of 1, thus there are more far away. Then $z_1 < z_\Omega$ and $z_3 < z_\Omega$ because as the further away we get away from the center and the more lower we go. If we name d the distance from (Oz) , we have $x^2 + y^2 = d^2$, so $z = \sqrt{1 - \frac{d^2}{R^2}}$. If d goes up, z goes down.

But here is the contradiction: it is impossible to have a segment that passes through three distinct points where the altitude of one of them is different from the other two.

Thus we cannot have a stabilised table.

0.4 Any convex quadrilateral

The section of a sphere by a plan is a circle. Then the quadrilateral has to be in a circle to stand on a sphere. If the convex quadrilateral cannot stand in a circle, he cannot stand on a sphere.

The convex quadrilateral is contained in a plane. Then any quadrilateral table is grounded on a plane.

Question 2

We will which of the above quadrilaterals could the table be wobbly or/and stabilised.

$$y = - \sqrt{\left(\frac{l^2 R^2}{4(R^4 - 2R^2 + 1)} + \frac{l^2}{4(R^4 - 2R^2 + 1)} - \frac{\sqrt{l^4 R^2 - 4l^2 R^6 - 4l^2 R^2 + 16R^8 - 16R^6 + 16R^4}}{2(R^4 - 2R^2 + 1)} - \frac{R^4}{R^4 - 2R^2 + 1} - \frac{R^2}{R^4 - 2R^2 + 1} \right)}$$


$$y = \sqrt{\left(\frac{l^2 R^2}{4(R^4 - 2R^2 + 1)} + \frac{l^2}{4(R^4 - 2R^2 + 1)} - \frac{\sqrt{l^4 R^2 - 4l^2 R^6 - 4l^2 R^2 + 16R^8 - 16R^6 + 16R^4}}{2(R^4 - 2R^2 + 1)} - \frac{R^4}{R^4 - 2R^2 + 1} - \frac{R^2}{R^4 - 2R^2 + 1} \right)}$$

$$y = - \sqrt{\left(\frac{l^2 R^2}{4(R^4 - 2R^2 + 1)} + \frac{l^2}{4(R^4 - 2R^2 + 1)} + \frac{\sqrt{l^4 R^2 - 4l^2 R^6 - 4l^2 R^2 + 16R^8 - 16R^6 + 16R^4}}{2(R^4 - 2R^2 + 1)} - \frac{R^4}{R^4 - 2R^2 + 1} - \frac{R^2}{R^4 - 2R^2 + 1} \right)}$$

$$y = \sqrt{\left(\frac{l^2 R^2}{4(R^4 - 2R^2 + 1)} + \frac{l^2}{4(R^4 - 2R^2 + 1)} + \frac{\sqrt{l^4 R^2 - 4l^2 R^6 - 4l^2 R^2 + 16R^8 - 16R^6 + 16R^4}}{2(R^4 - 2R^2 + 1)} - \frac{R^4}{R^4 - 2R^2 + 1} - \frac{R^2}{R^4 - 2R^2 + 1} \right)}$$

Enlarge Data Customize Plain Text

14: Solutions from Wolfram Alpha for y



15: Slice of the 3D representation

0.5 Stabilised

On a plane, the table is always stabilised. Because the table is contained in a plane.
For a sphere, if the table is contained in a circle, then the table is stabilised.

0.6 Wobbly

For a sphere, if the table cannot be contained in a circle, then the table is wobbly. Because the section of a sphere by a plane is a circle.

Question 3

0.7 Inclined Plane

Our table is on the plane $P : ax + by - z = 0$. A normal vector of this plane is $\vec{n} = \begin{pmatrix} a \\ b \\ -1 \end{pmatrix}$

And a normal vector of the horizontal plane $P' : z = 0$ is $\vec{n}' = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$

The angle α between the two planes is defined as follow:

$$\cos \alpha = \frac{|\vec{n} \cdot \vec{n}'|}{\|\vec{n}\| \times \|\vec{n}'\|}$$

Here, we have:

$$\cos \alpha = \frac{|\vec{n} \cdot \vec{n}'|}{\|\vec{n}\| \times \|\vec{n}'\|} = \frac{|-1|}{\sqrt{a^2 + b^2 + 1} \times 1} = \frac{1}{\sqrt{a^2 + b^2 + 1}}$$

Then, $\alpha = \arccos\left(\frac{1}{\sqrt{a^2 + b^2 + 1}}\right)$