

Technical Verification Companion to the ECH Spin-Torsion Program: Λ CDM+ ΔN_{eff} MCMC Proxy, NaMaster Pipeline Recovery, and a Birefringence Consistency Check with a Spectator-ALP Model

Houston Golden^{1,*}

¹*Independent Researcher, Los Angeles, California, USA*

(Dated: June 9, 2026 PDT)

We report the technical verification material for the Einstein-Cartan-Holst (ECH) spin-torsion cosmology no-go program of Paper I(a) [1]. Three analyses are documented. (1) *Stock-CAMB Λ CDM+ ΔN_{eff} MCMC proxy* (Cobaya v3.6.1, **309,189** frozen samples across two converged dataset combinations, plus a third Planck-only combination still accumulating): this run uses stock CAMB with ΔN_{eff} as a free parameter and carries *no torsion modifications to the Boltzmann equations*; it is reported as a null-consistency test of an extra radiation-like degree of freedom, not as evidence for or against the ECH spin-torsion framework. Both frozen dataset combinations find ΔN_{eff} consistent with zero (-0.020 ± 0.169 full-tension; $+0.065 \pm 0.17$ Planck+BAO+SN) and H_0 consistent with standard Λ CDM (67.68 ± 1.06 full-tension; 67.79 ± 1.09 Planck+BAO+SN, both in $\text{km s}^{-1} \text{Mpc}^{-1}$). (2) *NaMaster pseudo- C_ℓ pipeline validation* on synthetic Λ CDM CMB polarization skies with an ACT-like footprint mask ($N_{\text{side}} = 512$, $\ell_{\text{max}} = 1024$, $f_{\text{sky}} = 0.32$, $10 \mu\text{K} \cdot \text{arcmin}$ white noise, 500 Monte Carlo realizations): injecting the spectator-ALP fiducial value $\beta = 0.27^\circ$ recovers $\hat{\beta} = 0.238^\circ$ (pipeline-recovery bias $\hat{\beta} - \beta_{\text{inj}} = -0.032^\circ$). *Scope of the validation*: the test confirms the algebraic pseudo- C_ℓ $E \rightarrow B$ deconvolution under MASTER mode coupling, NOT the physical separation of the cosmic-rotation angle β from the instrumental-miscalibration angle α which strictly requires unrotated galactic foregrounds (the synthetic CMB-only skies contain no galactic foregrounds, so the very component that breaks the β - α degeneracy in published Planck/ACT DR6 measurements is absent by construction). The MC recovery is therefore a pipeline-validation figure, not a sky-detection significance claim. The primary sky detection significance is the published Planck/ACT DR6 2.7–2.9 σ [2, 3];^a the pipeline SNR figures refer to recovery of injected MC signals and are *not* competitive sky measurements. (3) *Spectator-ALP consistency check*: a field with $f_a \sim M_{\text{Pl}}$, $m \sim H_0$ is consistent with the published joint WMAP+Planck value $\beta = 0.342^\circ \pm 0.094^\circ$ (3.6 σ) [2]. *Spectator-status caveat*: for $\theta_i \sim 1$ the ALP energy density $\rho_a \sim m^2 f_a^2 \theta_i^2 \sim H_0^2 M_{\text{Pl}}^2$ is of order the critical density today, so the spectator label is only consistent under the explicit restriction $\theta_i \ll 1$ (which constitutes fine-tuning of the misalignment initial condition); at $\theta_i \sim 1$ the ALP must instead be treated as the dark-energy field itself, which is outside the scope of this companion-paper consistency check. The same birefringence arises in standard GR with an identical ALP; it is not a distinctive ECH prediction, and the spectator-status fine-tuning is required regardless of whether the underlying cosmology is a bounce or Λ CDM. A reproducibility manifest is included in Appendix A.

PACS numbers: 98.80.-k, 95.36.+x, 04.50.Kd

CONTENTS

I. Introduction	2	III. Stock-CAMB Λ CDM+ ΔN_{eff} MCMC: Generic Radiation-Proxy Test (Not a Spin-Torsion Theory Module)	2
II. Cosmological Tensions: H_0 and σ_8	2	IV. Data Methods: CMB E - B Analysis	5
		V. Cosmological Fits and Model Comparison	7
		A. Datasets and Configuration	7
		B. Results	7
		VI. Cosmic Birefringence: Spectator ALP Consistency Check	8
		VII. Conclusions	10
		Data and Code Availability	11
		Acknowledgments	11
		A. Reproducibility Materials	11

* houston@hubify.com

^a Eskilt & Komatsu 2022 disambiguation: the published PRD paper [2] (PRD 106:063503, arXiv:2205.13962) analyzes *Planck PR3 + WMAP9*; the public reproduction code released by the authors at github.com/LilleJohs/Cosmic_Birefringence was subsequently updated to use *Planck PR4 / NPIPE*. Throughout this paper, the labels “PR4/NPIPE” attached to the Eskilt+Komatsu likelihoods refer to the code-repository dataset (which is what the ALP-MCMC re-runs actually use); the abstract $\beta = 0.342^\circ \pm 0.094^\circ$ (3.6 σ) headline is from the published PR3+WMAP9 joint analysis. The repository README is the authoritative source for the dataset attribution in the executed pipeline.

B. Claims Classification	12
C. ALP-MCMC Sampled Parameters, Priors, and Likelihood Stack	12
References	12

I. INTRODUCTION

This companion paper provides the technical verification layer for the ECH structural-closure no-go result reported in Paper I(a) [1]. The main paper establishes 14 independent structural constraints closing minimal-ECH dark-energy routes and proves a perturbation-transparency theorem for the Holst sector. The present paper documents the three numerical analyses that support and contextualize those results.

Scope of this paper.—Three verification analyses are reported here, each with explicit scope limitations:

1. Λ CDM+ ΔN_{eff} MCMC proxy (Secs. II and III): stock CAMB Boltzmann solver with ΔN_{eff} as a free parameter. **Not a spin-torsion theory module.** The Boltzmann code carries no torsion modifications; this run tests whether the data prefer an extra radiation-like degree of freedom in standard cosmology.
2. NaMaster CMB E-B analysis (Sec. IV): a bias-injection Monte Carlo validation of the pseudo- C_ℓ deconvolution pipeline on synthetic Λ CDM polarization skies. **Not a competitive sky detection.** The high pipeline template-fit SNR figures (e.g., 20.32) refer to recovery of injected MC signals, not to the significance of the CMB sky measurement, which remains the published 2.7–2.9 σ from Planck/ACT.
3. Spectator-ALP consistency check (Sec. VI): a standard GR+ALP computation showing that the observed signal is consistent with an ALP having natural parameters (taken at scan-prior midpoint values; the $\sim 25\times$ misalignment tuning required to reconcile the headline result with the spectator-consistent corner is disclosed in Sec. VI and fn. 5). **Not a distinctive ECH prediction.** The same $\beta \approx 0.27^\circ$ arises in any GR+ALP setup with the same parameters; no ECH-specific derivation connects the Holst action to the photon-torsion coupling required.

What is NOT in this paper.—The 13 logically-independent structural barriers, the perturbation-transparency theorem, the 14-barrier table, and the surviving matter-bounce-specific test predictions ($f_{\text{NL}} = -35/8$, valid strictly for the minimal scalar-only $w = 0$ matter-dominated contraction class, not the broader bouncing-cosmology landscape which encompasses ekpyrotic, Cuscuton, string-gas, and quintom variants with different f_{NL} signatures, and ALP birefringence) are in Paper I(a) [1]. The SPHEREx multi-tracer Fisher forecast is in Paper II [4]. The multi-survey anomaly catalog

is in Paper III [5]. The galaxy chirality catalog is in Paper IV [6].

Cross-paper citations.—When this companion reports MCMC values (H_0 , σ_8 , etc.) that are referenced in the main paper, those values come from Secs. III and V here.

II. COSMOLOGICAL TENSIONS: H_0 AND σ_8

The bounce scenario motivates extending Λ CDM by ΔN_{eff} (particle production at the bounce) as a phenomenological proxy parameter; $(\omega/H)_0$ (angular momentum transfer) and Ω_k are fixed to zero in the actual sampled MCMC configuration of this paper (per the explicit parameter-scope clarification in Sec. VA; the $(\omega/H)_0$ parameter is discussed in Paper I(a) as a phenomenological bounce-class indicator but is not separately sampled here). The full-tension dataset combination includes the SH0ES [7] M_B -anchor likelihood (H0.riess2020Mb, entering through the supernova absolute magnitude rather than as a direct H_0 prior) in the Cobaya likelihood configuration, but the high-precision Planck NPIPE (PR4) CamSpec high- ℓ TT-TEEE + Planck 2018 low- ℓ TT/EE + Planck 2018 lensing likelihoods carry sufficient inverse-variance weight that the posterior H_0 in the proxy run is pulled to 67.68 ± 1.06 (Planck-dominated value) rather than to the simple Gaussian-combination value ~ 70 that would emerge if SH0ES and Planck were equally weighted. We do not therefore claim that the SH0ES tension is resolved or even moved by adding ΔN_{eff} in stock CAMB; in this configuration the ΔN_{eff} posterior is consistent with zero and the H_0 posterior is dominated by Planck. The spin-torsion framework alone does not resolve cosmological tensions at the present data precision.

III. STOCK-CAMB Λ CDM+ ΔN_{eff} MCMC: GENERIC RADIATION-PROXY TEST (NOT A SPIN-TORSION THEORY MODULE)

Scope statement.—“MCMC verification” refers throughout to a stock CAMB run of Λ CDM extended by ΔN_{eff} as a free parameter. *No custom CAMB modifications are used; no torsion-modified Boltzmann equations are solved.* The MCMC therefore tests whether the data prefer an extra radiation-like degree of freedom, treated as a generic phenomenological proxy for the spin-torsion sector’s possible effective radiation contribution. It does *not* verify the spin-torsion theory module itself; that would require a bespoke modified Boltzmann code.

The proxy run (Cobaya v3.6.1 with Planck NPIPE (PR4) CamSpec high- ℓ TTTEEE + Planck 2018 low- ℓ TT/EE + Planck 2018 lensing likelihoods, i.e. `planck_NPIPE_highl_CamSpec.TTTEEE + planck_2018_lowl.TT + planck_2018_lowl.EE + planck_2018_lensing.click` in the Cobaya YAML) has produced two frozen dataset combinations with

publication-quality convergence, plus a third Planck-only run currently at sub-convergence sample count and not aggregated into any frozen-posterior summary statistic in this paper. Frozen MCMC program: **309,189** raw samples across 2 frozen dataset combinations (176,240 + 132,949). We report this as a Λ CDM+ ΔN_{eff} null-consistency test: the data are consistent with $\Delta N_{\text{eff}} = 0$ in stock CAMB. Sampling configuration (per the on-disk Cobaya YAMLs): ΔN_{eff} enters as `nnu` with a flat prior $N_{\text{eff}} \in [2.046, 5.046]$ (i.e. $\Delta N_{\text{eff}} \in [-1, +2]$); the neutrino mass is held at the CAMB default $\Sigma m_\nu = 0.06$ eV with one massive eigenstate (`num_massive_neutrinos=1`), and Y_{He} follows the CAMB BBN-consistent default (no explicit override in any of the four YAMLs).¹

a. Scope of the ΔN_{eff} proxy: bounce-class discrimination, not a direct test of the spin-torsion sector. We emphasize that the stock-CAMB Λ CDM+ ΔN_{eff} proxy run does *not* test the ECH spin-torsion sector directly. The Hehl-Datta-Mercuri parity-even four-fermion contact interaction that survives torsion elimination [8, 9] is dimension-6 and M_{Pl}^{-2} -suppressed²: its leading Boltzmann effect is a scattering-amplitude shift, not a relativistic species, and it does *not* produce a ΔN_{eff} at recombination. The *minimal* matter-bounce class [10] (here “minimal” = the single-scalar $w = 0$ matter-dominated contraction phase, distinguished from the broader matter-bounce family including ekpyrotic, cyclic, Cuscuton, string-gas, and quintom-bounce variants) predicts $\Delta N_{\text{eff}} \approx 0$ by construction (no light bounce-internal species are thermalized at recombination); the proxy run *confirming* $\Delta N_{\text{eff}} = -0.020 \pm 0.169$ (full-tension) and

TABLE I. Λ CDM+ ΔN_{eff} proxy MCMC results from frozen Cobaya v3.6.1 chains (CAMB v1.6.5, stock; *no torsion modifications*). All values are posterior means $\pm 1\sigma$. Reported as a null-consistency cross-check, *not* as evidence for the spin-torsion theory.

Parameter	Full-tension	Planck+BAO+SN
H_0 [km/s/Mpc]	67.68 ± 1.06	67.79 ± 1.09
ΔN_{eff}	-0.020 ± 0.169	$+0.065 \pm 0.17$
σ_8	0.803 ± 0.008	0.812 ± 0.009
S_8	0.814 ± 0.008	0.831 ± 0.018
Ω_m	0.308 ± 0.005	0.312 ± 0.006
τ	0.054 ± 0.007	0.056 ± 0.007
n_s	0.965 ± 0.006	0.967 ± 0.006
Chains	6	6
Total samples	176,240	132,949
Worst $\hat{R} - 1^a$	0.001	0.003
Min ESS	4,744	4,692

^a Worst row is n_s in the full-tension combination at $\hat{R} - 1 = 9.74 \times 10^{-4}$; all sampled parameters across both frozen combinations satisfy $\hat{R} - 1 < 3 \times 10^{-3}$. The full-tension chain samples 17 parameters: 7 cosmological + 9 Planck likelihood nuisance (A_{planck} , amp_{143} , amp_{217} , $\text{amp}_{143 \times 217}$, n_{143} , n_{217} , $n_{143 \times 217}$, calTE , calEE) + 1 supernova absolute magnitude (M_b , entering via the Pantheon+ SN and SH0ES H0.riess2020Mb likelihoods rather than the Planck likelihood stack). The Planck+BAO+SN chain samples 16 parameters (7 cosmological + the same 9 Planck nuisance; M_b is not explicitly sampled there because `sn.pantheonplus` runs with `use_abs_mag: false`, analytically marginalizing the absolute magnitude in the absence of the SH0ES M_b likelihood; verified by direct chain-header / `.updated.yaml` blocking-list inspection). Full per-parameter convergence tables ($\hat{R}$, ESS, drift, per dataset) are archived at `reproducibility/cosmology/convergence_latest.csv` and mirrored in the HuggingFace chain-diagnostics dataset (Appendix A).

¹ Sample-count stratification (reconciliation): **309,189** is the sum of the two frozen combinations (176,240 + 132,949 raw accepted samples). After removing the first 30% of each chain as burn-in, $176,240 \times 0.7 + 132,949 \times 0.7 \approx 216,432$ post-burnin samples remain across both frozen chains (`convergence_summary.json`). For the full-tension subset specifically, $176,240 \times 0.7 \approx 123,368$ post-burnin (the **119,617** figure in Fig. 1 reflects additional `getdist` effective-sample weight-based thinning of this subset only). The post-burnin count of the full-tension subset alone is 123,129 (within $\pm 1\%$ of the 123,368 exact computation, with the small offset reflecting the chain-end-truncation of partial samples at the burn-in cut); the correct both-chains post-burnin total is 216,432. The third (Planck-only) dataset combination (114,992 raw samples; $\hat{R} - 1 \sim 0.05$) is still accumulating samples, is *not* reported in Table I (which contains only the two frozen combinations), and is *not* aggregated into the 309,189-sample headline anywhere in this paper.

² The dimension-6 four-fermion contact operator is interpreted here in the low-energy effective field theory (EFT) below the strong-coupling / torsion-resolution scale $\Lambda_{\text{strong}} \sim M_{\text{Pl}}/\sqrt{\gamma_{\text{BI}}}$ set by the inverse Barbero-Immirzi parameter γ_{BI} (cf. [9], where the contact interaction is derived; the strong-coupling scale itself is a heuristic EFT-validity estimate); above this scale the contact-operator description breaks down and the full Holst-sector dynamics with propagating torsion must be retained. All recombination-era and late-time observables addressed in this paper sit at energies $E \ll \Lambda_{\text{strong}}$, where the EFT treatment is controlled.

+0.065 $\pm$ 0.17 (Planck+BAO+SN) is therefore consistent with the minimal matter-bounce prediction, but the absence of a ΔN_{eff} detection is not a discriminator between minimal-ECH and standard Λ CDM at the present data precision. We frame the proxy as a bounce-class *compatibility check* (minimal matter-bounce-class scenarios without prolonged post-bounce inflation predict $\Delta N_{\text{eff}} \approx 0$), not as a posterior-preference test against a competing model.

Physics interpretation (Table II).—The converged $w_0 w_a$ posterior *disfavors* (in the marginal-tail sense; see fn. a) the LCDM point $(w_0, w_a) = (-1, 0)$ at the joint level: w_0 departs by $+4.3\sigma$ and w_a departs by -3.6σ , with $w_0 + w_a = -1.48 \pm 0.15$ requiring phantom crossing in the redshift range probed by DESI DR2 BAO + DES-Y5 + Pantheon+. This is the canonical quintom signature and is consistent with the bounce / pre-Big-Bang scenario discussed in Paper I(a)’s § Structural Tension as a constraint that single-field quintessence cannot accommodate without a second degree of freedom (phantom / quintom-B). In the converged chain there are zero free- $w_0 w_a$ samples at the LCDM point; the chain is centered well into quintom-B territory at $w_0 + w_a \approx -1.48$.

TABLE II. DESI DR2 $w_0 w_a$ posterior summary ($N = 128,385$ accepted samples across 16 chains, $\hat{R} - 1 = 0.00820$; 8 cosmological + 9 nuisance parameters). Likelihood stack: DESI DR2 BAO + Planck NPIPE (PR4) CamSpec high- ℓ TTTEEE + Planck 2018 low- ℓ TT/EE + Planck 2018 lensing (**native**) + DES-Y5 + Pantheon+. Full parameter list and chain diagnostics are available in the companion data repository.

Parameter	Mean $\pm \sigma$	vs LCDM
<i>Dark-energy extension</i>		
w_0	-0.8122 ± 0.0436	(marg.-tail, $+4.3\sigma$) ^a
w_a	-0.6666 ± 0.1864	-3.6σ from 0
$w_0 + w_a$	-1.4788 ± 0.1485	phantom-crossing required
w_{pivot}	-0.952 ± 0.019	$+2.5\sigma$ from -1 ^b
<i>Standard cosmology</i>		
H_0 [km/s/Mpc]	67.185 ± 0.455	—
Ω_m	0.3142 ± 0.0045	—
σ_8	0.8057 ± 0.0083	—
S_8	0.8245 ± 0.0089	—
n_s	0.9655 ± 0.0036	—
$10^9 A_s$	2.087 ± 0.030	—
τ	0.0537 ± 0.0072	—
$\omega_b h^2$	0.02224 ± 0.000125	—
$\omega_c h^2$	0.11892 ± 0.000819	—
$100\theta_{\text{MC}}$	1.04087 ± 0.000239	—
Age [Gyr]	13.763 ± 0.019	—
<i>Goodness-of-fit decomposition</i>		
χ^2_{total}	14037.4 ± 5.6^c	—
χ^2_{BAO}	10.6 ± 1.8	DESI DR2 ^d
χ^2_{CMB}	10983.9 ± 5.3	Planck PR4 + lensing
χ^2_{SN}	3043.0 ± 1.6	DES-Y5 + Pantheon+

^a Marginal-tail *posterior-extrapolation* departure: LCDM is unsampled by this chain (no $w_0 = -1, w_a = 0$ samples in the present Metropolis-Hastings run), so the $+4.3\sigma$ figure is a posterior-tail extrapolation distance only, *not* a Bayes-factor or $\ln B$ exclusion and *not* a frequentist tension. A Savage-Dickey readout is not viable at this tail depth; robust $\ln B$ is left to a follow-up nested-sampling analysis (see § Headline-result discussion).

^b $w_{\text{pivot}} \equiv w_0 + (1 - a_p) w_a$ with the pivot a_p chosen so that w_{pivot} and w_a are uncorrelated in the posterior covariance, i.e. $\text{Cov}(w_0, w_a) + (1 - a_p) \text{Var}(w_a) = 0$, giving $1 - a_p = -\text{Cov}(w_0, w_a) / \text{Var}(w_a)$. Direct readout of the converged iter2 chain (30% burn-in): $\text{Cov}(w_0, w_a) = -0.00729$ (correlation coefficient $\rho = -0.90$), so $1 - a_p = 0.210$, $a_p = 0.790$, and $z_p = 1/a_p - 1 = 0.27$; this pivot is internal to the dataset stack (DESI DR2 BAO + Planck NPIPE + DES-Y5 + Pantheon+) and is not the literature de Putter–Linder $z_p \approx 0.4$ value. The decorrelated combination has $\sigma_{w_{\text{pivot}}}^2 = \sigma_{w_0}^2 - \text{Cov}^2(w_0, w_a) / \sigma_{w_a}^2 = (0.0436)^2 - (0.00729 / 0.1864)^2 = (0.0193)^2$, reproducing the ± 0.019 value above, with mean $w_{\text{pivot}} = -0.8122 + 0.210 \times (-0.6666) = -0.952$. The choice of a_p depends on the dataset stack; rerunning the chain with SH0ES or DESI DR2 BAO alone would shift z_p and w_{pivot} at the linear-Fisher level.

^c The mean-of-total χ^2 here is GetDist’s weighted-sample average over the full posterior, which differs from the sum of the individual-channel means ($10.6 + 10983.9 + 3043.0 = 14037.5$) by a 0.1-unit arithmetic-rounding artifact; the two are formally identical to within sampling precision.

^d The DESI DR2 BAO likelihood (`baos.desi_dr2.desi_bao_all`) comprises 13 BAO data points, so $\chi^2_{\text{BAO}} / N \approx 0.8$ — an unremarkable goodness of fit.

At the decorrelation pivot ($z_p = 0.27$; fn. b), the effective equation of state is $w_{\text{pivot}} = -0.952 \pm 0.019$, i.e. $+2.5\sigma$ above -1 : the departure from LCDM persists, at reduced significance, even in the best-determined direction of the (w_0, w_a) posterior, with the largest single-axis departures carried by w_0 ($+4.3\sigma$) and the time-varying w_a term (-3.6σ).

Caveats.—(a) The robust Bayesian evidence / Bayes factor $\ln B$ against LCDM is NOT reported here; standard posterior MCMC samples do not give a robust $\ln B$, and a Savage-Dickey density ratio at the LCDM point is *not* viable on this chain because the LCDM point $(w, w_a) = (-1, 0)$ lies at $> 4\sigma$ in the joint marginal tails (Table II: w_0 departs by $+4.3\sigma$ and w_a by -3.6σ) and is therefore unsampled by the Metropolis-Hastings

chain; any KDE-based Savage-Dickey ratio at an unsampled point yields arbitrary kernel-dependent noise rather than a controlled posterior density (with zero free- $w_0 w_a$ samples at the LCDM point, a KDE estimator on the converged 2D (w, w_a) marginal fails catastrophically). The robust $\ln B$ recompute therefore requires dedicated nested sampling (e.g., PolyChord or MultiNest) or thermodynamic integration on the same likelihood stack rather than a KDE readout from the converged Metropolis-Hastings chain; robust $\ln B$ computation is left to a follow-up nested-sampling analysis. (b) τ is constrained empirically by the low- ℓ EE + TT likelihoods listed in the caption (`planck_2018_low1.EE` + `planck_2018_low1.TT`) and is sampled as a free parameter rather than fixed by a Gaussian prior at the

Planck best-fit value (the chain has τ free with the standard low- ℓ data constraint, not a Gaussian prior at the Planck best-fit). (c) The SH0ES H_0 prior is *not* included in this chain (BAO + CMB + SN-only, no local-distance ladder); the joint posterior $H_0 = 67.185 \pm 0.455$ km/s/Mpc is therefore the no-SH0ES result. The interaction with the SH0ES prior is the subject of the separate *full-tension* chain reported in Table I, which carries the `H0.riess2020Mb` likelihood active in its YAML (verified by direct `.input.yaml` inspection and by chain sample-mean readout: $M_B = -19.263 \pm 0.049$ mag, agreeing with the Riess+2020 SH0ES value $M_B = -19.253 \pm 0.027$ mag at 0.2σ). The full-tension chain returns $H_0 = 67.68 \pm 1.06$ km/s/Mpc with $\Delta N_{\text{eff}} = -0.020 \pm 0.169$, exhibiting the canonical 3.6σ Hubble tension with Riess $H_0 = 73.04 \pm 1.04$ km/s/Mpc that the $\Lambda\text{CDM} + \Delta N_{\text{eff}}$ extension is unable to resolve: the `H0.riess2020Mb` likelihood constrains the SN Ia absolute magnitude M_B (which is correctly pulled toward the Riess value), but H_0 is determined predominantly by the BAO+CMB acoustic scale through Pantheon+, and the ΔN_{eff} extension lacks the degrees of freedom to shift H_0 enough to accommodate both data sets. Direct YAML inspection confirms the SH0ES likelihood is active and the M_B posterior is correctly pulled; the 3.6σ tension persists because the model cannot shift H_0 enough to satisfy both data sets simultaneously — the canonical Hubble-tension result, not a YAML omission. (d) The `iter2` chain's $S_8 = 0.8245 \pm 0.0089$ (Table II) sits 2.5σ above the DES-Y3 weak-lensing value $S_8 = 0.776 \pm 0.017$ [11] ($\Delta = 0.049$ against a combined uncertainty of 0.019): the S_8 tension is not relieved in this chain either.

M_B - H_0 joint-posterior offset check. The joint posterior mean ($M_B = -19.263$, $H_0 = 67.68$) is fully consistent with an active `sn.pantheonplus` likelihood: `sn.pantheonplus` enforces a soft constraint on the combination $M_B - 5\log_{10}(H_0) \approx \text{const}$ along the SN distance-modulus degeneracy. At the Riess+2020 anchor ($M_B = -19.253$, $H_0 = 73.04$), the constant is $-19.253 - 5\log_{10}(73.04) = -28.571$. At the chain joint mean ($M_B = -19.263$, $H_0 = 67.68$), the constant is $-19.263 - 5\log_{10}(67.68) = -28.415$, i.e. a 0.156 mag offset from the Riess anchor along the Pantheon+ constraint axis. This offset is $\sim 3.2\sigma$ relative to the chain's $\sigma_{M_B} = 0.049$ marginal width and is the same Hubble tension manifesting in the M_B axis (3.2σ in chain- σ units, versus the canonical 3.6σ when the tension is expressed in distance-ladder terms in the H_0 axis). The chain posterior is therefore the correct compromise of three inputs: the Riess M_B -prior preferring $M_B = -19.253$ ($\sigma = 0.027$); the Pantheon+ likelihood preferring $M_B - 5\log_{10} H_0 = -28.571$ along the SN degeneracy; and the Planck-CMB-likelihood preferring $H_0 \approx 67.5$ via the acoustic-scale calibration. With ΔN_{eff} bounded near zero by the joint data, the model cannot resolve the tension and the chain settles into the 3.6σ -discrepant joint posterior — NOT a YAML alias failure; the parameters are correctly aliased per the `spin.torsion.input.yaml`




FIG. 1. Full-tension MCMC corner plot (119,617 post-burnin samples, `getdist`-thinned from 176,240 raw; footnote 1) over Planck+BAO+SN+ H_0 + S_8 . The ΔN_{eff} posterior is consistent with zero (-0.020 ± 0.169), confirming no additional relativistic species at recombination.

configuration (Mb is a single nuisance parameter sampled jointly by both `sn.pantheonplus` and `H0.riess2020Mb`).

Key finding.—Both frozen datasets find ΔN_{eff} consistent with zero and H_0 consistent with Planck ΛCDM at 0.3σ , confirming that the ΔN_{eff} extension alone does not resolve the Hubble tension. Current data neither require nor exclude a small positive ΔN_{eff} from the spin-torsion sector; CMB-S4 ($\sigma(N_{\text{eff}}) \sim 0.03$) will provide the first precision test.

Independent cross-validation.—Liu *et al.* [12] constrained an EC torsion model using DESI DR2 [13] + Pantheon+ [14] + DES-SN5YR [15] + Planck 2018, finding torsion preferred by AIC ($\Delta\text{AIC} = -5.7$ to -6.6) but with the torsion parameter itself consistent with zero ($\alpha = -0.00066 \pm 0.00098$). Their headline values $H_0 = 68.41 \pm 0.32$ km/s/Mpc and $S_8 = 0.812 \pm 0.006$ agree with our Planck+BAO+SN chain at 0.5σ in H_0 ($|67.79 - 68.41|/\sqrt{1.09^2 + 0.32^2}$) and 1.0σ in S_8 ($|0.831 - 0.812|/\sqrt{0.018^2 + 0.006^2}$); their torsion-consistent-with-zero result parallels our ΔN_{eff} -consistent-with-zero null finding.

IV. DATA METHODS: CMB E - B ANALYSIS

Birefringence measurements are adopted from the published literature: $\beta = 0.30^\circ \pm 0.11^\circ$ (Planck NPIPE [16]) and $\beta = 0.215^\circ \pm 0.074^\circ$ (ACT DR6 [3]). The spectrator ALP analysis (Sec. VI) uses these published values.




FIG. 2. ΔN_{eff} marginal posterior comparison across the two frozen dataset combinations of Table I (176,240 and 132,949 samples). Panel (a): Gaussian summaries of the ΔN_{eff} marginal posteriors at the Table I means $\pm 1\sigma$, with the Standard-Model value $\Delta N_{\text{eff}} = 0$ marked. Panel (b): all seven Table I parameters, normalized to the full-tension mean and σ . Both combinations recover ΔN_{eff} consistent with zero at $< 1\sigma$; the headline “full-tension” stack (Planck+BAO+SN+ H_0 + S_8) gives $\Delta N_{\text{eff}} = -0.020 \pm 0.169$. The standard-ECH route to dark energy via additional relativistic species at recombination is therefore not viable as an amplitude-level explanation of either tension.

Scope note.—The NaMaster pseudo- C_ℓ analysis below is a bias-injection Monte Carlo validation of the deconvolution pipeline on synthetic skies, not a cosmological measurement. The high pipeline template-fit SNR figures (e.g., 20.32, 25.71; fn. 3) refer to recovery of injected MC signals and must not be conflated with the published Planck/ACT DR6 2.7–2.9 σ sky detection.

Pipeline configuration.—The pseudo- C_ℓ analysis follows the NaMaster framework [17]. *Simulated skies.*—Each Monte Carlo realization is a synthetic Λ CDM CMB Q/U sky generated directly at $N_{\text{side}} = 512$ (healpy.synfast, harmonic content to $\ell_{\text{max}} = 2N_{\text{side}} = 1024$) from a semi-analytic fit to the Planck-2018 EE spectrum plus a lensing-like BB component ($C_\ell^{BB} = 0.05 C_\ell^{EE}$). No real Planck map enters the Monte Carlo, and no instrumental beam is applied (the synthetic skies and the recovery template share the same spectra, so a common beam would cancel in the β estimate).

Mask.—An ACT-like footprint (Galactic cut $|b| > 20^\circ$ plus declination cut $\text{dec} \in [-65^\circ, +25^\circ]$), apodized by Gaussian smoothing of the binary mask at 2° FWHM, giving $f_{\text{sky}} = 0.32$. No E/B purification is applied (purify_b=False); residual $E \rightarrow B$ leakage from the apodized mask is absorbed into the measured pipeline-recovery bias, which is the quantity this validation is designed to characterize.

Mode-coupling matrix and binning.—The $M_{\ell\ell'}$ matrix is computed via NmtWorkspace.compute_coupling_matrix on the apodized mask, retaining the full $EE/EB/BB$ block structure. Spectra are band-power-binned into 20 linear bins spanning $30 \leq \ell \leq 3N_{\text{side}} = 1536$ (bins above the map band limit $\ell = 1024$ carry noise only). Only this single binning/ ℓ -range configuration is exercised; an ℓ -range robustness sweep is not part of the present MC

suite.

Noise model and injections.—Each realization adds white noise at the ACT-like level $\Delta_P = 10 \mu\text{K} \cdot \text{arcmin}$ (a conservative worst-case bias check; no $1/f$ or anisotropic component). The $\beta = 0.27^\circ$, $\beta = 0.342^\circ$, and $\beta = 0$ injections rotate $Q + iU$ via $e^{2i\beta}(Q + iU)$. The angle is recovered per configuration by an unweighted χ^2 template fit of the decoupled C_ℓ^{EB} band-powers to $\sin(2\beta) \cos(2\beta) C_\ell^{EE}$ on a β grid. Only non-negative injections were run; a negative- β injection (sign-symmetry/linearity check) is not part of the present MC suite and is an acknowledged validation gap.

Reproducibility.—The canonical driver script, deterministic seeds (seed.base=42), and output summary are archived at reproducibility/pi_namaster_500mc/ (mirroring the executed pod run pipelines/h200_results/pod1_namaster_umap_2026-04-29/).

Production 500-realization run (April 2026).—We performed the NaMaster pseudo- C_ℓ Monte Carlo described above with 500 realizations per injection. Injecting the spectator-ALP fiducial $\beta = 0.27^\circ$ (consistent with, but not derived from, the ECH action) recovers:

$$\hat{\beta}_{\text{NaMaster}} = 0.238^\circ \quad (500\text{-MC sample mean of } \hat{\beta}). \quad (1)$$

The pipeline-recovery bias, defined throughout as $\Delta\hat{\beta} \equiv \hat{\beta} - \beta_{\text{inj}}$, is -0.032° (consistent with the apodized-mask power suppression expected from a 2° apodization scale).³ For $\beta = 0.342^\circ$ (the published joint WMAP+Planck value [2]), the pipeline recovers 0.302° at template-fit SNR = 25.71; for $\beta = 0$, the recovered angle is 0.000° with template-fit SNR 0.0 (null check; summary.json). The pipeline-recovery bias is $\Delta\hat{\beta} = -0.032^\circ$ at injection $\beta = 0.27^\circ$ ($\hat{\beta} = 0.238^\circ$) and $\Delta\hat{\beta} = -0.040^\circ$ at injection $\beta = 0.342^\circ$ ($\hat{\beta} = 0.302^\circ$). The absolute bias grows mildly ($\sim 25\%$) with injected amplitude, while the

³ The “pipeline-recovery SNR= 20.32” figure (and analogously 25.71 for the $\beta = 0.342^\circ$ injection below) is the template-fit SNR of the driver script, $\text{SNR}^{\text{tmpl}} \equiv [\sum_b (C_b^{EB,\text{th}}/\sigma_b^{\text{MC}})^2]^{1/2}$, where $C_b^{EB,\text{th}} = \sin(2\beta) \cos(2\beta) C_b^{EE}$ is the injected EB band-power template and σ_b^{MC} is the per-bin scatter across the 500 realizations — the matched-template significance of the injected signal against single-realization noise, the appropriate quantity for evaluating the deconvolution pipeline. It is *not* the significance of the recovered angle itself. The sky-fraction sweep artifact (c1_fsky_sweep.json) confirms the identification: its snr_template_canonical_def values 32.98 and 28.81 at $f_{\text{sky}} = 0.85$ and 0.65 equal $20.32 \sqrt{f_{\text{sky}}/0.32}$. The per-realization angle-recovery ratio $\hat{\beta}/\sigma_\beta$, with σ_β the per-realization scatter of the recovered angle, is measured in the same artifact as 8.1 ($\sigma_\beta = 0.029^\circ$, $f_{\text{sky}} = 0.85$) and 7.2 ($\sigma_\beta = 0.033^\circ$, $f_{\text{sky}} = 0.65$); σ_β was not recorded for the canonical $f_{\text{sky}} = 0.32$ run, but the $\sqrt{f_{\text{sky}}}$ scaling validated between the two measured points implies $\sigma_\beta(0.32) \approx 0.029^\circ \sqrt{0.85/0.32} \approx 0.047^\circ$ and hence $\hat{\beta}/\sigma_\beta \approx 0.238/0.047 \approx 5$ per realization at the canonical mask. The standard error of the 500-MC mean is smaller by $\sqrt{N} = 22.4$.




FIG. 3. NaMaster pipeline-recovery validation on synthetic Λ CDM polarization skies ($N_{\text{side}} = 512$, $\Delta_P = 10 \mu\text{K} \cdot \text{arcmin}$ white noise, $N = 500$ MC realizations per point; Sec. IV). Panel (a): 500-MC mean recovered $\hat{\beta}$ vs. injected $\beta_{\text{inj}} \in \{0, 0.27^\circ, 0.342^\circ\}$ at the canonical apodized $f_{\text{sky}} = 0.32$ ACT-like mask, annotated with the pipeline-recovery bias $\hat{\beta} - \beta_{\text{inj}}$ (0.000° , -0.032° , -0.040°). Panel (b): the $\beta_{\text{inj}} = 0.27^\circ$ sky-fraction sweep; outer (light) error bars are the per-realization scatter σ_β (0.029° at $f_{\text{sky}} = 0.85$, 0.033° at 0.65), inner bars the standard error of the 500-MC mean $\sigma_\beta/\sqrt{N}$; per-realization σ_β was not recorded in the canonical $f_{\text{sky}} = 0.32$ artifact, so that point carries the mean only. The worst-case $|\text{bias}| = 0.040^\circ$ is the NaMaster systematic floor adopted in Eq. 1–fn. 3. Underlying data artifacts are listed in the *Reproducibility* paragraph of Sec. IV.

multiplicative under-recovery is amplitude-independent at $\sim 12\%$ ($0.238/0.27 = 0.302/0.342 = 0.88$). The deconvolution is therefore unbiased at the $|\Delta\hat{\beta}| \leq 0.040^\circ$ level in the worst-case injection, which we carry forward as the NaMaster systematic floor; this is a methodology cross-check, not a competitive sky measurement.

Sky-fraction sweep.—Because published birefringence analyses use larger sky fractions than our canonical $f_{\text{sky}} = 0.32$ validation mask, we repeat the $\beta = 0.27^\circ$ injection-recovery exercise at $f_{\text{sky}} = 0.85$ (Planck-like) and $f_{\text{sky}} = 0.65$ (ACT-DR6-like) galactic-cut masks with the same apodization recipe, noise level, and $N = 500$ MC realizations ($N_{\text{side}} = 512$). The recovered values are $\hat{\beta} = 0.237^\circ$ ($f_{\text{sky}} = 0.85$, per-realization $\sigma_\beta = 0.029^\circ$) and $\hat{\beta} = 0.236^\circ$ ($f_{\text{sky}} = 0.65$, $\sigma_\beta = 0.033^\circ$), i.e. a recovery bias of -0.033° to -0.034° — statistically indistinguishable from the canonical-mask bias of -0.032° (the 0.001° difference is $0.8\times$ the standard error of the 500-realization mean, $\sigma_\beta/\sqrt{500} \approx 0.029^\circ/22.4 \approx 0.0013^\circ$). The $\sim 12\%$ multiplicative under-recovery is therefore a sky-fraction-independent property of the apodized pipeline (consistent with apodization-induced power suppression), not an artifact of the small canonical mask, and remains well below the published observational uncertainty $\sigma_\beta^{\text{obs}} = 0.094^\circ$ at every sky fraction tested. Artifact: `reproducibility/p1_namaster_500mc/results/c1_fsky_sweep.json`.

V. COSMOLOGICAL FITS AND MODEL COMPARISON

A. Datasets and Configuration

We analyze four dataset combinations: (1) Planck CMB (NPIPE (PR4) CamSpec high- ℓ TTTEEE + 2018 low- ℓ TT/EE + 2018 lensing) [18] (the NPIPE/PR4 release provides the high- ℓ CamSpec likelihood; the low- ℓ and lensing likelihoods are taken from the 2018 release, the standard pairing in the Cobaya likelihood stack); (2) +SDSS BAO — eBOSS DR16 LRG/QSO/Ly α (auto and $\times$ QSO) [19] + SDSS DR7 MGS [20] + 6dFGS [21]; (3) +Pantheon+ [14]; (4) +SH0ES M_B -anchor [7] + DES Y3 S_8 [11]. No DESI BAO likelihood enters the frozen Λ CDM+ ΔN_{eff} chains; DESI DR2 BAO enters only the separate iter2 $w_0 w_a$ chain of Table II. The exact Cobaya likelihood blocks of all five chains are listed in Table III. Parameter estimation uses Cobaya [22] (v3.5 original; v3.6.1 verification) with stock CAMB and ΔN_{eff} as a free parameter—no custom CAMB modifications. The extended parameter space adds ΔN_{eff} to Λ CDM, with both $(\omega/H)_0$ and Ω_k fixed to zero in the actual sampled YAML configuration (Sec. II). Reproducibility materials at <https://github.com/Hubify-Projects/bigbounce/tree/main/reproducibility>.

B. Results

a. Model-comparison statistics: deferred to a dedicated nested-sampling run. We do not report χ_{eff}^2 , AIC,

TABLE III. Per-chain Cobaya likelihood blocks, verbatim from the on-disk YAML configurations (`cobaya*.yaml`; frozen-chain `spin_torsion.input.yaml`; iter2 `spin_torsion.dr2.input.yaml`). All four Λ CDM+ ΔN_{eff} chains share the Planck block `planck_NPIPE_high1.CamSpec.TTTEE + planck.2018_low1.TT + planck.2018_low1.EE + planck.2018_lensing.clik`; the SDSS BAO block is `bao.sixdf.2011.bao + bao.sdss.dr7_mgs + bao.sdss.dr16.baoplus_{lrg,qso,lyauto,lyxqso}`. The first four rows are incremental: each adds to the stack of the row above. The iter2 chain uses `planck.2018_lensing.native` in place of `.clik`.

Chain	Likelihood stack	Status
Planck-only	Planck block	accumulating (not frozen)
Planck+BAO	+ SDSS BAO block	exploratory
Planck+BAO+SN	+ <code>sn.pantheonplus</code>	frozen (Table I)
Full-tension	+ <code>H0.riess2020Mb</code> + DES-Y3 S_8 Gaussian	frozen (Table I)
iter2 w_0w_a	<code>bao.desi.dr2.desi.bao.all</code> + Planck block + <code>sn.desy5</code> + <code>sn.pantheonplus</code>	converged (Table II)

BIC, or $\ln B$ Bayes-factor model-comparison numbers in this paper. Robust evaluation of these statistics on the present quintom posterior requires either a dedicated nested-sampling run (PolyChord or MultiNest) on the identical likelihood stack, or thermodynamic integration on a joint quintom/ Λ CDM run; a Savage-Dickey density-ratio readout from the converged Metropolis-Hastings chain used here is not viable because the Λ CDM point $(w_0, w_a) = (-1, 0)$ sits at $> 4\sigma$ in the joint marginal tails and is unsampled by the chain, so any KDE-based estimator yields kernel-dependent noise rather than a controlled density. The nested-sampling recompute is omitted; we report only the parameter posteriors below, which are unaffected by the model-comparison-statistic deferral. The two posterior parameter measurements that remain are $\Delta N_{\text{eff}} = -0.020 \pm 0.169$ (full-tension) and $+0.065 \pm 0.17$ (Planck+BAO+SN), both consistent with zero; these are the load-bearing numbers used elsewhere in the paper and they are unaffected by the model-comparison-statistic deferral. The χ^2 goodness-of-fit decomposition (BAO, CMB, SN, and total contributions) is reported in Table II; the AIC, BIC, and $\ln B$ evidence metrics are *not* reported there. The headline result is $w_0 = -0.812 \pm 0.044$ (departing from the Λ CDM point $w_0 = -1$ at $+4.3\sigma$; fn. a) and $w_a = -0.667 \pm 0.186$ (departing from $w_a = 0$ at -3.6σ), with $w_0 + w_a = -1.48 \pm 0.15$ requiring phantom crossing (the canonical quintom signature). Robust $\ln B$ computation requires nested sampling and is left to a follow-up analysis.

The Λ CDM+ ΔN_{eff} proxy thus offers neither posterior preference nor exclusion at the present data precision. The ΔN_{eff} extension alone does not resolve the Hubble tension.

VI. COSMIC BIREFRINGENCE: SPECTATOR ALP CONSISTENCY CHECK

Note.—This subsection presents a secondary consistency check using a spectator ALP model. The ALP produces cosmic birefringence independently of the gravitational theory; the same prediction $\beta \approx 0.27^\circ$ arises in standard GR with an identical ALP Lagrangian and natural parameters (taken at scan-prior midpoint values;

the $\sim 25\times$ misalignment tuning required for the headline result is disclosed below and in fn. 5). It is not derived from minimal ECH (which does not produce the required photon-torsion coupling) and is not a distinctive ECH prediction. The model class was previously studied by Fujita *et al.* [23].

Headline observational constraint.—The primary observational reference adopted in this analysis is the published Eskilt & Komatsu joint WMAP+Planck value $\beta = 0.342^\circ \pm 0.094^\circ$ (3.6σ) [2] (the joint WMAP9 + Planck PR4/NPIPE analysis; ACT DR6 enters only via the separate $\beta = 0.215^\circ \pm 0.074^\circ$ measurement [3], used below only in the auxiliary inverse-variance combination), which accounts for shared calibration systematics. The simplified inverse-variance combination below (3.9σ) is retained as an auxiliary cross-check only and is explicitly *not* used as the headline number anywhere in this paper.

ALP field evolution.—Numerical integration of the ALP equation of motion $\ddot{\phi} + 3H\dot{\phi} + m^2 f_a \sin(\phi/f_a) = 0$ in a Λ CDM background⁴ yields the field displacement from recombination to today:

$$\Delta\phi/f_a \approx 0.65 \quad (m = H_0, \theta_i = 1). \quad (2)$$

Across the natural parameter range $m/H_0 \in [1, 3]$, $\theta_i \in [0.5, 2]$: $\Delta\phi/f_a \in [0.2, 1.1]$.⁵

⁴ The ALP ODE is integrated on a Λ CDM late-time $H(z)$ here as an empirical background, distinct from the quintom-bounce dynamics that supply the early-universe / contracting-phase $H(z)$ in the underlying ECH cosmology [24]. The Λ CDM-background choice is conservative for the ALP-MCMC: a quintom late-time w_0w_a background (as in Sec. V) shifts $H(z)$ at $z \lesssim 1$ by $\sim$ few percent, propagating to a $\lesssim$ few-percent systematic on $\Delta\phi/f_a$ — well below the $\sim 30\%$ prior-width envelope on θ_i and m/H_0 dominating the β prediction.

⁵ Backreaction disclosure: the ALP backreaction fraction scales as $\Omega_a \sim \rho_a/\rho_{\text{crit}} \sim (m^2 f_a^2/H_0^2 M_{\text{Pl}}^2) \theta_i^2$, so the abstract’s spectator-status restriction $\theta_i \ll 1$ (see the *Spectator-status caveat* in the abstract) implies $\Omega_a \ll 1$ only in the sub-natural sliver $\theta_i \sim 0.1$. The numerical scan range $\theta_i \in [0.5, 2]$ is RETAINED here for completeness of the parameter envelope, but the natural-prior-anchored spectator-consistent result sits at $\theta_i \sim 0.1$: at $\theta_i = 0.1$ vs the scan-midpoint $\theta_i = 0.5$ the backreaction is

Birefringence value.—For $C_{a\gamma} = 8$, $\theta_i = 1$, $m \approx 2H_0$:

$$\beta \approx \frac{\alpha_{\text{EM}} \times 8}{4\pi} \times 1.07 = 4.97 \times 10^{-3} \text{ rad} \times \frac{180^\circ}{\pi} \approx 0.29^\circ. \quad (3)$$

The fiducial value $\beta \approx 0.27^\circ$ corresponds to the midpoint $m \approx 1.8H_0$, $\Delta\phi/f_a \approx 1.0$. The prediction spans $\beta \approx 0.17\text{--}0.43^\circ$ over $C_{a\gamma} \in [4, 12]$, $m/H_0 \in [1, 3]$, $\theta_i \in [0.5, 2]$, comfortably bracketing the observed value across this photon-coupling / mass / misalignment envelope (see Appendix C for the full ALP-MCMC sampled-parameter list, priors, and dataset-likelihood configuration). The spectator-consistent corner of this envelope ($\theta_i \sim 0.1$, per fn. 5) does require a $\sim 25\times$ tuning of the misalignment initial condition relative to the natural prior midpoint $\theta_i \sim 0.5$, so the model *accommodates* the observed signal but is not free of misalignment-prior tuning. The range $[0.17, 0.43]^\circ$ is obtained from a joint-trajectory scan over the *coupled* ($C_{a\gamma}, m/H_0, \theta_i$) space and not from an independent-extremes product (which would give the wider naive envelope $[0.027, 0.44]^\circ$); $\Delta\phi/f_a$ is a function of m/H_0 and θ_i along ALP trajectories rather than an independent variable.

Summary-likelihood combination (auxiliary cross-check).—Combining $\beta = 0.30^\circ \pm 0.11^\circ$ (Planck NPIPE [16]) and $\beta = 0.215^\circ \pm 0.074^\circ$ (ACT DR6 [3]) via inverse-variance weighting:

$$\beta_{\text{combined}} = 0.241^\circ \pm 0.061^\circ \quad (3.9\sigma) \quad (4)$$

(Auxiliary cross-check only; the 3.9σ is the significance of β_{combined} relative to the null $\beta = 0$, i.e. $0.241/0.061$.) This neglects shared calibration systematics, which produce positively correlated errors between the two measurements. Positively correlated errors *underestimate* the inverse-variance-combined σ , and therefore *overestimate* the significance: the naive 3.9σ figure is an upper bound on the true significance, not a lower bound. The published joint analysis at 3.6σ [2], which properly accounts for the shared polarization-angle calibration systematics via a joint covariance matrix with the Tau A self-calibration nuisance parameters, is the headline; the 3.9σ figure here is retained only as an internal cross-check demonstrating that the uncorrelated-errors approximation is qualitatively consistent with the published result.

MCMC parameter estimation.—Dedicated MCMC sampling of the ALP parameter space (3 configurations, 9,720 total accepted samples) yields: $\beta_{\text{ALP}} = 0.336^\circ \pm 0.107^\circ$ ($C_{a\gamma} = 8$ fixed; the direct-sample priors are on the underlying ALP parameters ($\theta_i, m/H_0$), not on $\Delta\phi/f_a$ which is a derived quantity along each ALP trajectory. The natural-envelope range

$\Delta\phi/f_a \in [0.2, 1.1]$ stated above corresponds to the corners of the ($\theta_i, m/H_0$) natural-prior box at $\theta_i \in [0.5, 2]$, $m/H_0 \in [1, 3]$. The MCMC posterior, anchored to the Gaussian summary likelihood on the published Eskiit–Komatsu joint WMAP+Planck measurement (fn. a; not to the *EB* spectra themselves), pulls ($\theta_i, m/H_0$) toward the upper edge of the natural-prior box — the data-preferred posterior implies the joint product $C_{a\gamma}(\Delta\phi/f_a) \approx 10.3$, which at the quoted $C_{a\gamma} = 8$ fixed corresponds to $\Delta\phi/f_a \approx 1.29$, $\sim 17\%$ above the natural envelope upper bound; this is the same $\sim 25\times$ -fine-tuning regime flagged in fn. 5 and is consistent with the model accommodating but not naturally explaining the observed $\beta_{\text{obs}} = 0.342^\circ$, consistent with the model-independent fit $\beta_{\text{free}} = 0.344^\circ \pm 0.096^\circ$ (our internal model-independent MCMC fit, with β as a free parameter, to a Gaussian summary likelihood on the published Eskiit–Komatsu joint WMAP+Planck birefringence measurement $\beta_{\text{obs}} = 0.342^\circ \pm 0.094^\circ$ [2] — not a re-analysis of the *EB* spectra themselves; 9,720 accepted samples across the 3 ALP-MCMC configurations described in Sec. VI (configurations $C_{a\gamma} = 4, 8, 12$; full priors and likelihood details in Appendix C); β_{free} denotes the unconstrained-amplitude fit distinct from β_{ALP} which has $C_{a\gamma} = 8$ fixed) and the observed $\beta_{\text{obs}} = 0.342^\circ \pm 0.094^\circ$. All three within 1σ . The combined coupling-displacement product entering the birefringence formula is $C_{a\gamma}(\Delta\phi/f_a) \approx 10.3$ ($\beta = 0.342^\circ$ in radians is 5.97×10^{-3} , the prefactor $\alpha_{\text{EM}}/(4\pi)$ is 5.8×10^{-4} , giving $C_{a\gamma} \Delta\phi/f_a = \beta/[\alpha_{\text{EM}}/(4\pi)] \approx 10.3$); with the field-displacement range $\Delta\phi/f_a \in [0.2, 1.1]$ from the numerical ALP-EOM integration, the required $C_{a\gamma}$ spans ~ 9 to ~ 51 . Both ends are larger than the standard KSVZ/DFSZ benchmark range, which predicts $|C_{a\gamma}| \sim \mathcal{O}(1)$; the entire required range therefore lies outside minimal ALP photon-coupling benchmarks and requires non-minimal model building. The lower end (~ 9) can be accommodated in extended models with modest photon-coupling enhancement (e.g. chiral-fermion-loop enhancement, clockwork constructions), while the upper end (~ 51) requires either substantial UV-completion enhancement or a field displacement at the upper end of the $\Delta\phi/f_a \in [0.2, 1.1]$ range. The signal is therefore accommodated across the considered parameter space rather than fine-tuned only at one benchmark, but the upper-coupling end is not generic. We emphasize that the spectator-consistent corner $\theta_i \sim 0.1$ (fn. 5) inflates the required photon-coupling further: at fixed $\beta = 0.342^\circ$, $\Delta\phi/f_a \propto \theta_i$ along the underdamped trajectory, so a $5\times$ reduction in θ_i from the scan midpoint demands a correspondingly higher $C_{a\gamma}$ to hold the $C_{a\gamma} \Delta\phi/f_a \approx 10.3$ product, pushing the required enhancement well above standard KSVZ/DFSZ $\mathcal{O}(1)$ benchmarks. Because the EOM-required coupling range $[\sim 9, \sim 51]$ exceeds the $C_{a\gamma} \in [1, 30]$ prior of the original extended configuration (which truncated $\sim 28\%$ of the posterior mass above $C_{a\gamma} = 30$), we reran the $C_{a\gamma}$ -free fit with a continuous uniform prior $C_{a\gamma} \in [4, 60]$ covering

$\Omega_a(0.1)/\Omega_a(0.5) \sim 1/25$ (i.e., a $\sim 25\times$ fine-tuning of the misalignment initial condition is required to keep the ALP a true spectator). The $\Omega_a \sim 1$ regime at $\theta_i \sim 1$ is the dark-energy-ALP regime explicitly excluded from this spectator-consistency companion check.

the full required band (4 MPI chains, $\hat{R} - 1 = 0.0095$, 8,955 accepted samples). The resulting posterior is broad, as expected for a single-amplitude constraint on a three-parameter degeneracy: median $C_{a\gamma} = 20.7$ with 16–84% range [7.3, 45.6], and 69% of the posterior mass falls inside the EOM-required band [9, 51] ($\approx 10.3/1.1$ to $10.3/0.2$, the $C_{a\gamma} \Delta\phi/f_a \approx 10.3$ product divided across the $\Delta\phi/f_a \in [0.2, 1.1]$ envelope; 22% of the mass lies below $C_{a\gamma} = 9$, consistent with the high- $\Delta\phi/f_a$ tail, where the fixed product is met at smaller coupling). These mass fractions are prior-dependent statements (flat $C_{a\gamma} \in [4, 60]$, $\theta_i \in [0.01, \pi]$, $\log_{10} m_a$ priors), not prior-independent measurements. The recovered $\beta = 0.326^\circ \pm 0.099^\circ$ posterior matches the observed $0.342^\circ \pm 0.094^\circ$, confirming the consistency-check verdict over the full required coupling range rather than at fixed benchmarks. *Spectator-subset readout* (same chain, no additional sampling): the chain’s derived ALP energy fraction Ω_a satisfies $\Omega_a < 0.1$ for 44% and $\Omega_a < 0.01$ for 13% of the posterior mass; restricted to the $\Omega_a \leq 0.01$ spectator-safe subset, the rotation marginal is $\beta = 0.28^\circ \pm 0.10^\circ$, consistent with $\beta_{\text{obs}} = 0.342^\circ \pm 0.094^\circ$ at 0.5σ . The strict $\theta_i \leq 0.1$ sliver of fn. 5 carries only 0.4% of the posterior mass (too few samples for a stable marginal), quantifying the misalignment tuning of the spectator-consistent corner. Convergence: $\hat{R} - 1 < 0.01$ for all runs. The joint posterior structure of this continuous-prior configuration is shown in Fig. 4. Artifact: `research/branch_R_alp_birefringence/phase2_mcmc/chains/c5_continuous/`.

LiteBIRD forecast.—LiteBIRD is projected to achieve $\sigma(\beta) \approx 0.03^\circ$ [25]. For $\beta = 0.27^\circ$: $\sim 9\sigma$ statistical significance—either decisive confirmation or clean exclusion.

Caveats.—This birefringence prediction is independent of bounce cosmology: the ALP is a spectator field that does not participate in the bounce dynamics. The ECH framework provides heuristic motivation ($f_a \sim M_{\text{Pl}}$ from the Holst sector pseudoscalar structure) but no derived photon-torsion coupling connects the Holst action to a specific ALP potential. The model *accommodates* the observed signal for natural parameter values (taken at scan-prior midpoint values; the $\sim 25\times$ misalignment tuning required for the headline result is disclosed in Sec. VI and fn. 5); it does not uniquely predict it.

VII. CONCLUSIONS

This companion paper documents three technical verification analyses for the ECH spin-torsion structural closure program [1].

$\Lambda\text{CDM} + \Delta N_{\text{eff}}$ MCMC proxy.—The stock-CAMB Cobaya v3.6.1 run (309,189 frozen samples across two converged dataset combinations; an additional 114,992-sample Planck-only run is still accumulating at $\hat{R} - 1 \sim 0.05$, is *not* reported in Table I, and is *not* aggregated into the frozen headline) finds ΔN_{eff} consistent with




FIG. 4. Spectator-ALP joint posterior triangle from the continuous-prior cross-check configuration in which the photon anomaly coefficient is sampled freely (flat priors $C_{a\gamma} \in [4, 60]$ — widened from the earlier [1, 30] to cover the full EOM-required band [9, 51] — $\theta_i \in [0.01, \pi]$, $\log_{10}(m_a/\text{eV}) \in [-35, -30]$ — the mass prior corresponds to $m/H_0 \approx 7 \times 10^{-3}$ to 7×10^2 for $H_0 = 67.7 \text{ km/s/Mpc} = 1.44 \times 10^{-33} \text{ eV}$, far broader than the benchmark $m/H_0 \in [1, 3]$), broadening the three fixed- $C_{a\gamma}$ benchmark configurations of Appendix C; the direct-sample priors are on the underlying ALP parameters (θ_i, m_a), with $\Delta\phi/f_a$ derived along each trajectory. The rotation marginal, $\beta = 0.326^\circ \pm 0.099^\circ$, is consistent within 1σ with the fixed- $C_{a\gamma} = 8$ result $\beta_{\text{ALP}} = 0.336^\circ \pm 0.107^\circ$ and the observed $\beta_{\text{obs}} = 0.342^\circ \pm 0.094^\circ$ (an internal-consistency statement: all three are constrained by the same single published β_{obs} , not independent measurements). The θ_i – $C_{a\gamma}$ anti-correlation band traces the constant-product degeneracy enforced by the summary-likelihood constraint through $C_{a\gamma} (\Delta\phi/f_a) \approx 10.3$, and the m_a marginal piles toward the upper (heavier) edge of its prior range — the data-driven pull toward the upper edge of the natural-prior box discussed in the text, consistent with the model *accommodating* rather than naturally predicting the observed rotation (fn. 5).

zero in both frozen dataset combinations and recovers $H_0 = 67.68 \pm 1.06 \text{ km s}^{-1} \text{ Mpc}^{-1}$, consistent with standard Planck ΛCDM . The ΔN_{eff} extension does not resolve the Hubble tension. Current data neither require nor exclude a spin-torsion ΔN_{eff} contribution; CMB-S4 will provide the first precision test ($\sigma(N_{\text{eff}}) \sim 0.03$). Bayes factors and information criteria (ΔAIC , ΔBIC , $\ln B$) are deferred to a dedicated nested-sampling run (Sec. V, *Model-comparison statistics* paragraph); we report only the parameter posteriors (ΔN_{eff} , H_0 , etc.) as load-bearing here and explicitly omit Bayes-factor or information-criterion comparisons in this Metropolis-Hastings-only analysis.

NaMaster pipeline validation.—The 500-MC pseudo- C_ℓ analysis on synthetic Λ CDM polarization skies confirms that the deconvolution pipeline recovers injected birefringence angles with amplitude-dependent bias $\hat{\beta} - \beta_{\text{inj}} = -0.032$ to -0.040° (worst-case -0.040° at injection $\beta = 0.342^\circ$; Sec. IV) at template-fit SNR consistent with the ACT-noise floor. This is a methods validation, not a competitive sky detection; the primary observational evidence for cosmic birefringence remains the published Planck/ACT DR6 2.7–2.9 σ measurements.

Spectator-ALP consistency.—An ALP with $f_a \sim M_{\text{Pl}}$, $m \sim H_0$ is consistent with the published 3.6 σ joint signal, with the caveat that the spectator-consistent regime $\theta_i \sim 0.1$ (fn. 5) requires $\sim 25\times$ misalignment tuning relative to the natural prior midpoint ($C_{a\gamma} \Delta\phi/f_a \approx 10.3$ for $\beta = 0.342^\circ$, with $\Delta\phi/f_a$ in the natural range $[0.2, 1.1]$ giving $C_{a\gamma}$ between ~ 9 and ~ 51 ; see §VI for the explicit numerical derivation). The same result arises in standard GR; the ECH framework provides motivation but not a derivation. LiteBIRD will settle this at $\sim 9\sigma$ in the early 2030s.

Quintom-B empirical anchor.—The converged DESI DR2 + Planck NPIPE + Pantheon+ [14] + DES-SN5YR [15] Cobaya chain with the w_0w_a free-parameter extension (128,385 accepted samples across 16 MPI chains; $\hat{R} - 1 = 0.00820$, below the standard $\hat{R} - 1 < 10^{-2}$ publication target) supplies the GetDist w_0w_a posteriors of Table II, an empirical test of the quintom-B scenario [13].

Data and Code Availability

All materials are at:

<https://github.com/Hubify-Projects/bigbounce/tree/main/reproducibility>

The repository is program-wide: it includes the four Cobaya YAML configurations (one per dataset combination, stock CAMB, no custom modifications), NaMaster driver scripts, and the implementation map (IMPLEMENTATION_MAP.md), alongside galaxy-spin pipeline code used by Paper IV [6] but not by any analysis of this paper. MCMC chains: regenerate via `reproduce_cosmology.sh` (~ 4 – 12 h per configuration on 4 CPU cores).

ACKNOWLEDGMENTS

We thank the Planck, ACT, LiteBIRD, and Cobaya collaborations for providing the data, code, and observational infrastructure used in these analyses. The author acknowledges the use of Claude (Anthropic) as an AI research assistant during systematic analysis and manuscript preparation. All scientific claims, derivations, numerical results, and bibliographic attributions were independently verified by the author. No external

funding was received. Computational resources were self-funded (RunPod H200 instances).

Appendix A: Reproducibility Materials

Repository structure.—The public repository <https://github.com/Hubify-Projects/bigbounce> contains:

- Four Cobaya YAML configurations (`cobaya_planck.yaml`, `cobaya_planck_bao.yaml`, `cobaya_planck_bao_sn.yaml`, `cobaya_full_tension.yaml`)—stock CAMB, no torsion modifications.
- `galaxy_spins/spin_fit_stan.py`—hierarchical Bayesian model (CmdStanPy) fitting $A(z)$ to published aggregate CW/CCW galaxy counts (program-wide content used by Paper IV, not by this paper’s analyses).
- `data.build/build_galaxy_spin_dataset.py`—reproducible pipeline downloading Galaxy Zoo DECaLS [26] from Zenodo (DOI: 10.5281/zenodo.4573248, CC-BY-4.0; likewise Paper IV content).
- `docs/IMPLEMENTATION_MAP.md`—mapping from each result to its code artifact.
- `docs/KNOWN_GAPS.md`—honest disclosure of what cannot currently be reproduced.

What is NOT included.—MCMC chains are not pre-computed (regenerate via `reproduce_cosmology.sh`, ~ 4 – 12 h per config on 4 CPU cores). Bayes factors and information criteria (ΔAIC , ΔBIC , $\ln B$) are NOT reported in this manuscript: dedicated nested sampling via PolyChord or MultiNest on the identical likelihood stack is required for a controlled Bayesian evidence on the present quintom posterior (Sec. V, *Model-comparison statistics* paragraph). A Savage-Dickey readout from the present Metropolis-Hastings chain is not viable—the LCDM point $(w, w_a) = (-1, 0)$ lies at $> 4\sigma$ in the joint marginal tails and is unsampled, so a KDE-based estimator yields kernel-dependent noise rather than a controlled density. This is left to a dedicated future nested-sampling analysis. No CNN galaxy classifier is included; the hierarchical fit uses published catalog labels. No CMB polarization map analysis code is provided beyond the NaMaster driver script; all published birefringence values are literature citations.

HuggingFace datasets.—The following HuggingFace datasets accompany this work (links in the repository README):

1. MCMC chain diagnostics and convergence CSV files.
2. NaMaster pipeline artifacts (mask, MC seeds, output spectra).
3. ALP parameter MCMC chains.

Appendix B: Claims Classification

Appendix C: ALP-MCMC Sampled Parameters, Priors, and Likelihood Stack

The ALP-MCMC results quoted in Sec. VI ($\beta_{\text{ALP}} = 0.336^\circ \pm 0.107^\circ$ at $C_{a\gamma} = 8$ fixed; $\beta_{\text{free}} = 0.344^\circ \pm 0.096^\circ$ model-independent; 9,720 total accepted samples across 3 configurations) use the following setup.

Sampled parameters and priors (model-dependent ALP fit).

- $C_{a\gamma}$: fixed at one of $\{4, 8, 12\}$ across the three configurations (3,240 samples per configuration); not a sampled parameter within a single chain.
- m/H_0 : uniform prior on $[1, 3]$ (ultra-light ALP class; $m \sim H_0$ required for cosmological-time field evolution).
- θ_i : uniform prior on $[0.5, 2]$ (natural-misalignment range, $\theta_i = \phi_{\text{init}}/f_a$).⁶
- f_a : fixed at M_{Pl} (spectator-class theoretical input from the Holst-sector pseudoscalar structure; not sampled).

Sampled parameters and priors (model-independent β_{free} fit).

- β : uniform prior on $[-2^\circ, 2^\circ]$; sampled as a free amplitude with no ALP-model parametric structure.

Likelihood stack.—All fits use a Gaussian summary likelihood on the published Eskilt–Komatsu

joint WMAP+Planck isotropic-birefringence measurement $\beta_{\text{obs}} = 0.342^\circ \pm 0.094^\circ$ [2] (fn. a; the chain configuration encodes exactly `beta_obs: 0.342, sigma.beta: 0.094`) — i.e. the constraint enters at the level of the published β posterior, not as a direct re-analysis of the EB spectra. The separate ACT DR6 measurement $\beta = 0.215^\circ \pm 0.074^\circ$ [3] is an independent cross-check quoted in Sec. VI and does *not* enter any MCMC likelihood. The MCMC engine is Cobaya v3.6.1 with the Metropolis-Hastings sampler; convergence threshold $\hat{R} - 1 < 0.01$ across all 3 benchmark configurations (achieved at $N_{\text{tot}} = 9,720$ accepted samples post burn-in) and for the continuous-prior $C_{a\gamma} \in [4, 60]$ configuration ($\hat{R} - 1 = 0.0095$, 8,955 accepted samples). *Chain-to-number map*: $\beta_{\text{ALP}} = 0.336^\circ \pm 0.107^\circ$ — the three fixed- $C_{a\gamma} \in \{4, 8, 12\}$ benchmark chains (9,720 samples); $\beta_{\text{free}} = 0.344^\circ \pm 0.096^\circ$ — the model-independent single-parameter chain; median $C_{a\gamma} = 20.7$, 16–84% range $[7.3, 45.6]$, $\beta = 0.326^\circ \pm 0.099^\circ$, and the spectator-subset readouts — the continuous-prior c5 chain (8,955 samples).

Scope statement.—These fits constrain a specific ultra-light ALP class ($m \sim H_0$, $f_a \sim M_{\text{Pl}}$; the $C_{a\gamma} \in [4, 12]$ benchmark sweep plus the continuous-prior $C_{a\gamma} \in [4, 60]$ headline configuration). The consistency with β_{obs} established here is *not* a universal mechanism-independent statement; alternative parity-violating mechanisms (Chern-Simons, axion dark matter at higher m , etc.) require independent fits with their own model parameters.

-
- [1] H. Golden, Structural Closure of Einstein–Cartan–Holst Dark Energy: Perturbation Transparency, Inflation– f_{NL} Tension, and Surviving Matter-Bounce Tests, (in preparation) (2026), hUBIFY-2026-001A; companion paper, this volume.
 - [2] J. R. Eskilt and E. Komatsu, Improved constraints on cosmic birefringence from the WMAP and Planck cosmic microwave background polarization data, *Phys. Rev. D* **106**, 063503 (2022), arXiv:2205.13962 [astro-ph.CO].
 - [3] P. Diego-Palazuelos and E. Komatsu, Cosmic birefringence from the Atacama Cosmology Telescope data release 6, arXiv preprint (2025), arXiv:2509.13654 [astro-ph.CO].
 - [4] H. Golden, $f_{\text{NL}} = -35/8$ Forecast: SPHEREx Discrimination of Bounce vs. Inflation, (in preparation) (2026), hUBIFY-2026-002; companion paper, this volume.
 - [5] H. Golden, Spectrally Unusual Sources at Scale: A Multi-

Survey Catalog of 378,280 Anomalies and Native-Trained Novelty Rates from 37.3 Million Sources, (in preparation) (2026), hUBIFY-2026-003; companion paper, this volume.

- [6] H. Golden, Galaxy Chirality at Scale: 8.47M Galaxies Classified, Hemisphere Null at $p_{\text{LEE}} < 10^{-4}$, (in preparation) (2026), hUBIFY-2026-004; companion paper, this volume.
- [7] A. G. Riess, W. Yuan, L. M. Macri, *et al.*, A comprehensive measurement of the local value of the Hubble constant with 1 km/s/Mpc uncertainty from the Hubble Space Telescope and the SH0ES team, *The Astrophysical Journal Letters* **934**, L7 (2022), arXiv:2112.04510 [astro-ph.CO].
- [8] F. W. Hehl, P. von der Heyde, G. D. Kerlick, and J. M. Nester, General relativity with spin and torsion: Foundations and prospects, *Reviews of Modern Physics* **48**, 393 (1976).
- [9] S. Mercuri, Fermions in the Ashtekar-Barbero connection formalism for arbitrary values of the Immirzi parameter, *Physical Review D* **73**, 084016 (2006), arXiv:gr-qc/0601013 [gr-qc].
- [10] Y.-F. Cai, W. Xue, R. Brandenberger, and X. Zhang, Non-gaussianity in a matter bounce, *JCAP* **0905**, 011, arXiv:0903.0631.

⁶ Backreaction / spectator-status disclosure: see fn. 5 in Sec. VI. In brief, the $\theta_i \in [0.5, 2]$ prior is an envelope-completeness choice, not the spectator-consistent sub-range ($\theta_i \sim 0.1$, a $\sim 25\times$ tuning relative to the prior midpoint); posterior samples at $\theta_i \gtrsim 0.5$ belong to the dark-energy-ALP regime excluded from the spectator-consistency claim.

TABLE IV. Claims classification for this companion paper.

Claim	Type	Status	Notes
$\Delta N_{\text{eff}} = -0.020 \pm 0.169$ (full-tension)	MCMC	Verified	Stock CAMB proxy
$\Delta N_{\text{eff}} = +0.065 \pm 0.17$ (Planck+BAO+SN)	MCMC	Verified	Stock CAMB proxy
$H_0 = 67.68 \pm 1.06$ (full-tension)	MCMC	Verified	Recovers Λ CDM
$H_0 = 67.79 \pm 1.09$ (Planck+BAO+SN)	MCMC	Verified	Recovers Λ CDM
Model-comparison $\Delta\text{AIC}/\text{BIC}/\ln B$	Numerical	Omitted	Follow-up nested-sampling analysis
$\hat{\beta}_{\text{NaMaster}} = 0.238^\circ$ (500-MC)	Numerical	Verified	Pipeline; MC bias table
$\beta_{\text{ALP}} = 0.336^\circ \pm 0.107^\circ$	MCMC	Verified	ALP MCMC
Published 3.6σ ($\beta = 0.342 \pm 0.094^\circ$)	Lit.	Cited	Eskilt et al.
Stock CAMB proxy $\neq$ ECH theory module	Scope	Defn.	§III
ALP birefringence not distinctive ECH prediction	Scope	Defn.	§VI

- [11] DES Collaboration, T. M. C. Abbott, *et al.*, Dark Energy Survey Year 3 results: Cosmological constraints from galaxy clustering and weak lensing, *Physical Review D* **105**, 023520 (2022), [arXiv:2105.13549 \[astro-ph.CO\]](#).
- [12] T. Liu, X. Li, T. Xu, M. Biesiada, and J. Wang, Torsion cosmology in the light of DESI, supernovae and CMB observational constraints, *European Physical Journal C* (2025), [arXiv:2507.04265 \[gr-qc\]](#).
- [13] DESI Collaboration, M. Abdul-Karim, *et al.*, DESI DR2 results II: Measurements of baryon acoustic oscillations and cosmological constraints, *Physical Review D* **112**, 083515 (2025), [arXiv:2503.14738 \[astro-ph.CO\]](#).
- [14] D. Brout *et al.*, The Pantheon+ analysis: Cosmological constraints, *Astrophys. J.* **938**, 110 (2022), [arXiv:2202.04077 \[astro-ph.CO\]](#).
- [15] DES Collaboration, T. M. C. Abbott, *et al.*, The dark energy survey: Cosmology results with ~ 1500 new high-redshift type Ia supernovae using the full 5-yr data set, *Astrophys. J. Lett.* **973**, L14 (2024), [arXiv:2401.02929 \[astro-ph.CO\]](#).
- [16] P. Diego-Palazuelos, J. R. Eskilt, Y. Minami, M. Tristram, *et al.*, Cosmic birefringence from the Planck data release 4, *Phys. Rev. Lett.* **128**, 091302 (2022), reports $\beta = 0.30 \pm 0.11$ deg from Planck NPIPE (PR4), [arXiv:2201.07682 \[astro-ph.CO\]](#).
- [17] D. Alonso, J. Sanchez, and A. Slosar (LSST Dark Energy Science), A unified pseudo- C_ℓ framework, *Mon. Not. Roy. Astron. Soc.* **484**, 4127 (2019), [arXiv:1809.09603 \[astro-ph.CO\]](#).
- [18] Planck Collaboration, N. Aghanim, *et al.*, Planck 2018 results. VI. cosmological parameters, *Astronomy & Astrophysics* **641**, A6 (2020), [arXiv:1807.06209 \[astro-ph.CO\]](#).
- [19] S. Alam *et al.*, Completed SDSS-IV extended baryon oscillation spectroscopic survey: Cosmological implications from two decades of spectroscopic surveys at the Apache Point Observatory, *Phys. Rev. D* **103**, 083533 (2021), [arXiv:2007.08991 \[astro-ph.CO\]](#).
- [20] A. J. Ross, L. Samushia, C. Howlett, W. J. Percival, A. Burden, and M. Manera, The clustering of the SDSS DR7 main galaxy sample – I. a 4 per cent distance measure at $z = 0.15$, *Mon. Not. R. Astron. Soc.* **449**, 835 (2015), [arXiv:1409.3242 \[astro-ph.CO\]](#).
- [21] F. Beutler, C. Blake, M. Colless, D. H. Jones, L. Staveley-Smith, L. Campbell, Q. Parker, W. Saunders, and F. Watson, The 6dF Galaxy Survey: baryon acoustic oscillations and the local Hubble constant, *Mon. Not. R. Astron. Soc.* **416**, 3017 (2011), [arXiv:1106.3366 \[astro-ph.CO\]](#).
- [22] J. Torrado and A. Lewis, Cobaya: Code for Bayesian analysis of hierarchical physical models, *Journal of Cosmology and Astroparticle Physics* **05** (057), 057, [arXiv:2005.05290 \[astro-ph.IM\]](#).
- [23] T. Fujita, K. Murai, H. Nakatsuka, and S. Tsujikawa, Detection of isotropic cosmic birefringence and its implications for axionlike particles including dark energy, *Physical Review D* **103**, 043509 (2021), [arXiv:2011.11894 \[astro-ph.CO\]](#).
- [24] Y.-F. Cai, E. N. Saridakis, M. R. Setare, and J.-Q. Xia, Quintom Cosmology: Theoretical implications and observations, *Phys. Rept.* **493**, 1 (2010), canonical quintom-cosmology review (two-field DE with w crossing -1). Used in P1A Sec. VI to point readers to the bounce-class alternative DE mechanism that survives the 14 ECH-specific structural barriers., [arXiv:0909.2776 \[hep-th\]](#).
- [25] LiteBIRD Collaboration, E. Allys, *et al.*, Probing cosmic inflation with the LiteBIRD cosmic microwave background polarization survey, *Progress of Theoretical and Experimental Physics* **2023**, 042F01 (2023), [arXiv:2202.02773 \[astro-ph.IM\]](#).
- [26] M. Walmsley, C. Lintott, T. G eron, *et al.*, Galaxy Zoo DECaLS: Detailed visual morphology measurements from volunteers and deep learning for 314 000 galaxies, *Monthly Notices of the Royal Astronomical Society* **509**, 3966 (2022), [arXiv:2102.08414 \[astro-ph.GA\]](#).