

# Technical Verification Companion: $\Lambda$ CDM+ $\Delta N_{\text{eff}}$ MCMC Proxy, NaMaster Pipeline Recovery, and Spectator-ALP Consistency Check for the ECH Spin-Torsion Program

Houston Golden<sup>1,\*</sup>

<sup>1</sup>*Independent Researcher, Los Angeles, California, USA*

(Dated: May 15, 2026 PDT — v1B.0.9)

We report the technical verification material for the Einstein-Cartan-Holst (ECH) spin-torsion cosmology no-go program of Paper I(a) [1]. Three analyses are documented. (1) *Stock-CAMB  $\Lambda$ CDM+ $\Delta N_{\text{eff}}$  MCMC proxy* (Cobaya v3.6.1, **309,789** frozen samples across two converged dataset combinations, plus a third Planck-only combination ongoing): this run uses stock CAMB with  $\Delta N_{\text{eff}}$  as a free parameter and carries *no torsion modifications to the Boltzmann equations*; it is reported as a null-consistency test of an extra radiation-like degree of freedom, not as evidence for or against the ECH spin-torsion framework. Both frozen dataset combinations find  $\Delta N_{\text{eff}}$  consistent with zero ( $-0.020 \pm 0.169$  full-tension;  $+0.065 \pm 0.17$  Planck+BAO+SN) and  $H_0$  consistent with standard  $\Lambda$ CDM ( $67.68 \pm 1.06$  full-tension;  $67.79 \pm 1.09$  Planck+BAO+SN, both in  $\text{km s}^{-1} \text{Mpc}^{-1}$ ). (2) *NaMaster pseudo- $C_\ell$  pipeline validation* on the Planck Commander CMB polarization map ( $N_{\text{side}} = 512$ ,  $\ell_{\text{max}} = 1024$ ,  $f_{\text{sky}} = 0.32$ , 500 Monte Carlo realizations): injecting the spectator-ALP fiducial value  $\beta = 0.27^\circ$  recovers  $\hat{\beta} = 0.238^\circ$  (pipeline-recovery bias  $0.032^\circ$ , SNR= 20.32). The primary sky detection significance is the published Planck/ACT DR6 2.4–2.9 $\sigma$  [2, 3]; the pipeline SNR figures refer to recovery of injected MC signals and are *not* competitive sky measurements. (3) *Spectator-ALP consistency check*: a field with  $f_a \sim M_{\text{Pl}}$ ,  $m \sim H_0$  is consistent with the published joint Planck+ACT value  $\beta = 0.342^\circ \pm 0.094^\circ$  (3.6 $\sigma$ ) [4] without fine-tuning. The same birefringence arises in standard GR with an identical ALP; it is not a distinctive ECH prediction. A cross-paper status table and reproducibility manifest are included (Sec. VII).

PACS numbers: 98.80.-k, 95.36.+x, 04.50.Kd

## CONTENTS

|                                                                                                                                 |   |
|---------------------------------------------------------------------------------------------------------------------------------|---|
| I. Introduction                                                                                                                 | 1 |
| II. Cosmological Tensions: $H_0$ and $\sigma_8$                                                                                 | 2 |
| III. Stock-CAMB $\Lambda$ CDM+ $\Delta N_{\text{eff}}$ MCMC: Generic Radiation-Proxy Test<br>(Not a Spin-Torsion Theory Module) | 2 |
| IV. Data Methods: CMB $E$ - $B$ Analysis                                                                                        | 3 |
| V. Cosmological Fits and Model Comparison                                                                                       | 4 |
| A. Datasets and Configuration                                                                                                   | 4 |
| B. Results                                                                                                                      | 4 |
| VI. Cosmic Birefringence: Spectator ALP Consistency Check                                                                       | 4 |
| VII. Cross-Paper Verification Status                                                                                            | 5 |
| A. Free- $w_0 w_a$ chain status (anchor for Paper I(a) Table II ‡ footnote)                                                     | 5 |
| VIII. Conclusions                                                                                                               | 5 |
| Data and Code Availability                                                                                                      | 6 |
| Acknowledgments                                                                                                                 | 7 |

## A. Reproducibility Materials

7

## B. Claims Classification

7

## References

7

## I. INTRODUCTION

This companion paper provides the technical verification layer for the ECH structural-closure no-go result reported in Paper I(a) [1]. The main paper establishes 14 independent structural constraints closing minimal-ECH dark-energy routes and proves a perturbation-transparency theorem for the Holst sector. The present paper documents the three numerical analyses that support and contextualize those results.

*Scope of this paper.*—Three verification analyses are reported here, each with explicit scope limitations:

1.  $\Lambda$ CDM+ $\Delta N_{\text{eff}}$  MCMC proxy (Secs. II and III): stock CAMB Boltzmann solver with  $\Delta N_{\text{eff}}$  as a free parameter. **Not a spin-torsion theory module.** The Boltzmann code carries no torsion modifications; this run tests whether the data prefer an extra radiation-like degree of freedom in standard cosmology.
2. NaMaster CMB  $E$ - $B$  analysis (Sec. IV): a bias-injection Monte Carlo validation of the pseudo- $C_\ell$  deconvolution pipeline. **Not a competitive sky detection.** The high pipeline-recovery SNR figures (e.g.,

\* houston@hubify.com

20.32 $\sigma$ ) refer to recovery of injected MC signals, not to the significance of the CMB sky measurement, which remains the published 2.4–2.9 $\sigma$  from Planck/ACT.

3. *Spectator-ALP consistency check* (Sec. VI): a standard GR+ALP computation showing that the observed signal is consistent with an ALP having natural parameters. **Not a distinctive ECH prediction.** The same  $\beta \approx 0.27^\circ$  arises in any GR+ALP setup with the same parameters; no ECH-specific derivation connects the Holst action to the photon-torsion coupling required.

*What is NOT in this paper.*—The 13 logically-independent structural barriers (14 historical catalog entries; see Paper I(a) v1A.0.22), the perturbation-transparency theorem, the 14-barrier table, and the surviving mechanism-independent test predictions ( $f_{\text{NL}} = -35/8$  and ALP birefringence) are in Paper I(a) [1]. The SPHEREx multi-tracer Fisher forecast is in Paper II [5]. The multi-survey anomaly catalog is in Paper III [6]. The galaxy chirality catalog is in Paper IV [7].

*Cross-paper citations.*—When this companion reports MCMC values ( $H_0$ ,  $\sigma_8$ , etc.) that are referenced in the main paper, those values come from Secs. III and V here.

## II. COSMOLOGICAL TENSIONS: $H_0$ AND $\sigma_8$

The bounce scenario motivates extending  $\Lambda$ CDM by  $\Delta N_{\text{eff}}$  (particle production at the bounce) as a phenomenological proxy parameter;  $(\omega/H)_0$  (angular momentum transfer) and  $\Omega_k$  are fixed to zero in the actual sampled MCMC configuration of this paper (per the explicit parameter-scope clarification in Sec. V A; the  $(\omega/H)_0$  parameter is discussed in Paper I(a) as a phenomenological bounce-class indicator but is not separately sampled here). The full-tension dataset combination includes the SH0ES [8]  $H_0$  prior in the Cobaya likelihood configuration, but the high-precision Planck NPIPE CamSpec TTTEEE+lowl+lensing likelihoods carry sufficient inverse-variance weight that the posterior  $H_0$  in the proxy run is pulled to  $67.68 \pm 1.06$  (Planck-dominated value) rather than to the simple Gaussian-combination value  $\sim 70$  that would emerge if SH0ES and Planck were equally weighted. We do not therefore claim that the SH0ES tension is resolved or even moved by adding  $\Delta N_{\text{eff}}$  in stock CAMB; in this configuration the  $\Delta N_{\text{eff}}$  posterior is consistent with zero and the  $H_0$  posterior is dominated by Planck. The spin-torsion framework alone does not resolve cosmological tensions at the present data precision.

## III. STOCK-CAMB $\Lambda$ CDM+ $\Delta N_{\text{eff}}$ MCMC: GENERIC RADIATION-PROXY TEST (NOT A SPIN-TORSION THEORY MODULE)

*Scope statement.*—“MCMC verification” refers throughout to a stock CAMB run of  $\Lambda$ CDM extended by

$\Delta N_{\text{eff}}$  as a free parameter. *No custom CAMB modifications are used; no torsion-modified Boltzmann equations are solved.* The MCMC therefore tests whether the data prefer an extra radiation-like degree of freedom, treated as a generic phenomenological proxy for the spin-torsion sector’s possible effective radiation contribution. It does *not* verify the spin-torsion theory module itself; that would require a bespoke modified Boltzmann code.

The proxy run (Cobaya v3.6.1 with Planck NPIPE CamSpec TTTEEE + lowl TT/EE + lensing) has produced two frozen dataset combinations with publication-quality convergence, plus a third Planck-only run currently at sub-convergence sample count and not aggregated into any frozen-posterior summary statistic in this paper. Frozen MCMC program: **309,789** raw samples across 2 frozen dataset combinations (176,840 + 132,949). We report this as a  $\Lambda$ CDM+ $\Delta N_{\text{eff}}$  null-consistency test: the data are consistent with  $\Delta N_{\text{eff}} = 0$  in stock CAMB.<sup>1</sup>

a. *Scope of the  $\Delta N_{\text{eff}}$  proxy: bounce-class discrimination, not a direct test of the spin-torsion sector.* We emphasize that the stock-CAMB  $\Lambda$ CDM+ $\Delta N_{\text{eff}}$  proxy run does *not* test the ECH spin-torsion sector directly. The Hehl-Datta-Mercuri parity-even four-fermion contact interaction that survives torsion elimination [9, 10] is dimension-6 and  $M_{\text{Pl}}^{-2}$ -suppressed: its leading Boltzmann effect is a scattering-amplitude shift, not a relativistic species, and it does *not* produce a  $\Delta N_{\text{eff}}$  at recombination. The matter-bounce class [11] predicts  $\Delta N_{\text{eff}} \approx 0$  by construction (no light bounce-internal species are thermalized at recombination); the proxy run *confirming*  $\Delta N_{\text{eff}} = -0.020 \pm 0.169$  (full-tension) and  $+0.065 \pm 0.17$  (Planck+BAO+SN) is therefore consistent with the matter-bounce prediction, but the absence of a  $\Delta N_{\text{eff}}$  detection is not a discriminator between minimal-ECH and standard  $\Lambda$ CDM at the present data precision. We frame the proxy as a bounce-class *compatibility check* (matter-bounce-class scenarios without prolonged post-bounce inflation predict  $\Delta N_{\text{eff}} \approx 0$ ), not as a posterior-preference test against a competing model.

<sup>1</sup> Sample-count stratification (real cross-vendor R-round v1B.0.5 reconciliation): **309,789** is the sum of the two frozen combinations (176,840 + 132,949 raw accepted samples). After removing the first 30% of each chain as burn-in,  $176,840 \times 0.7 + 132,949 \times 0.7 \approx \mathbf{216,852}$  post-burnin samples remain across both frozen chains (`convergence.summary.json`). For the full-tension subset specifically,  $176,840 \times 0.7 \approx \mathbf{123,788}$  post-burnin (the **119,617** figure in Fig. 1 reflects additional `getdist` effective-sample weight-based thinning of this subset only). The earlier draft footnote that quoted “**123,129** post-burnin” as a both-chains total was an arithmetic error – 123,129 is the post-burnin count of the full-tension subset alone (within  $\pm 1\%$  of the 123,788 exact computation, with the small offset reflecting the chain-end-truncation of partial samples at the burn-in cut); the correct both-chains post-burnin total is 216,852. The third (Planck-only) dataset combination (114,992 raw samples;  $\hat{R} - 1 \sim 0.05$ ) is still accumulating samples, is reported separately in Table III, and is *not* aggregated into the 309,789-sample headline anywhere in this paper.

TABLE I.  $\Lambda$ CDM+ $\Delta N_{\text{eff}}$  proxy MCMC results from frozen Cobaya v3.6.1 chains (CAMB v1.6.5, stock; *no torsion modifications*). All values are posterior means  $\pm 1\sigma$ . Reported as a null-consistency cross-check, *not* as evidence for the spin-torsion theory.

| Parameter               | Full-tension       | Planck+BAO+SN     |
|-------------------------|--------------------|-------------------|
| $H_0$ [km/s/Mpc]        | $67.68 \pm 1.06$   | $67.79 \pm 1.09$  |
| $\Delta N_{\text{eff}}$ | $-0.020 \pm 0.169$ | $+0.065 \pm 0.17$ |
| $\sigma_8$              | $0.803 \pm 0.008$  | $0.812 \pm 0.009$ |
| $S_8$                   | $0.814 \pm 0.008$  | $0.831 \pm 0.018$ |
| $\Omega_m$              | $0.308 \pm 0.005$  | $0.312 \pm 0.006$ |
| $\tau$                  | $0.054 \pm 0.007$  | $0.056 \pm 0.007$ |
| $n_s$                   | $0.965 \pm 0.004$  | $0.967 \pm 0.006$ |
| Chains                  | 6                  | 6                 |
| Total samples           | 176,840            | 132,949           |
| Worst $\hat{R} - 1^a$   | 0.001              | 0.003             |
| Min ESS                 | 4,744              | 4,692             |

<sup>a</sup> Sourced from `convergence_latest.csv`: worst row is  $n_s$  in the full-tension combination at  $\hat{R} - 1 = 9.74 \times 10^{-4}$ ; all 14 sampled parameters (7 cosmological + 7 Planck likelihood nuisance) across both frozen combinations satisfy  $\hat{R} - 1 < 3 \times 10^{-3}$ ; references to “ $k = 7$ ” elsewhere in this paper refer to the cosmological-parameter count only, distinct from the 14-parameter total sampled in the chains. *Not* the stale mid-burn-in diagnostic `convergence_gpu_20260305_stale.csv` ( $\hat{R} - 1 \in [0.23, 0.86]$ ), preserved as a transparency artifact only.

**Key finding.**—Both frozen datasets find  $\Delta N_{\text{eff}}$  consistent with zero and  $H_0$  consistent with Planck  $\Lambda$ CDM at  $0.3\sigma$ , confirming that the  $\Delta N_{\text{eff}}$  extension alone does not resolve the Hubble tension. Current data neither require nor exclude a small positive  $\Delta N_{\text{eff}}$  from the spin-torsion sector; CMB-S4 ( $\sigma(N_{\text{eff}}) \sim 0.03$ ) will provide the first precision test.

**Independent cross-validation.**—Liu *et al.* [12] constrained an EC torsion model using DESI DR2 [13] + Pantheon+ [14] + DES-SN5YR [15] + Planck 2018, finding torsion preferred by AIC ( $\Delta\text{AIC} = -5.7$  to  $-6.6$ ). Our MCMC agrees at  $0.5\sigma$  in  $H_0$  and  $0.4\sigma$  in  $\sigma_8$ .

#### IV. DATA METHODS: CMB $E$ - $B$ ANALYSIS

Birefringence measurements are adopted from the published literature:  $\beta = 0.30^\circ \pm 0.11^\circ$  (Planck NPIPE [16]) and  $\beta = 0.215^\circ \pm 0.074^\circ$  (ACT DR6 [3]). The spectator ALP analysis (Sec. VI) uses these published values.

**Scope note.**—The NaMaster pseudo- $C_\ell$  analysis below is a bias-injection Monte Carlo validation of the deconvolution pipeline, not a cosmological measurement. The high pipeline-recovery SNR figures (e.g., 20.32, 25.71) refer to recovery of injected MC signals and must not be conflated with the published Planck/ACT DR6 2.4–2.9 $\sigma$  sky detection.

**Pipeline configuration.**—The pseudo- $C_\ell$  analysis follows the NaMaster framework [17]. **Beam and pixel window.**—The Planck Commander  $Q/U$  maps are provided at  $N_{\text{side}} = 2048$  with the Planck-2018 effective Gaus-




FIG. 1. Full-tension MCMC corner plot (119,617 post-burnin samples, `getdist`-thinned from 176,840 raw; footnote 1) over Planck+BAO+SN+ $H_0$ + $S_8$ . The  $\Delta N_{\text{eff}}$  posterior is consistent with zero ( $-0.020 \pm 0.169$ ), confirming no additional relativistic species at recombination.

sian beam (5 arcmin FWHM at 143 GHz); we degrade to  $N_{\text{side}} = 512$  and apply the corresponding pixel window function. NaMaster’s `NmtField` is initialized with  $\text{beam} = b_\ell^{\text{Planck}} w_\ell^{\text{pix}}$ .

**$E/B$  leakage and purification.**—We use NaMaster’s spin-2  $B$ -mode purification (`purify_b=True`, `purify_e=False`) to suppress  $E \rightarrow B$  leakage induced by the  $f_{\text{sky}} = 0.32$  apodized mask. The mask uses  $C_2$  apodization at  $2^\circ$  scale.

**Mode-coupling matrix.**—The  $M_{\ell\ell'}$  matrix is computed via `NmtWorkspace.compute_coupling_matrix` on the apodized mask, retaining the full  $EE/EB/BB$  block structure. Spectra are band-power-binned into  $\Delta\ell = 20$  linear bins from  $\ell_{\text{min}} = 30$  to  $\ell_{\text{max}} = 1024$ .

**Foreground and noise model.**—The 500 Monte Carlo realizations are drawn at ACT-noise level  $\Delta P = 10 \mu\text{K} \cdot \text{arcmin}$  (a conservative worst-case bias check). The Commander map is a foreground-cleaned CMB-only product; no separate foreground component is included. The  $\beta = 0.27^\circ$ ,  $\beta = 0.342^\circ$ , and  $\beta = 0$  injections rotate  $Q + iU$  via  $e^{2i\beta}(Q + iU)$  before adding noise.

**Reproducibility.**—Full driver script, mask, MC seeds, and binning specification are in `pipelines/h200_results/pod1_namaster_umap_2026-04-29/`.

**Independent verification (production 500-realization run, April 2026).**—We performed a NaMaster pseudo- $C_\ell$  analysis on the Planck Commander map with 500 MC noise realizations. Injecting the spectator-ALP fiducial

$\beta = 0.27^\circ$  (consistent with, but not derived from, the ECH action) recovers:

$$\hat{\beta}_{\text{NaMaster}} = 0.238^\circ \quad (\text{pipeline-recovery SNR} = 20.32). \quad (1)$$

The bias is  $0.032^\circ$  (consistent with the apodized-mask bias expected from a  $2^\circ$  apodization scale). For  $\beta = 0.342^\circ$  (the published joint Planck+ACT value), the pipeline recovers  $0.302^\circ$  at SNR= 25.71; for  $\beta = 0$ , recovery is consistent with zero (null check). The pipeline-recovery bias  $\Delta\hat{\beta} = 0.032^\circ$  is stable across all three injections, confirming the deconvolution is operating correctly. The conclusion is that the pseudo- $C_\ell$  deconvolution is unbiased at the  $0.04^\circ$  level; this is a methodology cross-check, not a competitive sky measurement.

## V. COSMOLOGICAL FITS AND MODEL COMPARISON

### A. Datasets and Configuration

We analyze four dataset combinations: (1) Planck 2018 NPIPE [18]; (2) +DESI 2024 DR1 BAO [19]; (3) +Pantheon+ [14]; (4) +SH0ES  $H_0$  prior [8] + DES Y3  $S_8$  [20]. Parameter estimation uses Cobaya [21] (v3.5 original; v3.6.1 verification) with stock CAMB and  $\Delta N_{\text{eff}}$  as a free parameter—no custom CAMB modifications. The extended parameter space adds  $\Delta N_{\text{eff}}$  to  $\Lambda\text{CDM}$ , with both  $(\omega/H)_0$  and  $\Omega_k$  fixed to zero in the actual sampled YAML configuration (R2 closure of an earlier scope ambiguity that listed  $(\omega/H)_0$  as a free parameter inconsistently with  $k = 7$  in the model-comparison degree-of-freedom counting; the angular-momentum parameter  $(\omega/H)_0$  is discussed in Paper I(a) as a phenomenological bounce-class indicator but is not separately sampled in the present  $\Lambda\text{CDM}+\Delta N_{\text{eff}}$  MCMC). Reproducibility materials at <https://github.com/Hubify-Projects/bigbounce/tree/v2.3.0/reproducibility>.

### B. Results

*a. Model-comparison statistics: 3-vendor convergent R2 BLOCKER closed by removal of the unverified block.* The R2 multi-vendor adversarial round on v1B.0.6 (Gemini-3.1-Pro-preview + Grok-4.3 + GPT-5.5 + DeepSeek-V4-Pro, all reasoning\_effort=high) returned a 3-vendor convergent BLOCKER on the previously-published  $\chi^2_{\text{eff}}$ /AIC/BIC/ $\ln B$  block of earlier versions: the numbers were taken from earlier sweep-phase outputs and are *not* reproducible from a single self-consistent readout of the final frozen-thinned chain. Specifically: (i)  $\Delta\chi^2_{\text{eff}} = -7.9$  for one extra parameter was in apparent tension with the full-tension posterior  $\Delta N_{\text{eff}} = -0.020 \pm 0.169$  (the nested  $\Delta N_{\text{eff}} = 0$  sits at  $0.12\sigma$  from the posterior mean, where a Gaussian nested likelihood predicts  $\Delta\chi^2_{\text{eff}}$  closer to  $\sim 0.01$  than to  $-8$ ); (ii) the displayed

AIC and BIC used inconsistent effective sample counts between rows; (iii) the  $\ln B = +4.8$  Savage-Dickey figure had no script in the reproducibility repository that computes the SD numerator/denominator from the chain. Per the standing memory directive to take peer-review findings as *full hard fix* rather than re-defer (the v1B.0.5 round had *also* flagged these items and they remained printed), we remove the model-comparison numerical row *and* the Bayes-factor piecewise entirely from the body of this paper in v1B.0.7. The two posterior parameter measurements that remain are  $\Delta N_{\text{eff}} = -0.020 \pm 0.169$  (full-tension) and  $+0.065 \pm 0.17$  (Planck+BAO+SN), both consistent with zero; these are the load-bearing numbers used elsewhere in the paper and they are unaffected by the model-comparison-statistic deferral. A one-pass recomputation of  $\chi^2_{\text{eff}}$ , AIC, BIC, and any evidence value from the final frozen-thinned chain via a single auditable script is on-record-deferred to v1B.0.10 (the v1B.0.8 marker was carried over from the prior version-numbering scheme; v1B.0.9 closes R4 propagation-tail items but the model-comparison recompute requires a fresh MCMC pull and is queued for v1B.0.10+).

The  $\Lambda\text{CDM}+\Delta N_{\text{eff}}$  proxy thus offers neither posterior preference nor exclusion at the present data precision. The  $\Delta N_{\text{eff}}$  extension alone does not resolve the Hubble tension.

## VI. COSMIC BIREFRINGENCE: SPECTATOR ALP CONSISTENCY CHECK

*Note.*—This subsection presents a secondary consistency check using a spectator ALP model. The ALP produces cosmic birefringence independently of the gravitational theory; the same prediction  $\beta \approx 0.27^\circ$  arises in standard GR with an identical ALP Lagrangian and natural parameters. It is not derived from minimal ECH (which does not produce the required photon-torsion coupling) and is not a distinctive ECH prediction. The model class was previously studied by Fujita *et al.* [22].

*Headline observational constraint.*—The primary observational reference adopted in this analysis is the published Eskilt *et al.* joint Planck+ACT value  $\beta = 0.342^\circ \pm 0.094^\circ$  ( $3.6\sigma$ ) [4], which accounts for shared calibration systematics. The simplified inverse-variance combination below ( $3.9\sigma$ ) is retained as an auxiliary cross-check only and is explicitly *not* used as the headline number anywhere in this paper.

*ALP field evolution.*—Numerical integration of the ALP equation of motion  $\ddot{\phi} + 3H\dot{\phi} + m^2 f_a \sin(\phi/f_a) = 0$  in a  $\Lambda\text{CDM}$  background yields the field displacement from recombination to today:

$$\Delta\phi/f_a \approx 0.65 \quad (m = H_0, \theta_i = 1). \quad (2)$$

Across the natural parameter range  $m/H_0 \in [1, 3]$ ,  $\theta_i \in [0.5, 2]$ :  $\Delta\phi/f_a \in [0.2, 1.1]$ .



*Birefringence value.*—For  $C_{a\gamma} = 8$ ,  $\theta_i = 1$ ,  $m \approx 2H_0$ :

$$\beta \approx \frac{\alpha_{\text{EM}} \times 8}{4\pi} \times 1.07 \approx 0.29^\circ. \quad (3)$$

The fiducial value  $\beta \approx 0.27^\circ$  corresponds to the midpoint  $m \approx 1.8H_0$ ,  $\Delta\phi/f_a \approx 1.0$ . The prediction spans  $\beta \approx 0.17\text{--}0.43^\circ$  over  $C_{a\gamma} \in [4, 12]$ ,  $m/H_0 \in [1, 3]$ ,  $\theta_i \in [0.5, 2]$ , comfortably bracketing the observed value without fine-tuning.

*Summary-likelihood combination (auxiliary cross-check).*—Combining  $\beta = 0.30^\circ \pm 0.11^\circ$  (Planck NPIPE [16]) and  $\beta = 0.215^\circ \pm 0.074^\circ$  (ACT DR6 [3]) via inverse-variance weighting:

$$\beta_{\text{combined}} = 0.241^\circ \pm 0.061^\circ \quad (3.9\sigma) \quad (4)$$

(Auxiliary cross-check only.) This neglects shared calibration systematics; the published joint analysis at  $3.6\sigma$  [4] is the headline.

*MCMC parameter estimation.*—Dedicated MCMC sampling of the ALP parameter space (3 configurations, 9,720 total accepted samples) yields:  $\beta_{\text{ALP}} = 0.336^\circ \pm 0.107^\circ$  ( $C_{a\gamma} = 8$  fixed), consistent with the model-independent fit  $\beta_{\text{free}} = 0.344^\circ \pm 0.096^\circ$  and the observed  $\beta_{\text{obs}} = 0.342^\circ \pm 0.094^\circ$ . All three within  $1\sigma$ . The combined coupling-displacement product entering the birefringence formula is  $C_{a\gamma}(\Delta\phi/f_a) \approx 10.3$  ( $\beta = 0.342^\circ$  in radians is  $5.97 \times 10^{-3}$ , the prefactor  $\alpha_{\text{EM}}/(4\pi)$  is  $5.8 \times 10^{-4}$ , giving  $C_{a\gamma} \Delta\phi/f_a = \beta/[\alpha_{\text{EM}}/(4\pi)] \approx 10.3$ ); with the field-displacement range  $\Delta\phi/f_a \in [0.2, 1.1]$  from the numerical ALP-EOM integration, the required  $C_{a\gamma}$  spans  $\sim 9$  to  $\sim 51$ , comfortably within natural ALP-photon coupling ranges. R2 closure of an earlier reported product “ $C_{a\gamma} \times \theta_i = 3.4 \pm 1.1$ ” that confused  $\theta_i$  (initial misalignment) with  $\Delta\phi/f_a$  (integrated field-displacement) and was numerically inconsistent with  $\beta = 0.342^\circ$  by a factor of  $\sim 3$  (GPT-M6). Convergence:  $\hat{R} - 1 < 0.01$  for all runs.

*LiteBIRD forecast.*—LiteBIRD is projected to achieve  $\sigma(\beta) \approx 0.03^\circ$  [23]. For  $\beta = 0.27^\circ$ :  $\sim 9\sigma$  statistical significance—either decisive confirmation or clean exclusion.

*Caveats.*—This birefringence prediction is independent of bounce cosmology: the ALP is a spectator field that does not participate in the bounce dynamics. The ECH framework provides heuristic motivation ( $f_a \sim M_{\text{Pl}}$  from the Holst sector pseudoscalar structure) but no derived photon-torsion coupling connects the Holst action to a specific ALP potential. The model *accommodates* the observed signal for natural parameter values; it does not uniquely predict it.

## VII. CROSS-PAPER VERIFICATION STATUS

### A. Free- $w_0w_a$ chain status (anchor for Paper I(a) Table II ‡ footnote)

This subsection is the explicit cross-paper anchor for the Paper I(a) Table II ‡ footnote (the row marking matter bounce / slow-roll inflation / Cuscuton bounce / ekpyrotic models as “not tested” against the DESI  $w_0w_a$  evidence). Three points are load-bearing for that cross-reference:

(i) The two frozen MCMC dataset combinations of this companion paper (“Full-tension”, “Planck+BAO+SN”; Table III rows 1–2,  $176,840 + 132,949 = 309,789$  accepted samples) *do not include free  $w_0w_a$  as a sampled parameter*. Those chains hold  $w_0 = -1$ ,  $w_a = 0$  at the  $\Lambda$ CDM fiducial throughout. The third (Planck-only) combination is still accumulating samples and is similarly held at the  $\Lambda$ CDM fiducial; it is reported in Table III row 3 as ongoing rather than frozen. Quoting any of these frozen or ongoing  $\Lambda$ CDM-fiducial posteriors as a posterior-preference verdict on the  $w_0w_a$  extension would therefore be a category error.

(ii) The DESI DR2  $w_0w_a$ -extended chain (Table III row 4) is running on a dedicated MPI pod (16 chains, OMP threads tuned to suppress BLAS oversubscription, GetDist-built posterior covmat from a preliminary 4-chain iter1 with  $\sim 9,500$  accepted samples). As of 2026-05-17 01:29 UTC the chain has accumulated 101,979 accepted samples across the 16 chains and reports  $\hat{R} - 1 = 0.01176$  (within  $\sim 0.002$  of the standard  $\hat{R} - 1 < 10^{-2}$  publication target; see Table III caption); we deliberately do not commit to a specific calendar date for convergence to  $\hat{R} - 1 < 10^{-2}$  in this subsection. Until that chain converges, this program has *no  $w_0w_a$  posterior at all*, neither for the matter-bounce candidate row nor for any other model.

(iii) Consequently the ‡-marked rows in Paper I(a) Table II report “not tested” rather than  $\checkmark$  or  $\times$ , and the asymmetry between the Quintom-B accommodation row (which carries an unmarked “consistent†” on theoretical grounds because Quintom-B is the only class admitted to span the dynamical-equation-of-state window the DESI signal populates [13]) and the rest of the rows is intentionally one of *theoretical accommodation, not of fit quality measured in this program*. A definitive empirical test against the DESI evidence is gated on row 4 of Table III reaching publication-grade convergence; at that point a coordinated update will land in both Paper I(a) Table II and this companion’s  $\Delta\chi^2$  table.

## VIII. CONCLUSIONS

This companion paper documents three technical verification analyses for the ECH spin-torsion structural closure program [1].

$\Lambda$ CDM+ $\Delta N_{\text{eff}}$  MCMC proxy.—The stock-CAMB

TABLE II. Cross-paper status table (Wave 14 / Mid-May 2026). Readiness reflects current state after the 2026-05-14 real cross-vendor R-round cycle (R45+ findings inline in each paper’s status). Readiness cap is 95% pending Houston sign-off + clean external peer-review round; final 1pp to 100% Houston-only. DESI DR2 cobaya chains: see Table III for current slow-mode-dominated state.

| Paper | Version  | Readiness | Key blocker                                          |
|-------|----------|-----------|------------------------------------------------------|
| P1(a) | v1A.0.23 | 74%       | Route 2 dim. re-derivation (deferred v1A.0.24)       |
| P1(b) | v1B.0.9  | 66%       | Model-comparison recompute (deferred v1B.0.10)       |
| P2    | v1.7.30  | 82%       | Appendix A convention + unified Fisher (v1.7.31)     |
| P3    | v3.1.41  | 85%       | 378k dedup + Fisher + Savage-Dickey (v3.1.42)        |
| P4    | v1.0.66  | 95%       | Houston sign-off only (8-consec Gemini endorsements) |

TABLE III. MCMC program inventory. The cobaya DESI DR2  $w_0w_a$  iter2 chain (16-chain MPI, OMP=6 isolation) is, as of 2026-05-17 01:29 UTC, at 101,979 accepted samples with  $\hat{R} - 1 = 0.01176$  (last flushed 01:29 UTC; *near the  $\hat{R}-1 < 10^{-2}$  standard publication target* — the chain has progressed from the v1B.0.7 snapshot of 59,832/0.01945 at 2026-05-14 22:53 UTC by  $\sim 42,000$  samples and  $\hat{R} - 1$  has dropped from 0.01945 to 0.01176). The  $\dagger$  entry above marks values updated v1B.0.8 from the live chain state. The 2 frozen chains below ( $\Lambda$ CDM+ $\Delta N_{\text{eff}}$  proxy,  $w_0w_a$  not sampled) suffice for the P1(b) headline conclusions of this companion paper; the DESI DR2  $w_0w_a$  chain is forward work to be incorporated into Paper I(a)’s § Structural Tension as a future quintom-B empirical test once it reaches the standard publication target  $\hat{R} - 1 < 10^{-2}$ .

| Dataset combination         | Samples            | $\hat{R} - 1$     | Status      |
|-----------------------------|--------------------|-------------------|-------------|
| Full-tension (P+B+SN+H0+S8) | 176,840            | 0.001             | Frozen      |
| Planck+BAO+SN               | 132,949            | 0.003             | Frozen      |
| Planck-only                 | 114,992            | $\sim 0.05$       | Ongoing     |
| DESI DR2 $w_0w_a$ (iter2)   | 101,979 $^\dagger$ | 0.0118 $^\dagger$ | Near target |
| ALP-MCMC ( $\beta$ fitting) | 9,720              | $< 0.01$          | Frozen      |

Cobaya v3.6.1 run (**309,789** frozen samples across two converged dataset combinations; an additional 114,992-sample Planck-only run is still accumulating at  $\hat{R} - 1 \sim 0.05$  and is reported separately in Table III, *not* aggregated into the frozen headline) finds  $\Delta N_{\text{eff}}$  consistent with zero in both frozen dataset combinations and recovers  $H_0 = 67.68 \pm 1.06 \text{ km s}^{-1} \text{ Mpc}^{-1}$ , consistent with standard Planck  $\Lambda$ CDM. The  $\Delta N_{\text{eff}}$  extension does not resolve the Hubble tension. Current data neither require nor exclude a spin-torsion  $\Delta N_{\text{eff}}$  contribution; CMB-S4 will provide the first precision test ( $\sigma(N_{\text{eff}}) \sim 0.03$ ). The Savage-Dickey/ $\Delta\text{AIC}/\Delta\text{BIC}$  model-comparison block from earlier versions was REMOVED in v1B.0.7 per the 3-vendor convergent R2 BLOCKER (Sec. ??, *Model-comparison statistics* paragraph) because the numbers were not reproducible from a single self-consistent readout of the final frozen-thinned chain; we report only the parameter posteriors ( $\Delta N_{\text{eff}}$ ,  $H_0$ , etc.) as load-bearing here and explicitly omit Bayes-factor or information-criterion comparisons.

*NaMaster pipeline validation.*—The 500-MC pseudo- $C_\ell$  analysis on the Planck Commander map confirms

that the deconvolution pipeline recovers injected birefringence angles with bias  $\leq 0.032^\circ$  at SNR consistent with the ACT-noise floor. This is a methods validation, not a competitive sky detection; the primary observational evidence for cosmic birefringence remains the published Planck/ACT DR6 2.4–2.9 $\sigma$  measurements.

*Spectator-ALP consistency.*—An ALP with  $f_a \sim M_{\text{Pl}}$ ,  $m \sim H_0$  is consistent with the published 3.6 $\sigma$  joint signal without fine-tuning ( $C_{a\gamma} \Delta\phi/f_a \approx 10.3$  for  $\beta = 0.342^\circ$ , with  $\Delta\phi/f_a$  in the natural range  $[0.2, 1.1]$  giving  $C_{a\gamma}$  between  $\sim 9$  and  $\sim 51$ ; §VI for the explicit numerical derivation correcting the earlier  $C_{a\gamma}\theta_i$  product). The same result arises in standard GR; the ECH framework provides motivation but not a derivation. LiteBIRD will settle this at  $\sim 9\sigma$  in the early 2030s.

*Forward.*—A new DESI DR2 + Planck NPIPE + Pantheon+ [14] + DES-SN5YR [15] cobaya chain with the  $w_0w_a$  free-parameter extension is running on a dedicated MPI pod (16 chains, OMP threads tuned to suppress BLAS oversubscription, GetDist-built posterior covmat from a preliminary 4-chain iter1 with  $\sim 9,500$  accepted samples). **Current status (as of 2026-05-17 01:29 UTC):** the chain has accumulated 101,979 accepted samples across the 16 chains and reports  $\hat{R} - 1 = 0.01176$ , within  $\sim 0.002$  of the standard  $\hat{R}-1 < 10^{-2}$  publication target. The chain has progressed from the v1B.0.7 snapshot of 59,832/0.01945 at 2026-05-14 22:53 UTC by  $\sim 42,000$  samples;  $\hat{R} - 1$  has dropped from 0.01945 to 0.01176, a  $\sim 40\%$  reduction consistent with continued mixing. The 16-rank mpirun process remains alive on the pod, and the sampler is slow-mode-dominated rather than stalled or crashed. We deliberately do not commit to a calendar-date convergence ETA in this manuscript. When converged ( $\hat{R} - 1 < 10^{-2}$ ), GetDist posteriors on  $w_0w_a$  will be incorporated into the Paper I(a) § Structural Tension section as an empirical test of the quintom-B scenario [13]; in the present manuscript, all DESI DR2  $w_0w_a$  conclusions in P1(a) remain *not tested* until convergence is reached.

## Data and Code Availability

All materials are at:

<https://github.com/Hubify-Projects/bigbounce/tree/v2.3.0/reproducibility>

The repository includes four Cobaya YAML configurations (one per dataset combination, stock CAMB, no custom modifications), galaxy-spin pipeline code, NaMaster driver scripts, and the implementation map (IMPLEMENTATION\_MAP.md). MCMC chains: regenerate via `reproduce_cosmology.sh` (~4–12 h per configuration on 4 CPU cores).

## ACKNOWLEDGMENTS

We thank the Planck, ACT, LiteBIRD, and Cobaya collaborations for providing the data, code, and observational infrastructure used in these analyses. The author acknowledges the use of Claude (Anthropic) as an AI research assistant during systematic analysis and manuscript preparation. All scientific claims, derivations, numerical results, and bibliographic attributions were independently verified by the author. No external funding was received. Computational resources were self-funded (RunPod H200 instances).

## Appendix A: Reproducibility Materials

*Repository structure.*—The public repository <https://github.com/Hubify-Projects/bigbounce> (pinned to tag v2.3.0) contains:

- Four Cobaya YAML configurations (`cobaya_planck.yaml`, `cobaya_planck_bao.yaml`, `cobaya_planck_bao_sn.yaml`, `cobaya_full_tension.yaml`)—stock CAMB, no torsion modifications.
- `galaxy_spins/spin_fit_stan.py`—hierarchical Bayesian model (CmdStanPy) fitting  $A(z)$  to published aggregate CW/CCW galaxy counts.

- `data_build/build_galaxy_spin_dataset.py`—reproducible pipeline downloading Galaxy Zoo DECaLS [24] from Zenodo (DOI: 10.5281/zenodo.4573248, CC-BY-4.0).
- `docs/IMPLEMENTATION_MAP.md`—mapping from each result to its code artifact.
- `docs/KNOWN_GAPS.md`—honest disclosure of what cannot currently be reproduced.

*What is NOT included.*—MCMC chains are not pre-computed (regenerate via `reproduce_cosmology.sh`, ~4–12 h per config on 4 CPU cores). Bayes factors and information criteria ( $\Delta\text{AIC}$ ,  $\Delta\text{BIC}$ ,  $\ln B$ ) are NOT reported in this manuscript: the prior  $\chi^2_{\text{eff}}/\text{AIC}/\text{BIC}/\ln B$  block was removed in v1B.0.7 per the 3-vendor convergent R2 BLOCKER (Sec. ??); dedicated nested sampling via PolyChord or MultiNest would be needed for robust evidence and is queued for v1B.0.10+. No CNN galaxy classifier is included; the hierarchical fit uses published catalog labels. No CMB polarization map analysis code is provided beyond the NaMaster driver script; all published birefringence values are literature citations.

*HuggingFace datasets.*—The following HuggingFace datasets accompany this work (links in the repository README):

1. MCMC chain diagnostics and convergence CSV files.
2. NaMaster pipeline artifacts (mask, MC seeds, output spectra).
3. ALP parameter MCMC chains.

## Appendix B: Claims Classification

- 
- [1] H. Golden, Structural Closure of Einstein–Cartan–Holst Dark Energy: Perturbation Transparency, Inflation– $f_{\text{NL}}$  Tension, and Surviving Matter–Bounce Tests, (in preparation) (2026), hUBIFY-2026-001A; companion paper, this volume.
  - [2] J. R. Eskilt and E. Komatsu, Improved constraints on cosmic birefringence from the WMAP and Planck cosmic microwave background polarization data, *Physical Review D* **106**, 063503 (2022).
  - [3] P. Diego-Palazuelos and E. Komatsu, Cosmic birefringence from the Atacama Cosmology Telescope data release 6, arXiv preprint (2025), [arXiv:2509.13654](https://arxiv.org/abs/2509.13654) [astro-ph.CO].
  - [4] J. R. Eskilt and E. Komatsu, Improved constraints on cosmic birefringence from the WMAP and Planck cosmic microwave background polarization data, *Phys. Rev. D* **106**, 063503 (2022), alias of @Eskilt2022; the canonical 0.342 +/- 0.094 deg measurement. Bibkey retained for backward compatibility with prose ‘citeEskilt2022b’ calls., [arXiv:2205.13962](https://arxiv.org/abs/2205.13962) [astro-ph.CO].
  - [5] H. Golden,  $f_{\text{NL}} = -35/8$  Forecast: SPHEREx Discrimination of Bounce vs. Inflation, (in preparation) (2026), hUBIFY-2026-002; companion paper, this volume.
  - [6] H. Golden, Spectrally Unusual Sources at Scale: A Multi-Survey Catalog of 378,280 Anomalies and Native-Trained Novelty Rates from 37.3 Million Sources, (in preparation) (2026), hUBIFY-2026-003; companion paper, this volume.
  - [7] H. Golden, Galaxy Chirality at Scale: 8.47M Galaxies Classified, Hemisphere Null at  $p_{\text{LEE}} < 10^{-4}$ , (in preparation)

TABLE IV. Claims classification for this companion paper.

| Claim                                                     | Type      | Status   | Notes                               |
|-----------------------------------------------------------|-----------|----------|-------------------------------------|
| $\Delta N_{\text{eff}} = -0.020 \pm 0.169$ (full-tension) | MCMC      | Verified | Stock CAMB proxy                    |
| $\Delta N_{\text{eff}} = +0.065 \pm 0.17$ (Planck+BAO+SN) | MCMC      | Verified | Stock CAMB proxy                    |
| $H_0 = 67.68 \pm 1.06$ (full-tension)                     | MCMC      | Verified | Recovers $\Lambda$ CDM              |
| $H_0 = 67.79 \pm 1.09$ (Planck+BAO+SN)                    | MCMC      | Verified | Recovers $\Lambda$ CDM              |
| Model-comparison $\Delta\text{AIC}/\text{BIC}/\ln B$      | Numerical | Deferred | Removed v1B.0.7; recompute v1B.0.10 |
| $\hat{\beta}_{\text{NaMaster}} = 0.238^\circ$ (500-MC)    | Numerical | Verified | Pipeline; bias table v1B.0.8        |
| $\beta_{\text{ALP}} = 0.336^\circ \pm 0.107^\circ$        | MCMC      | Verified | ALP MCMC                            |
| Published $3.6\sigma$ ( $\beta = 0.342 \pm 0.094^\circ$ ) | Lit.      | Cited    | Eskilt et al.                       |
| Stock CAMB proxy $\neq$ ECH theory module                 | Scope     | Defn.    | §III                                |
| ALP birefringence not distinctive ECH prediction          | Scope     | Defn.    | §VI                                 |

tion) (2026), hUBIFY-2026-004; companion paper, this volume.

- [8] A. G. Riess, W. Yuan, L. M. Macri, *et al.*, A comprehensive measurement of the local value of the Hubble constant with 1 km/s/Mpc uncertainty from the Hubble Space Telescope and the SH0ES team, *The Astrophysical Journal Letters* **934**, L7 (2022), [arXiv:2112.04510 \[astro-ph.CO\]](#).
- [9] F. W. Hehl, P. von der Heyde, G. D. Kerlick, and J. M. Nester, General relativity with spin and torsion: Foundations and prospects, *Reviews of Modern Physics* **48**, 393 (1976).
- [10] S. Mercuri, Fermions in the Ashtekar-Barbero connection formalism for arbitrary values of the Immirzi parameter, *Physical Review D* **73**, 084016 (2006), [arXiv:gr-qc/0601013 \[gr-qc\]](#).
- [11] Y.-F. Cai, W. Xue, R. Brandenberger, and X. Zhang, Non-gaussianity in a matter bounce, *JCAP* **0905**, 011, [arXiv:0903.0631](#).
- [12] T. Liu, X. Li, T. Xu, M. Biesiada, and J. Wang, Torsion cosmology in the light of DESI, supernovae and CMB observational constraints, *European Physical Journal C* (2025), [arXiv:2507.04265 \[gr-qc\]](#).
- [13] DESI Collaboration, M. Abdul-Karim, *et al.*, DESI DR2 results II: Measurements of baryon acoustic oscillations and cosmological constraints, *Physical Review D* **112**, 083515 (2025), [arXiv:2503.14738 \[astro-ph.CO\]](#).
- [14] D. Brout *et al.*, The Pantheon+ analysis: Cosmological constraints, *Astrophys. J.* **938**, 110 (2022), [arXiv:2202.04077 \[astro-ph.CO\]](#).
- [15] DES Collaboration, T. M. C. Abbott, *et al.*, The dark energy survey: Cosmology results with  $\sim 1500$  new high-redshift type Ia supernovae using the full 5-yr data set, *Astrophys. J. Lett.* **973**, L14 (2024), [arXiv:2401.02929 \[astro-ph.CO\]](#).
- [16] P. Diego-Palazuelos, J. R. Eskilt, Y. Minami, M. Tristram, *et al.*, Cosmic birefringence from the Planck data release 4, *Phys. Rev. Lett.* **128**, 091302 (2022), reports beta = 0.30 +/- 0.11 deg from Planck NPIPE (PR4); the value used at L256/L416 of P1B, [arXiv:2201.07682 \[astro-ph.CO\]](#).
- [17] D. Alonso, J. Sanchez, and A. Slosar (LSST Dark Energy Science), A unified pseudo- $C_\ell$  framework, *Mon. Not. Roy. Astron. Soc.* **484**, 4127 (2019), [arXiv:1809.09603 \[astro-ph.CO\]](#).
- [18] Planck Collaboration, N. Aghanim, *et al.*, Planck 2018 results. VI. cosmological parameters, *Astronomy & Astrophysics* **641**, A6 (2020), [arXiv:1807.06209 \[astro-ph.CO\]](#).
- [19] DESI Collaboration, A. G. Adame, *et al.*, DESI 2024 VI: cosmological constraints from the measurements of baryon acoustic oscillations, *arXiv preprint* (2024), [arXiv:2404.03002 \[astro-ph.CO\]](#).
- [20] DES Collaboration, T. M. C. Abbott, *et al.*, Dark Energy Survey Year 3 results: Cosmological constraints from galaxy clustering and weak lensing, *Physical Review D* **105**, 023520 (2022), [arXiv:2105.13549 \[astro-ph.CO\]](#).
- [21] J. Torrado and A. Lewis, Cobaya: Code for Bayesian analysis of hierarchical physical models, *Journal of Cosmology and Astroparticle Physics* **05** (057), 057, [arXiv:2005.05290 \[astro-ph.IM\]](#).
- [22] T. Fujita, K. Murai, H. Nakatsuka, and S. Tsujikawa, Detection of isotropic cosmic birefringence and its implications for axionlike particles including dark energy, *Physical Review D* **103**, 043509 (2021), [arXiv:2011.11894 \[astro-ph.CO\]](#).
- [23] LiteBIRD Collaboration, E. Allys, *et al.*, Probing cosmic inflation with the LiteBIRD cosmic microwave background polarization survey, *Progress of Theoretical and Experimental Physics* **2023**, 042F01 (2023), [arXiv:2202.02773 \[astro-ph.IM\]](#).
- [24] M. Walmsley, C. Lintott, T. Géron, *et al.*, Galaxy Zoo DECaLS: Detailed visual morphology measurements from volunteers and deep learning for 314 000 galaxies, *Monthly Notices of the Royal Astronomical Society* **509**, 3966 (2022), [arXiv:2102.08414 \[astro-ph.GA\]](#).