

Testing the Matter Bounce with Primordial Non-Gaussianity: Forecasts for SPHEREx and MegaMapper

Houston Golden^{ORCID}

Independent Researcher

houston@hubify.com

<https://bigbounce.hubify.app>

March 19, 2026

Abstract

A matter-dominated contracting phase preceding a nonsingular bounce produces a specific, parameter-free prediction for local-type primordial non-Gaussianity: $f_{\text{NL}}^{\text{local}} = -35/8 = -4.375$ [Cai et al., 2009]. This value is approximately 300 times larger than the standard single-field inflationary prediction and opposite in sign. We present forecasts for testing this prediction with the SPHEREx and MegaMapper galaxy surveys via the scale-dependent bias effect and the galaxy bispectrum. In the squeezed limit relevant for large-scale structure estimators, the matter-bounce bispectrum shape projects exactly onto the local template with no projection penalty. The SPHEREx multi-tracer galaxy bispectrum achieves $\sigma(f_{\text{NL}}) \approx 0.7$ [Heinrich et al., 2023], giving approximately 4–6 σ sensitivity to the bounce signal after accounting for photometric redshift degradation, PNG bias uncertainty, and relativistic projection effects. MegaMapper’s spectroscopic multi-tracer capability could reach $\sigma(f_{\text{NL}}) \approx 0.5$ under ideal conditions (3–7 σ realistic, conditional on ultra-large-scale systematics modeling). We perform a Bayesian model comparison using 800,000 Monte Carlo realizations across analytic, mock-based, and GR-aware frameworks, finding that a detection near $f_{\text{NL}} = -4.375$ would favor the bounce over standard single-field inflation at Bayes factor > 300 and over tuned multifield competitors at Bayes factor > 7 , with these conclusions robust to conservative treatment of relativistic projection effects. A null result from SPHEREx would disfavor the quasi-dust matter bounce at $> 4\sigma$ significance.

1 Introduction

The inflationary paradigm provides a remarkably successful framework for generating the observed spectrum of primordial perturbations. Standard single-field slow-roll inflation predicts a nearly scale-invariant, nearly Gaussian spectrum with a small, positive local-type non-Gaussianity $f_{\text{NL}} \approx (5/12)(1 - n_s) \approx 0.015$, set by the Maldacena consistency relation [Maldacena, 2003].

Bouncing cosmology offers an alternative origin for primordial perturbations: modes exit the Hubble radius during a contracting phase and re-enter after a nonsingular bounce. In particular, a matter-dominated contraction ($w \approx 0$) produces a scale-invariant scalar spectrum through the growth of the curvature perturbation ζ on superhorizon scales [Wands, 1999, Finelli and Brandenberger, 2002].

A distinctive prediction of the matter bounce is a large, negative, parameter-free local-type non-Gaussianity $f_{\text{NL}} = -35/8 = -4.375$ [Cai et al., 2009]. This value is determined entirely by the equation of state during contraction ($\epsilon = 3/2$ for matter) and the structure of the Maldacena cubic action, with no free parameters. The prediction is mechanism-independent: it holds regardless of whether the bounce is realized by loop quantum cosmology [Wilson-Ewing, 2013], Einstein-Cartan-Holst gravity, or other nonsingular completions. In minimal

Einstein-Cartan-Holst gravity, scalar perturbations reduce exactly to the standard Mukhanov-Sasaki sector because the Holst term becomes a topological invariant when torsion vanishes for canonical scalar field matter, rendering the Barbero-Immirzi parameter invisible in all scalar observables.

The next generation of galaxy surveys—SPHEREx [Dore et al., 2014] and MegaMapper [Schlegel et al., 2022]—will constrain local-type f_{NL} at unprecedented precision through the scale-dependent bias effect [Dalal et al., 2008] and the galaxy bispectrum [Heinrich et al., 2023]. In this paper, we present hardened forecasts for testing $f_{\text{NL}} = -35/8$ with these surveys, including a systematic assessment of the dominant observational fragilities and a Bayesian model comparison quantifying the discrimination power against inflationary alternatives.

2 The Matter-Bounce Bispectrum Benchmark

2.1 The Prediction

In a matter-dominated contracting universe with standard GR perturbation theory and Bunch-Davies vacuum, the curvature perturbation ζ grows as $|\eta|^{-3}$ on superhorizon scales during contraction. The cubic interactions, governed by the Maldacena action [Maldacena, 2003] specialized to $\epsilon = 3/2$, produce a bispectrum with shape function [Cai et al., 2009]:

$$A_T(k_1, k_2, k_3) = \frac{3}{256 k_1^2 k_2^2 k_3^2} P(k_1, k_2, k_3), \quad (1)$$

where P is a degree-9 homogeneous polynomial in the wavenumbers. The nonlinearity parameter in the squeezed limit is:

$$|B|_{\text{NL}} = \frac{10}{3} \frac{A_T}{\sum_i k_i^3} \rightarrow -\frac{35}{8} \quad \text{as} \quad k_1/k \rightarrow 0. \quad (2)$$

We have independently verified this result by evaluating the shape function at three distinct momentum configurations (Table 1 and Fig. 1).

Configuration	$ B _{\text{NL}}$ (this work)	$ B _{\text{NL}}$ (Cai et al.)
Squeezed ($k_1 \rightarrow 0$)	-4.375	-35/8
Equilateral ($k_1 = k_2 = k_3$)	-3.984	-255/64
Folded ($k_1 = 2k_2 = 2k_3$)	-2.250	-9/4

Table 1: Independent verification of the matter-bounce shape function at three benchmark momentum configurations. All values match the published results [Cai et al., 2009] exactly.

2.2 Mechanism Independence

The prediction depends only on: (a) matter-dominated contraction ($w \approx 0$, $\epsilon \approx 3/2$), (b) standard GR perturbation theory during contraction, and (c) Bunch-Davies vacuum initial conditions. It does not depend on the specific bounce mechanism. The bounce enters only through providing a nonsingular transition and transferring the contraction-phase perturbations into the expanding phase.

2.3 The Viable Model

The Wilson-Ewing Λ CDM quasi-dust model [Wilson-Ewing, 2013] provides a complete observational package: $n_s = 0.964$ (from $w = -0.003$, one free parameter), $r \approx 10^{-4}$ (from LQC quantum-geometry tensor suppression), and $f_{\text{NL}} = -35/8$ (parameter-free). This model has no current observational tensions.




Figure 1: Matter-bounce bispectrum shape function $|B|_{\text{NL}}(k_1, k, k)$ as a function of the squeeze ratio k_1/k , showing convergence to $-35/8$ in the squeezed limit. Red circle: squeezed benchmark. Orange square: equilateral. Green triangle: folded.

3 Observable Mapping to Large-Scale Structure

3.1 Scale-Dependent Bias

Primordial local non-Gaussianity induces a scale-dependent correction to galaxy bias [Dalal et al., 2008]:

$$\Delta b(k) = \frac{2(b_1 - 1) f_{\text{NL}} \delta_c}{D(z) T(k) \alpha(k)}, \quad (3)$$

where $\alpha(k)$ encodes the Poisson equation normalization. The signal grows as $1/k^2$ on the largest scales.

3.2 Template Projection

For LSS surveys using the scale-dependent bias estimator, the relevant bispectrum configurations are in the squeezed limit ($k_{\text{long}} \ll k_{\text{short}}$). In this limit, the matter-bounce bispectrum converges to exactly $-35/8$ —the local template value. There is no template-projection penalty for LSS estimators.

3.3 Galaxy Bispectrum

The galaxy bispectrum provides an independent measurement channel that accesses information at shorter wavelengths, reducing the dependence on ultra-large-scale modes [Heinrich et al., 2023]. This makes bispectrum-based constraints more robust to large-scale systematics than power-spectrum-based scale-dependent bias alone.

4 SPHEREx Forecast

SPHEREx is an all-sky spectrophotometric survey ($0.75\text{--}5\ \mu\text{m}$) with spectral resolution $R \approx 40\text{--}130$ and approximately 450 million galaxies. A dedicated multi-tracer bispectrum analysis [Hein-

rich et al., 2023] forecasts $\sigma(f_{\text{NL}}) = 0.7$ from the bispectrum alone, with $\sigma(f_{\text{NL}}) = 0.5$ when combined with the power spectrum.

For our target signal $f_{\text{NL}} = -4.375$, the detection significance ranges from 6.3σ (bispectrum only, ideal) to 3.4σ (conservative, with GR marginalization at $\sigma_{\text{GR}} = 1.0$). The detection significance across survey scenarios is summarized in Fig. 2. SPHEREx provides the most robust near-term test because: (a) the bispectrum channel avoids ultra-large-scale mode dependence, (b) lower redshift ($z \approx 1.5$) reduces GR projection contamination, and (c) multi-tracer across redshift bins provides effective cosmic variance cancellation.




Figure 2: Detection significance for $f_{\text{NL}} = -35/8$ across survey configurations. Error bars show optimistic-to-conservative ranges accounting for multi-tracer, photo- z , bias, and GR systematics.

5 MegaMapper Forecast

MegaMapper is a proposed Stage-V spectroscopic survey targeting ~ 10 million Lyman-break galaxies at $z = 2-5$ with multi-tracer capability [Schlegel et al., 2022]. Published forecasts give $\sigma(f_{\text{NL}}) \approx 0.5$ under ideal conditions.

The significance ranges from 8.75σ (design goal) to $3-5\sigma$ (conservative, after GR marginalization and b_ϕ uncertainty). MegaMapper’s forecast is more sensitive than SPHEREx’s to: (a) relativistic projection effects, which create substantial GR-induced bias at $z > 2$ [Jolicoeur et al., 2025]; (b) PNG bias parameter b_ϕ uncertainty, which can degrade constraints if uncalibrated [Barreira, 2022]; and (c) multi-tracer implementation quality.

6 Inflation Mimicry and Bayesian Comparison

6.1 Can Inflation Reproduce the Signal?

Standard single-field slow-roll inflation predicts $f_{\text{NL}} = (5/12)(1 - n_s) \approx +0.015$ [Maldacena, 2003]—wrong by a factor of 300 and opposite in sign. Non-canonical single-field models (DBI, etc.) produce equilateral-shape f_{NL} , not local.

Non-attractor single-field inflation naturally gives $f_{\text{NL}} = +5/2$ (wrong sign) [Cai et al., 2018]. Reaching -4.375 requires engineering the attractor-to-slow-roll transition. The standard

quadratic curvaton gives minimum $f_{\text{NL}} \approx -1.25$ (insufficient). Self-interacting curvatons or curved field-space models can reach -4.375 but require ≥ 2 tuned parameters.




Figure 3: f_{NL} landscape: matter bounce vs. inflationary alternatives. The bounce prediction (red diamond) is parameter-free; inflationary alternatives require free parameters to reach the same region. SPHEREx 1σ error bar shown in blue.

6.2 The Kinematic vs. Parametric Asymmetry

The bounce predicts $f_{\text{NL}} = -35/8$ *kinematically*, with zero free parameters in the cubic sector. Inflation can only *accommodate* this value parametrically, requiring extra fields and tuned couplings. This asymmetry drives a natural Bayesian preference for the bounce.

6.3 Quantitative Bayesian Comparison

We performed model comparison using 800,000 Monte Carlo realizations across three frameworks (analytic closed-form, mock power spectrum generation with fitting, and GR-aware marginalization).

For a detection at $f_{\text{NL}} = -4.375$ by SPHEREx ($\sigma = 0.7$), the median Bayes factors are (Table 2):

Comparison	Median Bayes Factor	$P(\text{BF} > 3)$
Bounce vs. standard single-field	$> 10^5$	97%
Bounce vs. tuned multifield $[-15, +15]$	10–23	87–96%

Table 2: Bayesian model comparison for a mock detection at $f_{\text{NL}} = -4.375$ with $\sigma = 0.7$. Ranges reflect different GR treatment scenarios.

These results are robust to prior sensitivity: the Bayes factor vs. tuned multifield ranges from 7 (narrow prior) to 57 (broad prior) under all reasonable prior choices.

7 Systematics and GR-Aware Robustness

7.1 Dominant Fragilities

Our Fisher robustness scan identified three dominant threats to the forecast: (1) ultra-large-scale mode access (k_{min}), where the SDB signal is concentrated in the lowest k -modes (Fig. 4);

(2) relativistic projection effects, which create a GR-induced bias at large scales; and (3) PNG bias (b_ϕ) uncertainty, which degrades the SDB calibration [Barreira, 2022].




Figure 4: Left: $\sigma(f_{NL})$ vs. minimum accessible wavenumber for MegaMapper (orange) and SPHEREx SDB-only (blue). The SPHEREx bispectrum channel ($\sigma = 0.7$, dotted) avoids the ultra-large-scale fragility. Right: corresponding detection significance for $f_{NL} = -35/8$.

7.2 GR-Aware Analysis

We performed GR-aware Bayesian comparison across five scenarios (Table 3). Even under conservative GR marginalization ($\sigma_{GR} = 1.0$), the bounce is favored at Bayes factor > 300 over standard inflation and > 7 over tuned multifield.

GR Treatment	BF vs. SSFSR	BF vs. Tuned	$P(\text{BF} > 3)$ vs. SSFSR
Ideal (no GR)	3.3×10^6	10.9	98%
Marginalized ($\sigma_{GR} = 0.5$)	4.1×10^4	9.4	97%
Marginalized ($\sigma_{GR} = 1.0$)	329	7.9	96%
Corrected (10% residual)	3.3×10^6	10.9	98%

Table 3: GR-aware Bayesian comparison. The bounce-vs-inflation comparison survives all treatment scenarios.

8 Discussion

8.1 The Staged Observational Strategy

SPHEREx (~ 2028) provides the first real test via the galaxy bispectrum at ~ 4 – 6σ significance. MegaMapper ($\sim 2032+$, if funded) provides the more powerful but fragile follow-up at 3 – 7σ via scale-dependent bias.

8.2 Decision Thresholds

A measurement of $f_{NL} = -4 \pm 1$ by SPHEREx would provide strong evidence favoring a contracting/bounce origin over standard single-field inflation. A null result (f_{NL} consistent with zero at the 2σ level) would strongly disfavor the quasi-dust matter bounce.




Figure 5: Observational decision thresholds. Green: strongly favors bounce. Red: kills the matter-bounce hypothesis. Blue vertical line: bounce prediction $f_{\text{NL}} = -35/8$. Error bars: SPHEREx ($\sigma = 0.7$) and MegaMapper conservative ($\sigma = 1.5$).

8.3 Caveats

We emphasize that a detection of $f_{\text{NL}} \approx -4$ would constitute strong evidence *favoring* the bounce over standard inflation, not unique proof of a pre-Big-Bang contracting phase. Exotic multifield inflationary constructions can in principle accommodate this value, though at the cost of additional free parameters and engineering. The detection significance is conditional on the quality of GR projection modeling at ultra-large scales and the PNG bias parameter calibration.

We also note that appending late-time dynamical-dark-energy freedom (e.g., CPL parametrization) to a bounce model can improve cosmological fits at the parameter level, as explored in recent bounce + dark-energy literature. However, such phenomenological freedom does not derive from the bounce physics itself and does not constitute first-principles evidence for a contracting phase. Our analysis restricts attention to the parameter-free prediction $f_{\text{NL}} = -35/8$, which is controlled by the contraction dynamics.

9 Conclusion

The quasi-dust matter bounce makes a specific, parameter-free, falsifiable prediction: $f_{\text{NL}}^{\text{local}} = -35/8$. This value is mechanism-independent, approximately 300 times larger than the standard inflationary prediction, and opposite in sign. We have shown that SPHEREx can test this prediction at 4–6 σ significance through the multi-tracer galaxy bispectrum, with MegaMapper providing a more powerful but systematics-sensitive follow-up.

Our Bayesian model comparison, based on 800,000 Monte Carlo realizations across multiple frameworks including realistic GR contamination treatment, demonstrates that a detection near $f_{\text{NL}} = -4.375$ would robustly favor the bounce over standard single-field inflation (Bayes factor > 300 even under conservative systematics) and over tuned multifield competitors (Bayes factor > 7). These conclusions are robust to prior sensitivity and survive conservative treatment of all identified systematic effects.

The matter-bounce bispectrum provides what may be the sharpest single observable for distinguishing the bounce paradigm from standard inflation. SPHEREx data, expected around 2028, will provide the first meaningful test.

Acknowledgments

The author thanks the developers of the open-source tools used in this work, including NumPy, SciPy, Matplotlib, mpmath, and Tectonic. Computations were performed on consumer hardware; no dedicated HPC resources were required for any result in this paper. The author

acknowledges helpful interactions with AI research assistants during the systematic audit and verification phases of this work.

References

- Yi-Fu Cai, Wei Xue, Robert Brandenberger, and Xinmin Zhang. Non-gaussianity in a matter bounce. *JCAP*, 0905:011, 2009.
- Chen Heinrich, Olivier Dore, and Elisabeth Krause. Measuring f_{nl} with the spherex multi-tracer redshift space bispectrum. 2023.
- Juan Maldacena. Non-gaussian features of primordial fluctuations in single field inflationary models. *JHEP*, 0305:013, 2003.
- David Wands. Duality invariance of cosmological perturbation spectra. *Phys. Rev. D*, 60:023507, 1999.
- Fabio Finelli and Robert Brandenberger. On the generation of a scale-invariant spectrum of adiabatic fluctuations in cosmological models with a contracting phase. *Phys. Rev. D*, 65:103522, 2002.
- Edward Wilson-Ewing. The matter bounce scenario in loop quantum cosmology. *JCAP*, 1303:026, 2013.
- Olivier Dore et al. Cosmology with the spherex all-sky spectral survey. 2014.
- David J. Schlegel et al. The megamapper: A stage-5 spectroscopic instrument concept. 2022.
- Neal Dalal, Olivier Dore, Dragan Huterer, and Alexander Shirokov. The imprints of primordial non-gaussianities on large-scale structure. *Phys. Rev. D*, 77:123514, 2008.
- Sheean Jolicoeur et al. Unbiased analysis of primordial non-gaussianity: the multipoles of the full relativistic power spectrum. 2025.
- Alexandre Barreira. Can we actually constrain f_{nl} using the scale-dependent bias effect? 2022.
- Yi-Fu Cai, Xingang Chen, M. H. Namjoo, Misao Sasaki, Dong-Gang Wang, and Ziwei Wang. Revisiting non-gaussianity from non-attractor inflation models. *JCAP*, 2018.