

Cosmic Birefringence from a Planck-Scale Axion-Like Particle: Predictions, Constraints, and LiteBIRD Forecasts

Houston Golden

Independent Researcher, Los Angeles, California, USA

houston@hubify.com

June 9, 2026

Abstract

We present predictions and constraints for cosmic birefringence from a spectator axion-like particle (ALP) with Planck-scale decay constant $f_a \sim M_{\text{Pl}}$ and mass $m \sim H_0$. For order-unity initial misalignment $\theta_i \sim \mathcal{O}(1)$ and order-unity photon anomaly coefficient $C_0 \sim \mathcal{O}(1)$, this minimal setup yields a birefringence rotation angle $\beta \approx 0.27^\circ$, consistent with the 3.6σ isotropic birefringence signal ($\beta_{\text{obs}} = 0.342 \pm 0.094^\circ$ from the ESKILT et al. joint Planck + ACT analysis). *Scope of the naturalness claim:* along an ALP trajectory in a Λ CDM background, the photon-coupling combination $g_{a\gamma} \propto 1/f_a$ and the field displacement $\Delta\phi \propto f_a \theta_i F(m/H_0)$ both scale with f_a , so the rotation amplitude $\beta = (g_{a\gamma}/2) \Delta\phi \approx (C_0 \theta_i/2) F(m/H_0)$ is *independent of f_a* ; the “ $f_a \sim M_{\text{Pl}}$ ” choice is required by EFT consistency for a gravitationally coupled pseudoscalar and by the spectator-condition energy-density constraint $\Omega_\phi \ll 1$ (see Sec. 5), not by tuning to β_{obs} . The match to the observed signal depends on θ_i and C_0 both being $\mathcal{O}(1)$ at their natural prior values. We perform a Gaussian summary-likelihood inference using Planck HFI and ACT DR6 data, finding $\beta = 0.242 \pm 0.061^\circ$ (3.9σ from zero) with an effective photon coupling $f_{\text{photon}} \times C_0 = 1.73 \pm 0.44$ (order-unity, no fine-tuning). The Bayes factor in favor of nonzero rotation is $\ln B = 5.17$ (indicative; prior-dependent, see Sec. 3.4). We forecast that LiteBIRD, with $\sigma(\beta) \approx 0.03^\circ$, will test this prediction at 9σ significance—either confirming the signal or ruling out the ALP explanation decisively. This birefringence prediction is independent of bounce cosmology and can be tested regardless of whether the universe underwent a contracting phase.

1 Introduction

Cosmic birefringence—the uniform rotation of the polarization plane of CMB photons as they propagate from the last scattering surface to the observer—has emerged as a compelling signal in recent CMB data. The Planck HFI analysis [Minami and Komatsu, 2020] reported $\beta = 0.35 \pm 0.14^\circ$ (2.5σ), and the ACT DR6 analysis confirmed the signal at comparable significance. Combined, the evidence exceeds 3.5σ .

The leading theoretical explanation is a light pseudoscalar (axion-like particle, ALP) that couples to the electromagnetic Chern-Simons term $\phi F\tilde{F}$. If the ALP field evolves between recombination and today, the accumulated phase difference between left- and right-circular polarizations produces a net rotation $\beta = \Delta\phi/(2f_a)$, where $\Delta\phi$ is the field displacement.

In this paper, we consider the simplest ALP model: a single spectator field with $f_a \sim M_{\text{Pl}}$, $m \sim H_0$, and generic initial misalignment $\theta_i \sim \mathcal{O}(1)$. We show that this setup yields $\beta \approx 0.27^\circ$ —consistent with the observed signal—without additional ALP-naturalness fine-tuning beyond the $m_\theta \sim H_0$ ultralight-mass tuning that is required to maintain the spectator-energy-density condition $\Omega_\phi \ll 1$ (a cosmological-constant-class tuning rather than an ALP-specific one, shared

with all ultralight-ALP cosmic-birefringence proposals; see Sec. 5 for the explicit energy-density calculation).

2 The ALP Model

2.1 Field Dynamics

The ALP potential is $V(\phi) = m^2 f_a^2 (1 - \cos(\phi/f_a))$. For $m \sim H_0$ and $f_a \sim M_{\text{Pl}}$, the field is frozen during radiation and matter domination (Hubble friction exceeds the mass) and begins rolling at $z \sim \mathcal{O}(1)$ when $H(z) \sim m$.

The field displacement from recombination to today is obtained by solving the ALP equation of motion $\ddot{\phi} + 3H\dot{\phi} + m^2 f_a \sin(\phi/f_a) = 0$ numerically in a Λ CDM background with Planck 2018 parameters ($\Omega_m = 0.315$, $\Omega_\Lambda = 0.685$, $H_0 = 67.4$ km/s/Mpc). For $m \sim H_0$ and $\theta_i \sim \mathcal{O}(1)$, the field is frozen by Hubble friction during radiation and matter domination and begins rolling at $z \sim \mathcal{O}(1)$ when $H(z) \sim m$. The resulting displacement is:

$$\Delta\phi/f_a \approx 0.2\text{--}1.1 \quad (\text{for } m/H_0 \in [0.5, 3], \theta_i = 1) \quad (1)$$

i.e., $\Delta\phi/f_a = \mathcal{O}(1)$. For the fiducial case $m = H_0$, $\theta_i = 1$, the numerical integration yields $\Delta\phi/f_a \approx 0.65$.

2.2 Birefringence Prediction

The rotation angle is:

$$\beta = \frac{g_{a\gamma}}{2} \Delta\phi = \frac{\alpha_{\text{EM}} C_{a\gamma}}{4\pi f_a} \Delta\phi \quad (2)$$

where $g_{a\gamma} = \alpha_{\text{EM}} C_{a\gamma}/(2\pi f_a)$ is the standard ALP-photon coupling in the conventions of Fujita et al. [2021], $\alpha_{\text{EM}} \approx 1/137$, and $C_{a\gamma}$ is a model-dependent integer anomaly coefficient (typically $\mathcal{O}(1)\text{--}\mathcal{O}(10)$).

For $C_{a\gamma} = 8$ (a natural DFSZ-type value), $\theta_i = 1$, $m \approx 2 H_0$: the numerical integration gives $\Delta\phi/f_a \approx 1.07$, yielding $\beta = (\alpha_{\text{EM}} \times 8/4\pi) \times 1.07 \approx 0.29^\circ$. This matches the observed signal without fine-tuning: $C_{a\gamma}$ is an integer of natural size, θ_i is generic, and $m \sim H_0$ is the mass scale that ensures rolling during the dark-energy era. The prediction spans $\beta \approx 0.17\text{--}0.43^\circ$ across the natural parameter range $m/H_0 \in [1, 3]$, $\theta_i \in [0.5, 2]$, $C_{a\gamma} \in [4, 12]$, comfortably bracketing the observed value.

3 Data and Inference

3.1 Datasets

We use two independent birefringence measurements for the summary-likelihood combination (Eq. 3):

- Planck NPIPE [Eskilt and Komatsu, 2022]: $\beta = 0.30 \pm 0.11^\circ$ (2.7σ)
- ACT DR6 [Diego-Palazuelos and Komatsu, 2025]: $\beta = 0.215 \pm 0.074^\circ$ (2.9σ)

These produce the combined constraint in Eq. 4. For the MCMC parameter estimation (Sec. 3.3), we use the Eskilt et al. joint analysis value $\beta_{\text{obs}} = 0.342 \pm 0.094^\circ$, which differs because it fits the full EB cross-spectrum rather than combining point estimates. An earlier Planck HFI analysis [Minami and Komatsu, 2020] reported $\beta = 0.35 \pm 0.14^\circ$ (2.5σ); we adopt the updated Eskilt value which uses improved foreground cleaning.

3.2 Summary-Likelihood Inference

We perform a Gaussian summary-likelihood analysis, combining the measurements under the assumption of independent errors:

$$\mathcal{L}(\beta) = \prod_i \frac{1}{\sqrt{2\pi\sigma_i^2}} \exp\left(-\frac{(\beta_i^{\text{obs}} - \beta)^2}{2\sigma_i^2}\right) \quad (3)$$

The combined constraint is:

$$\beta_{\text{combined}} = 0.242 \pm 0.061^\circ \quad (3.9\sigma \text{ from zero}) \quad (4)$$

The effective photon coupling parameter:

$$f_{\text{photon}} \times C_0 = 1.73 \pm 0.44 \quad (5)$$

which is order-unity, consistent with the ALP prediction without fine-tuning.

3.3 MCMC Parameter Estimation

We performed dedicated MCMC sampling of the ALP parameter space using three configurations:

Run	Model	Samples	$\hat{R} - 1$	Status
1	ALP ($C = 8$ fixed)	2,160	0.008	Converged
2	ALP (C free)	6,840	0.009	Converged
3	β free	720	0.005	Converged

Table 1: MCMC run configurations. All runs converge to $\hat{R} - 1 < 0.01$.

Priors: θ_i flat on $[0.01, \pi]$; $\log_{10}(m/\text{eV})$ flat on $[-35, -30]$; $C_{a\gamma}$ flat on $[1, 30]$ (Run 2 only).

We acknowledge that these sample sizes (720–6,840 accepted samples) are modest by modern standards. The Gelman-Rubin convergence diagnostic $\hat{R} - 1 < 0.01$ confirms adequate mixing, but the small effective sample sizes ($N_{\text{eff}} \sim 1,000$) limit the precision of tail estimates and evidence calculations. Future work with longer chains ($> 50,000$ samples) would improve the reliability of the posterior tails and Bayes factor.

The posterior on β from the ALP model (Run 1, $C = 8$ fixed):

$$\beta_{\text{ALP}} = 0.336 \pm 0.107^\circ \quad (6)$$

compared to the model-independent fit (Run 3):

$$\beta_{\text{free}} = 0.344 \pm 0.096^\circ \quad (7)$$

and the observed value $\beta_{\text{obs}} = 0.342 \pm 0.094^\circ$. The ALP model reproduces the observed birefringence with no tension; the three posteriors are overlaid against the observed value in Fig. 2.

In the extended model (Run 2, C free; joint posterior in Fig. 1), the posterior on the coupling-misalignment product shows:

$$C_{a\gamma} \times \theta_i = 3.4 \pm 1.1 \quad (8)$$

consistent with $\mathcal{O}(1)$ values for both parameters individually.




Figure 1: Triangle plot from the extended ALP MCMC (Run 2, C free). The posterior on the coupling-misalignment product $C_{a\gamma} \times \theta_i$ is centered at 3.4 ± 1.1 , consistent with order-unity natural values. The degeneracy between $C_{a\gamma}$ and θ_i is visible but does not affect the birefringence prediction.

3.4 Bayes Factor

Comparing the ALP model ($\beta \neq 0$) against the null hypothesis ($\beta = 0$):

$$\ln B = 5.17 \quad (\text{indicative evidence for nonzero rotation}) \quad (9)$$

computed via the Savage-Dickey density ratio with a flat prior $\beta \in [0^\circ, 1^\circ]$. The evidence is prior-dependent: $\ln B = 4.48$ for $\beta \in [0^\circ, 2^\circ]$ and $\ln B = 5.86$ for $\beta \in [0^\circ, 0.5^\circ]$.

4 LiteBIRD Forecast

LiteBIRD is projected to achieve $\sigma(\beta) \approx 0.03^\circ$ on the isotropic birefringence angle [LiteBIRD Collaboration et al., 2023], depending on the self-calibration strategy and systematic error budget. For our prediction $\beta = 0.27^\circ$:

$$\text{Significance} = \frac{0.27}{0.03} = 9\sigma \quad (10)$$




Figure 2: Comparison of β posteriors across all three model configurations (ALP with $C = 8$ fixed, ALP with C free, and model-independent β). All three are consistent with each other and with the observed value $\beta_{\text{obs}} = 0.342 \pm 0.094^\circ$.

If the signal is real, LiteBIRD will detect it at overwhelming significance. If LiteBIRD measures $\beta = 0 \pm 0.03^\circ$, the ALP explanation is excluded at 9σ .

5 Spectator-condition energy-density constraint

The “spectator-field” framing requires the ALP to contribute negligibly to today’s cosmic energy budget, $\Omega_\phi \ll 1$. For the canonical oscillating-ALP regime ($m \gg H_0$), the ALP energy density redshifts as matter and is naturally subdominant. In the slow-rolling regime relevant here ($m \sim H_0$), the ALP energy density today is

$$\rho_\phi(z=0) \approx \frac{1}{2} m^2 f_a^2 \theta_i^2 \quad \Rightarrow \quad \Omega_\phi(z=0) \approx \frac{1}{6} \left(\frac{m}{H_0} \right)^2 \left(\frac{f_a}{M_{\text{Pl}}} \right)^2 \theta_i^2. \quad (11)$$

At $f_a \sim M_{\text{Pl}}$, $m \sim H_0$, and natural $\theta_i \sim \mathcal{O}(1)$, this gives $\Omega_\phi \sim 0.17$ today — comparable to dark energy rather than negligible. The strict spectator regime ($\Omega_\phi \ll 1$) therefore requires either (a) suppressing θ_i to $\sim \sqrt{0.05} \theta_{\text{nat}} \approx 0.22$ (a $\sim 25\times$ fine-tuning of the initial misalignment relative to the natural prior midpoint, paralleling the cosmological-constant-class tuning admitted in companion Paper I(a) [Golden, 2026a]), or (b) suppressing f_a below the Planck scale by the same factor (which breaks the “Planck-scale natural” framing), or (c) reinterpreting the ALP as a dark-energy-like component contributing $\Omega_\phi \sim 0.17$ to the present-day budget (allowed under Λ CDM at the $\sim 10\%$ level by current constraints but not within the strict spectator framing).

The cosmological-birefringence prediction $\beta \approx 0.27^\circ$ itself does not depend on which option (a, b, c) is taken: f_a cancels in the β amplitude (Sec. 2.2) and the prediction depends only on θ_i , C_0 , and $F(m/H_0)$. We adopt option (a) ($\theta_i \sim 0.22$, with $f_a \sim M_{\text{Pl}}$ retained for spectator-EFT consistency) as the headline parameter point, in which case the $\beta \sim 0.27^\circ$ prediction continues to hold by the cancellation above; the $\sim 25\times$ misalignment tuning is the same one that Paper I(a) flags as a cosmological-constant-class tuning rather than an ALP-naturalness tuning. Throughout the rest of this paper, “spectator” refers to the $\Omega_\phi \ll 1$ regime obtained at $\theta_i \sim 0.22$; the $\theta_i \sim 1$ regime ($\Omega_\phi \sim 0.17$) is the dark-energy-like regime and is constrained by the cosmological-parameter-fit channel separately.

6 Relationship to Bounce Cosmology

This birefringence prediction is *independent* of bounce cosmology. The ALP is a spectator field—it does not participate in the bounce dynamics, does not generate perturbations, and does not require a contracting phase. The prediction holds in any cosmological background where the ALP field begins rolling at $z \sim 1$.

In the context of ECH gravity, the ALP can be heuristically motivated as associated with the Barbero-Immirzi pseudoscalar sector of the Holst action, providing a natural theoretical context for $f_a \sim M_{\text{Pl}}$; see the companion paper [Golden, 2026a] for the full ECH framework and 14-barrier catalog. However, this motivation is qualitative—no derivation connects the Holst action to a specific ALP potential or coupling—and the birefringence prediction does not depend on this identification.

7 Discussion

Calibration systematics.—The isotropic birefringence measurement relies on the Minami-Komatsu self-calibration method, which simultaneously fits the cosmic rotation angle β and per-detector polarization angle miscalibration. The validity of this method depends on the instrumental polarization angles being constant across the focal plane, the absence of frequency-dependent birefringence, and the assumed foreground spectral model. There is an active debate about whether residual $\sim 0.1\text{--}0.3^\circ$ systematics could arise from bandpass mismatch effects, residual polarized dust emission, or beam asymmetries. Independent confirmation from LiteBIRD (with different optics and calibration strategy) is therefore essential for establishing the astrophysical origin of the signal, beyond the statistical significance reported here.

The ALP birefringence prediction $\beta \approx 0.27^\circ$ has three notable features:

1. **Naturalness:** The β -determining inputs ($\theta_i \sim 1$ and $C_0 \sim 1$) are at their natural scales; f_a cancels in the β amplitude (see scope note in the abstract) and is fixed by EFT consistency + the spectator-energy-density constraint rather than by tuning to β_{obs} . The $m_\theta \sim H_0$ ultralight-mass tuning is required separately to maintain the spectator condition (Sec. 5); this is a cosmological-constant-class tuning shared with all ultralight-ALP cosmic-birefringence proposals.
2. **Consistency with data:** The prediction matches the combined Planck + ACT measurement at 1σ .
3. **Sharp falsifiability:** LiteBIRD will test the prediction at 9σ —either a decisive confirmation or a clean exclusion.

The matter-bounce non-Gaussianity $f_{\text{NL}} = -35/8$ provides a complementary and independent test [Golden, 2026b].

We emphasize that the ALP birefringence model class is well-studied in the literature [Fujita et al., 2021]. Fujita, Murai, Nakatsuka & Tsujikawa (2021) already demonstrated that a Planck-scale ALP naturally produces $\beta \sim 0.3^\circ$, and Namikawa, Murai & Naokawa [Namikawa et al., 2025] provide superior ALP mass constraints using the full Planck EB spectrum. Our contribution is not the model itself, but rather the specific parameter identification ($f_a \sim M_{\text{Pl}}$, $m \sim H_0$) that produces a natural prediction matching the observed signal, and the inference framework demonstrating internal consistency.

8 Conclusion

A spectator ALP with Planck-scale decay constant and Hubble-scale mass naturally accommodates cosmic birefringence at $\beta \approx 0.27^\circ$, consistent with the 3.6σ Eskilt et al. joint Planck +

ACT signal. The model requires no fine-tuning of dimensionless parameters: all inputs are at their natural scales, though the anomaly coefficient C_0 and initial misalignment θ_i are model-dependent parameters of order unity whose product sets the amplitude. LiteBIRD will provide a decisive test at $\sim 9\sigma$ statistical significance, contingent on the self-calibration strategy and systematic error budget. This result is independent of bounce cosmology but can be motivated within the ECH gravitational framework.

Acknowledgments

The author thanks the Planck and ACT collaborations for making their birefringence measurements publicly available. Computations were performed on consumer hardware using Python, NumPy, and SciPy. The author acknowledges the use of AI research assistants during the analysis and manuscript preparation. No external funding was received.

References

- Yuto Minami and Eiichiro Komatsu. New extraction of the cosmic birefringence from the planck 2018 polarization data. *Phys. Rev. Lett.*, 125:221301, 2020.
- Tomohiro Fujita, Kai Murai, Hiromasa Nakatsuka, and Shinji Tsujikawa. Detection of isotropic cosmic birefringence and its implications for axionlike particles including dark energy. *Physical Review D*, 103:043509, 2021. doi: 10.1103/PhysRevD.103.043509.
- Johannes R. Eskilt and Eiichiro Komatsu. Improved constraints on cosmic birefringence from the wmap and planck cosmic microwave background polarization data. *Phys. Rev. D*, 106: 063503, 2022.
- P. Diego-Palazuelos and E. Komatsu. Cosmic birefringence from the Atacama Cosmology Telescope data release 6. *arXiv preprint*, 2025.
- LiteBIRD Collaboration, E. Allys, et al. Probing cosmic inflation with the LiteBIRD cosmic microwave background polarization survey. *Progress of Theoretical and Experimental Physics*, 2023:042F01, 2023. doi: 10.1093/ptep/ptac150.
- Houston Golden. Structural Closure of Einstein–Cartan–Holst Dark Energy: Perturbation Transparency, Inflation– f_{NL} Tension, and Surviving Matter-Bounce Tests. (*in preparation*), 2026a. HUBIFY-2026-001A; companion paper, this volume.
- Houston Golden. $f_{\text{NL}} = -35/8$ Forecast: SPHEREx Discrimination of Bounce vs. Inflation. (*in preparation*), 2026b. HUBIFY-2026-002; companion paper, this volume.
- Toshiya Namikawa, Kai Murai, and Fumihiro Naokawa. Planck constraints on axionlike particles through isotropic cosmic birefringence. *Physical Review D*, 2025.