

Spin-Torsion Cosmology and the Search for Geometric Dark Energy: Structural Barriers, Perturbation Transparency, and Surviving Predictions

Houston Golden^{1,*}

¹*Independent Researcher, Los Angeles, California, USA*

(Dated: April 27, 2026 — v2.3.0)

We investigate whether Einstein-Cartan-Holst (ECH) spin-torsion gravity can produce late-time dark energy or distinctive cosmological signatures. Within this framework, LQC holonomy corrections yield a nonsingular quantum bounce at $\rho_{\text{crit}} \approx 0.27\text{--}0.41 \rho_{\text{Pl}}$ (depending on the entropy-counting scheme for the Barbero-Immirzi parameter), while the ECH parity structure generates a unique four-fermion interaction. Through systematic analysis of 7 foundation studies and 17 research branches, we establish 14 independent structural barriers that close all minimal routes from the bounce to dark energy. Our central theoretical result is a perturbation-transparency result: for canonical scalar field matter, torsion vanishes at all perturbation orders, the Holst term reduces to a topological invariant, and the Barbero-Immirzi parameter γ is invisible in all perturbation observables—generalizing the well-known observation that Einstein-Cartan theory with spinless matter reduces to GR [1]. Independent MCMC verification (Cobaya v3.6.1, 424,181 samples across three dataset combinations, two frozen) finds ΔN_{eff} consistent with zero (-0.020 ± 0.169 full-tension; $+0.065 \pm 0.17$ Planck+BAO+SN) and $H_0 = 67.68 \pm 1.06 \text{ km s}^{-1} \text{ Mpc}^{-1}$, recovering standard Λ CDM values. The conclusion: ECH provides a clean bounce mechanism but cannot produce late-time dark energy or distinctive perturbation-level observables through any standard mechanism tested. The surviving science case rests on predictions of the matter bounce scenario [2], which is compatible with (but not derived from) the ECH framework—principally $f_{\text{NL}} = -35/8$ (following the convention of Cai *et al.* [2]; the alternate convention $-35/16$ of Li *et al.* [3] gives an identical bispectrum up to permutation counting), testable by SPHEREx at $4\text{--}6\sigma$.

PACS numbers: 98.80.-k, 04.50.Kd, 04.60.Pp, 95.36.+x

CONTENTS

		V. Data Methods: Galaxy Spin Analysis	9
I. Introduction	2	VI. Data Methods: CMB E - B Analysis	9
A. Theoretical Foundations and Novel Synthesis	3	VII. Cosmological Fits and Model Comparison	10
B. Original Contributions	3	A. Datasets and Configuration	10
C. Paper Organization	3	B. Results	10
II. Theoretical Framework	4	VIII. Systematic Analysis	10
A. Loop Quantum Cosmology and the Holst Action	4	IX. Falsification Criteria	10
1. Einstein-Cartan-Holst Action	4	X. Related Work	11
2. Derivation of the Parity-Odd Term	4		
3. Parameter Naturalness	6	XI. Structural Barriers to First-Principles Dark Energy	11
B. Black Hole Interior and Quantum Bounce	6	A. Barrier 1: Mass-Coupling Lock (Foundation A)	11
C. Cosmic Rotation and Dark Energy	6	B. Barrier 2: Topological-Shift Duality (Foundation B)	12
1. Inflationary Suppression	7	C. Barrier 3: Scalar-Tensor Universality (Foundation C)	12
2. Galaxy Spin Alignment Mechanism	7	D. Barrier 4: Planck Suppression (Foundation D)	12
III. Observational Signatures and Evidence	7	E. Barrier 5: Scale Separation (Foundation E)	12
A. CMB E - B Cross-Correlations	7	F. Barrier 6: Attractor-Sensitivity Dilemma (Foundation F)	12
B. Galaxy Spin Asymmetry: A Contested Anomaly	8	G. Barrier 7: Parameter Immunity (Foundation G)	13
C. Cosmological Tensions: H_0 and σ_8	8		
D. Independent Verification Results	8		
IV. Enhanced Theoretical Derivations	9		

* houston@hubify.com

H. Barrier 8: Parity-Even Interaction (Branch H)	13
I. Barrier 9: Liouville Conservation (Branch J)	13
J. Barrier 10: UV→IR Specificity Dilemma (Branch L)	13
K. Barrier 11: Decoupling Universality (Branches L/M)	14
L. Barrier 12: Vacuum Amplification Ceiling (Branch M)	14
M. Barrier 13: Gravitational Democracy (Branches N/O)	14
N. Barrier 14: Perturbation Transparency	14
XII. The Perturbation-Transparency Result	14
A. Statement	14
B. Proof (Scalar Sector)	15
C. Extension to Tensor Sector	15
D. Explicit Verification: The Holst Term in Perturbation Theory	15
E. What Would Break the Transparency	15
F. Implications	16
G. Discrimination Among Bouncing Cosmologies	16
XIII. The Hybrid Dark-Energy Loophole	17
XIV. Discussion	17
A. The Inflationary Suppression Factor	17
B. Theoretical Implications	17
C. Cosmic Birefringence: Spectator ALP Consistency Check	17
XV. Limitations and Future Directions	18
A. Current Limitations	18
1. Theoretical	18
2. Observational	18
B. Robustness to Galaxy Spin Null Results	19
C. Future Observational Prospects	19
D. Theoretical Research Program	20
1. Higher-Loop and Non-Perturbative Verification	20
2. Galaxy Spin Dipole: Resolved	20
3. Bounce-to-Inflation Transition Dynamics	21
4. Parity-Odd Primordial Gravitational Waves	21
5. Black Hole Interior Numerical Relativity	21
6. Connection to Other Quantum Gravity Approaches	21
E. Structural Tension: Dark Energy vs. Bounce f_{NL}	21
F. Structural Closure	22
G. Open Questions	22
XVI. Conclusions	22
Data and Code Availability	24

Acknowledgments	24
A. Complete Parameter Summary	25
B. Dimensional Analysis	25
C. Reproducibility Materials	25
D. Claims Classification	26
References	26

I. INTRODUCTION

The nature of dark energy remains one of the most profound challenges in modern physics. While the Λ CDM model successfully accounts for the observed cosmic acceleration [4], it faces severe theoretical difficulties—most notably the cosmological constant problem, where the observed vacuum energy density $\rho_\Lambda \sim (2.3 \text{ meV})^4$ is $\sim 10^{120}$ times smaller than the naive quantum field theory estimate $\sim M_{\text{Pl}}^4$ [5]. This spectacular fine-tuning has motivated extensive theoretical work on dynamical dark energy, modified gravity, and quantum gravitational approaches.

Simultaneously, precision cosmological measurements have revealed growing tensions within Λ CDM. The Hubble constant measured from the early universe via the cosmic microwave background (CMB) by Planck ($H_0 = 67.36 \pm 0.54 \text{ km s}^{-1} \text{ Mpc}^{-1}$) [4] disagrees at $\sim 4.9\sigma$ with late-universe determinations from the SH0ES collaboration ($H_0 = 73.04 \pm 1.04 \text{ km s}^{-1} \text{ Mpc}^{-1}$) [6].¹ The matter clustering amplitude σ_8 inferred from Planck ($\sigma_8 = 0.811 \pm 0.006$) exceeds weak lensing measurements from KiDS-1000 ($S_8 = 0.759^{+0.024}_{-0.021}$) [7] and DES Y3 [8] by $2\text{--}3\sigma$. The DESI 2024 BAO results [9] suggest dynamical dark energy at $2.5\text{--}3.9\sigma$, with the subsequent DESI DR2 analysis [10] strengthening this preference to 3.1σ (BAO+CMB) and up to 4.2σ (with supernovae), adding urgency to the search for extensions of the standard model.

This work presents a phenomenological framework that addresses these challenges through quantum gravitational effects in spin-torsion cosmology. Our approach builds on three well-established theoretical pillars, synthesized here for the first time into a unified cosmological model:

1. *Loop Quantum Cosmology* (LQC), providing a non-singular quantum bounce replacing the classical Big Bang singularity [11]. The bounce occurs at a critical density $\rho_{\text{crit}} \simeq 0.27\text{--}0.41 \rho_{\text{Pl}}$ (depending on the entropy-counting scheme for the Barbero-Immirzi parameter; see Sec. II B).

¹ $(73.04 - 67.36)/\sqrt{1.04^2 + 0.54^2} = 4.86\sigma$; we quote $\sim 4.9\sigma$ throughout.

2. *Einstein-Cartan theory* incorporating fermionic spin-torsion coupling, which generates four-fermion contact interactions and prevents gravitational singularities through torsion-induced repulsion at extreme densities [1, 12].
3. *Black hole universe origin*, where a rotating parent black hole spawns a non-singular baby universe beyond its event horizon through torsion-regulated gravitational collapse [13]. Torsion as an alternative to cosmic inflation was explored in Ref. [14]. The baby universe inherits angular momentum, establishing a preferred cosmic axis.

A. Theoretical Foundations and Novel Synthesis

Our framework synthesizes well-established theoretical components into a unified cosmological model with testable outputs:

Einstein-Cartan Theory.—The inclusion of spacetime torsion coupled to fermionic spin has been extensively studied since the pioneering work of Hehl *et al.* [1]. Poplawski [12, 15] demonstrated that torsion-induced four-fermion interactions can generate a small positive cosmological constant through condensate formation, and that the parity-violating pseudoscalar density modifies this interaction, providing a candidate mechanism for dark energy generation. The Einstein-Cartan-Sciama-Kibble (ECSK) framework avoids singularities through torsion-induced repulsion at extreme densities. Recent work by Liu *et al.* [16] showed that Einstein-Cartan torsion is preferred over Λ CDM by AIC criteria, providing independent statistical support for torsion-based cosmology.

Loop Quantum Gravity and Parity Violation.—The connection between LQG’s Barbero-Immirzi parameter and parity-violating effects has been rigorously established by Freidel, Minic & Takeuchi [17] and Mercuri [18, 19]. They showed that the Holst term in LQG, when coupled to fermions, generates effective four-fermion axial interactions that violate parity. The Barbero-Immirzi parameter γ , originally introduced as an ambiguity in the LQG quantization, acquires physical significance through its role in parity-violating observables.

Black Hole Universe Origin.—Poplawski [13] pioneered the “universe in a black hole” scenario, demonstrating that every black hole with torsion spawns a non-singular, closed baby universe beyond its event horizon. His subsequent work [14] explored torsion as an alternative to cosmic inflation, with the baby universe inheriting angular momentum and developing a preferred cosmic axis, providing a candidate origin for observed cosmic anomalies.

B. Original Contributions

Each theoretical ingredient above has been studied independently. Our contribution is to assemble them into a single quantitative framework and test it systematically. The original contributions are:

1. *14-barrier catalog and perturbation-transparency result:* Through 7 foundation studies and 17 research branches, we establish 14 independent structural barriers closing all minimal routes from the bounce to dark energy. The central result is that minimal ECH gravity is perturbation-transparent: for canonical scalar field matter, torsion vanishes at all perturbation orders and γ is invisible in all observables.
2. *MCMC verification revealing null result:* Independent Cobaya verification across two frozen dataset combinations demonstrates that ΔN_{eff} is consistent with zero, H_0 recovers the standard Planck Λ CDM value of 67.68 ± 1.06 , and the spin-torsion extension alone does not resolve cosmological tensions.
3. *ALP birefringence consistency check (secondary):* A spectator ALP with natural parameters ($f_a \sim M_{\text{Pl}}$, $m \sim H_0$) accommodates the observed $\beta = 0.242^\circ \pm 0.061^\circ$ (3.9σ) without fine-tuning. This ALP model class was previously studied by Fujita *et al.* [20]; our contribution is the ECH-motivated parameter identification and independent MCMC inference, not the model itself. The result is a consistency check, not a prediction of minimal ECH (which does not derive the photon-torsion coupling; see Sec. XIV C).

C. Paper Organization

This paper is organized as follows. Section II develops the theoretical framework. Section III presents observational signatures and evidence, including the independent MCMC verification (Sec. IIID). Sections IV–VI summarize derivations and data methods (extended details available upon request from the author). Section VII presents cosmological fits and model comparison. Sections VIII–X provide condensed systematic analysis, falsification criteria, and related work. The core results occupy Secs. XI–XIII: the 14 structural barriers, perturbation-transparency result, and hybrid loophole rejection. Section XIV discusses the inflationary suppression factor and birefringence analysis. Sections XV–XVI address limitations and conclusions. Appendices provide the parameter summary (Appendix A), dimensional analysis (Appendix B), reproducibility materials (Appendix C), and claims classification (Appendix D).

TABLE I. Executive summary: what the investigation found. Most routes fail; the surviving testable prediction is the matter-bounce $f_{\text{NL}} = -35/8$ [2].

Question	Result	Status
Does the framework resolve H_0 tension?	$H_0 = 67.68 \pm 1.06$ km/s/Mpc ^a	No. Standard Λ CDM recovered.
Does the framework resolve σ_8/S_8 tension?	$\sigma_8 = 0.803$; $S_8 = 0.814$ ^b	No. Below Planck at $\sim 2\sigma$.
Can the bounce derive dark energy?	14 barriers close all routes	No. Scaling ansatz only ^c .
Is there a nonsingular bounce?	LQC: $\rho_c \simeq 0.27\text{--}0.41 \rho_{\text{Pl}}$	Yes (LQC holonomy correction, not EC)
Is ECH visible in perturbations?	Perturbation-transparency result	No. γ invisible at all orders.
Is there a testable prediction?	$f_{\text{NL}} = -35/8$ (SPHEREx $\sim 5.5\sigma$, template-corrected)	Yes (matter bounce, mechanism-indepe

^a $(67.68 - 73.04)/\sqrt{1.06^2 + 1.04^2} = 3.61\sigma$; the ΔN_{eff} extension does not by itself resolve the Hubble tension (the original lower H_0 values were SH0ES-prior-driven; independent Cobaya v3.6.1 also recovers $H_0 = 67.79 \pm 1.09$ on Planck+BAO+SN — see Sec. IIID). ^b $\sigma_8 = 0.803 \pm 0.008$, $S_8 = 0.814 \pm 0.008$ (full-tension), in mild tension with Planck ($S_8 = 0.832$, $(0.832 - 0.814)/0.008 = 2.25\sigma$; $\sigma_8 = 0.811$, 1.0σ); the ΔN_{eff} extension does not reduce the σ_8/S_8 tension. ^cReparameterized as sensitivity to N_{tot} , not solved; see Sec. XIV A.

II. THEORETICAL FRAMEWORK

A. Loop Quantum Cosmology and the Holst Action

1. Einstein-Cartan-Holst Action

The fundamental action combining Einstein-Cartan theory with the Holst term is

$$S_{\text{ECH}} = \frac{1}{16\pi G} \int d^4x e \left[e_a^\mu e_b^\nu R_{\mu\nu}^{ab} + \frac{1}{\gamma} \varepsilon^{abcd} e_a^\mu e_b^\nu R_{cd\mu\nu} + \frac{1}{4} T^{abc} T_{abc} \right] + S_{\text{matter}}, \quad (1)$$

where $e = \det(e_\mu^a)$ is the tetrad determinant, $R_{\mu\nu}^{ab}$ is the curvature of the Lorentz connection, γ is the Barbero-Immirzi parameter, and T^{abc} is the torsion tensor. The Holst term $\varepsilon^{abcd} e_a^\mu e_b^\nu R_{cd\mu\nu} / \gamma$ is topological in the absence of torsion but contributes non-trivially when fermions are present. This construction builds directly on the foundational work of Freidel, Minic & Takeuchi [17], who established that the Barbero-Immirzi parameter becomes physically observable through its coupling to fermionic matter.

Equation (1) presents the *second-order* (metric) form of the ECH action, obtained after algebraic elimination of the torsion tensor via its own equation of motion (Eq. 3 below). In this form, the $\frac{1}{4} T^{abc} T_{abc}$ term is a shorthand for the four-fermion contact interaction that results from integrating out the non-propagating torsion (see, e.g., Ref. [1]). The coefficient $\frac{1}{4}$ is fixed by the Einstein-Cartan algebraic constraint $T_{bc}^a = \kappa \epsilon^a_{bcd} \bar{\psi} \gamma^d \gamma^5 \psi$; it is not a free parameter. We do not vary the action with respect to torsion as an independent field—the variation is performed with respect to the metric $g_{\mu\nu}$ and the matter fields only. In the first-order (Palatini) formulation, torsion is determined algebraically by the spin density, and this term is not independently specified. The results of this paper depend only on the standard EC torsion equation of motion, not on an independent kinetic term for torsion.

The Barbero-Immirzi parameter is fixed by the LQG black hole entropy calculation [11, 21]:

$$\gamma = 0.274 \pm 0.020, \quad (2)$$

Different entropy-counting schemes yield different values: the Ashtekar-Baez-Corichi-Krasnov (ABCK) calculation [21] gives $\gamma_{\text{ABCK}} \approx 0.274$, while the Domagala-Lewandowski-Meissner (DLM) full $\text{SU}(2)$ state counting [22, 23] gives $\gamma_{\text{DLM}} \approx 0.2375$. We adopt $\gamma = 0.274$ (ABCK) throughout this paper; the DLM value would shift ρ_{crit} upward to $\simeq 0.41 \rho_{\text{Pl}}$ without qualitatively changing any conclusion. Whether γ constitutes a genuine physical observable or a quantization artifact absorbable by canonical transformations remains debated [17, 18]. This distinction does not affect our results at leading order: if γ is non-physical, the torsion-fermion coupling factor $\gamma^2/(\gamma^2 + 1)$ in Eq. (4) reverts to unity (the standard ECSK value), modifying the parity-odd coefficient by $\mathcal{O}(10\%)$ without eliminating parity violation, which originates from the Holst term's coupling to fermions independently of the Barbero-Immirzi interpretation.

2. Derivation of the Parity-Odd Term

Starting with the complete action $S = S_{\text{gravity}} + S_{\text{Holst}} + S_{\text{fermion}}$, we derive the parity-odd effective action through four steps:

Step 1: Torsion Activation.—When fermions are minimally coupled to gravity with torsion, the torsion tensor is determined algebraically by the fermionic spin density:

$$T^{abc} = 8\pi G S^{abc}, \quad (3)$$

where $S^{abc} = \frac{1}{4} \bar{\psi} \gamma^{[a} \gamma^{bc]} \psi$ is the fermionic spin density tensor (we use natural units $c = \hbar = 1$ throughout; see Appendix XVI). Unlike in general relativity, torsion is not dynamical but is determined instantaneously by the matter content.




FIG. 1. Energy density hierarchy from the Planck scale to the observed dark energy scale. The Einstein-Cartan-Holst action generates a parity-odd vacuum energy $\rho_{\text{vac}} \sim [(\alpha/M) M_{\text{Pl}}] M_{\text{Pl}}^4$ at one loop. Inflationary dilution by $\sim e^{-3N_{\text{tot}}}$ bridges the gap to the observed $\rho_{\Lambda} \approx (2.3 \text{ meV})^4$. Left axis: comparison of the residual fine-tuning in this framework (10^5 , from uncertainty in N_{tot}) versus $\Lambda_{\text{CDM}} (10^{120})$.

Step 2: Four-Fermion Contact Interaction.—Substituting Eq. (3) back into the action and integrating out torsion yields an effective four-fermion interaction [1]:

$$\mathcal{L}_{\text{int}} = -\frac{3\pi G_N}{2} \times \frac{\gamma^2}{\gamma^2 + 1} \times J_\mu^5 J^{5\mu}, \quad (4)$$

where $J^{5\mu} = \bar{\psi} \gamma^\mu \gamma^5 \psi$ is the axial current. The Barbero-Immirzi parameter enters through the $\gamma^2/(\gamma^2 + 1)$ factor, which is purely a consequence of the Holst modification and would be absent ($\rightarrow 1$) in standard Einstein-Cartan theory.

Step 3: Parity-Odd Effective Action.—The effective action acquires a parity-odd term through quantum corrections, as shown by Mercuri [19]. The origin of this term is the Holst term: in first-order variables (tetrad e^I and Lorentz connection ω^{IJ} , with $I, J, \dots$ internal Lorentz indices), the ECH action's parity-odd piece is

$$S_{\text{Holst}} = \frac{M_{\text{Pl}}^2}{\gamma} \int e_I \wedge e_J \wedge R^{IJ}(\omega), \quad (5)$$

where $R^{IJ} = d\omega^{IJ} + \omega^I_K \wedge \omega^{KJ}$ is the curvature 2-form. This is a manifestly diffeomorphism-invariant 4-form (two tetrad 1-forms $\wedge$ one curvature 2-form). When the connection is decomposed as $\omega^{IJ} = \tilde{\omega}^{IJ} + K^{IJ}$ (Levi-Civita + contorsion), Eq. (5) generates terms involving contorsion coupled to curvature. After integrating out torsion (which is algebraically determined by fermion

spin density, Eq. 3) and including quantum corrections at one loop, the effective parity-odd action takes the form

$$S_{\text{eff}} = \frac{\alpha}{M} \int e_I \wedge e_J \wedge \mathcal{F}^{IJ}[K, \tilde{R}], \quad (6)$$

where $\mathcal{F}^{IJ}$ collects the contorsion-dependent pieces ($\tilde{D}K^{IJ} + K^I_K \wedge K^{KJ}$ at leading order), $M = M_{\text{area-gap}} \sim \sqrt{\gamma} M_{\text{Pl}}$ is the LQG area-gap mass scale, and α is a *dimensionless* coupling ($[\alpha] = M^0$). In components, the leading contribution reduces to

$$S_{\text{eff}} = \int d^4x \sqrt{-g} \frac{\alpha}{M} \varepsilon^{\mu\nu\rho\sigma} e_\mu^I e_\nu^J \mathcal{F}_{IJ\rho\sigma}, \quad (7)$$

where the operator $\varepsilon^{\mu\nu\rho\sigma} e_\mu^I e_\nu^J \mathcal{F}_{IJ\rho\sigma}$ has mass dimension +2 and α/M has dimension -1, giving a total Lagrangian density of mass dimension +1—three short of the required +4 (see Appendix B for the full counting).

We emphasize that the operator in Eq. (7) has naive mass dimension $[\mathcal{L}_{\text{odd}}] = +1$ in the Lagrangian density, three units short of the required +4. The evaluation at Planck-scale densities ($K \sim M_{\text{Pl}}$, $R \sim M_{\text{Pl}}^2$) that yields $\rho_{\Lambda} = \Xi M_{\text{Pl}}^4$ is therefore a *scaling ansatz*, not a controlled effective field theory calculation. A rigorous EFT treatment would require either (a) identifying the correctly dimensioned operator from the full set of dimension-4 parity-odd invariants, or (b) demonstrating that the on-shell evaluation is protected by a symmetry (e.g., a non-renormalization theorem in the torsionful sector). We proceed with the scaling ansatz as a phenomenological parameterization, noting that the 14 structural barriers identified in Secs. XI–XIII close the bounce-to-dark-energy route independently of whether this dimensional gap can be resolved. Throughout this paper, we adopt the convention that the Einstein-Hilbert action carries the standard $M_{\text{Pl}}^2/2$ prefactor, while the parity-odd sector carries α/M independently.

This operator is parity-odd (it changes sign under spatial reflection) but fully diffeomorphism-covariant. General covariance does not forbid parity-odd terms; it constrains only the tensor structure. Since classical gravity preserves parity, the coefficient α/M must be generated through quantum corrections.

Notation.—For brevity, we sometimes write the component-form integrand as “ $\varepsilon^{abcd} K_{ab} R_{cd}$ ” (where a, b, c, d are internal Lorentz indices) as shorthand for the full 4-form expression Eq. (6). This shorthand suppresses the tetrad factors that make the expression a proper 4-form; all variational calculations use the complete first-order expression.

Torsion remains algebraic.—The effective action S_{eff} (Eq. 6) contains derivatives of the contorsion through $\tilde{D}K^{IJ}$ in $\mathcal{F}^{IJ}$, which could in principle promote torsion from algebraic to dynamical. However, this kinetic contribution is suppressed by $\alpha/M \sim 10^{-21} \text{ GeV}^{-1}$ relative to the original EC torsion equation (sourced by M_{Pl}^2), making the correction $\mathcal{O}(\alpha/M M_{\text{Pl}}^2) \sim 10^{-3}$ to the torsion equation of motion. Torsion therefore remains effectively

algebraic (determined instantaneously by the spin density) at all energy scales below the Planck scale. A full stability analysis (ghost/tachyon checks) of the modified torsion sector is warranted but is not expected to reveal pathologies at this suppression level.

Step 4: Parity-Odd Coefficient.—The existence of a finite parity-odd coefficient is motivated by one-loop estimates in a torsionful background. Following the one-loop analyses of Freidel *et al.* [17] and Shapiro & Teixeira [24], which established the divergence structure and renormalization of the effective Barbero-Immirzi parameter, the expected order of magnitude is

$$\frac{\alpha}{M} \sim \frac{g^2}{32\pi^2} \frac{\gamma}{M} \ln\left(\frac{\Lambda_{\text{UV}}^2}{\mu^2}\right) + \delta_{\text{NY}}, \quad (8)$$

where g is the effective dimensionless axial-torsion Yukawa coupling defined by the vertex $g\bar{\psi}\gamma^5\gamma^a\psi K_a$ (with $g^2 \sim 8\pi G_N M^2 \gamma^2/(\gamma^2 + 1) \sim \mathcal{O}(1)$ at the area-gap scale), Λ_{UV} is a UV scale, μ is the renormalization scale, and δ_{NY} encodes the finite Nieh–Yan contribution whose precise value depends on the regularization scheme—particularly the treatment of γ_5 in dimensional regularization [24] (the ’t Hooft–Veltman and Breitenlohner–Maison prescriptions yield different finite parts for parity-odd traces). We therefore **treat α/M as a phenomenological parameter constrained by data** (see Table VI), with Eq. (8) providing the theoretical motivation for its existence and order of magnitude ($[(\alpha/M) M_{\text{Pl}}] \sim 10^{-2}$), not a controlled first-principles prediction.

A one-loop RG estimate suggests logarithmic running:

$$\frac{d\alpha}{d\ln\mu} \sim -\frac{g^2}{16\pi^2} \times 2N_f, \quad (9)$$

where N_f is the number of fermion species contributing at scale μ . However, this equation inherits the same scheme dependence as the coefficient itself: its derivation requires specifying how γ_5 is handled in $d = 4 - \epsilon$ dimensions and which counterterm conventions are adopted for the Nieh–Yan sector. The physical content is that α runs weakly (the loop factor $g^2/(16\pi^2) \sim 10^{-3}$ ensures perturbative suppression), so its value is approximately scale-independent over cosmological history. A controlled derivation of α/M from non-perturbative quantum gravity (spin-foam amplitudes or lattice methods) remains an important open problem (Sec. XV).

3. Parameter Naturalness

The parent black hole mass must exceed $M_{\text{crit}} = (M_{\text{Pl}}^4/\rho_{\text{vac}})^{1/3} \approx 10^{-3} M_{\odot}$, easily satisfied by any astrophysical black hole. The required dilution of inherited rotation is naturally achieved through ~ 50 e -folds of inflation.

B. Black Hole Interior and Quantum Bounce

In Loop Quantum Cosmology (LQC), holonomy corrections to the gravitational Hamiltonian constraint produce a non-singular bounce at Planck-scale densities [11]. The effective Friedmann equation is²

$$H^2 = \frac{8\pi G}{3} \rho \left[1 - \frac{\rho}{\rho_{\text{crit}}}\right], \quad (10)$$

where the critical density is

$$\rho_{\text{crit}} = \frac{3}{8\pi G \gamma^2 \Delta} = \frac{\sqrt{3}}{32\pi^2 \gamma^3} \rho_{\text{Pl}} \simeq 0.27 \rho_{\text{Pl}}, \quad (11)$$

with $\Delta = 4\sqrt{3}\pi\gamma\ell_P^2$ being the LQG area gap and $\rho_{\text{Pl}} \equiv c^5/(\hbar G^2)$. The numerical value $\rho_{\text{crit}} \simeq 0.27 \rho_{\text{Pl}}$ corresponds to $\gamma = 0.274$ (ABCK); the DLM value $\gamma = 0.2375$ gives $\rho_{\text{crit}} \simeq 0.41 \rho_{\text{Pl}}$ [11]. The range $\gamma \in [0.2375, 0.294]$ brackets both counting schemes and gives $\rho_{\text{crit}} \in [0.22, 0.41] \rho_{\text{Pl}}$. The factor $(1 - \rho/\rho_{\text{crit}})$ ensures that $H^2 \rightarrow 0$ as $\rho \rightarrow \rho_{\text{crit}}$, producing a smooth bounce. Key properties of the bounce:

- For $\rho \ll \rho_{\text{crit}}$: standard GR is recovered exactly.
- For $\rho \rightarrow \rho_{\text{crit}}$: $H \rightarrow 0$ (the universe momentarily stops expanding/contracting).
- For ρ slightly below ρ_{crit} (contracting phase): $\dot{H} > 0$, triggering the bounce.
- The bounce creates a new expanding region with initial conditions completely determined by the parent black hole properties, with no free parameters.

In a rotating black hole, the collapsing matter carries angular momentum through the bounce. The baby universe inherits this rotation, establishing a preferred cosmic axis $\hat{\omega}^a$. This is the physical origin of cosmic parity violation in our framework.

C. Cosmic Rotation and Dark Energy

The effective cosmological constant in our framework is parameterized as:

$$\boxed{\Lambda_{\text{eff}} \equiv \Xi M_{\text{Pl}}^2 + c_{\omega}\omega^2}, \quad \Xi \equiv \left[\frac{\alpha}{M} M_{\text{Pl}}\right] \mathcal{D}_{\text{inf}}, \quad (12)$$

² The LQC bounce (from quantum geometry holonomy corrections) is physically distinct from the Einstein–Cartan torsion bounce of Popławski [12], which produces a qualitatively similar ρ^2 repulsion through classical spin-torsion coupling at high fermion density. In this paper, we adopt the LQC effective equation for its well-studied perturbation dynamics. The ECH framework provides the broader theoretical context for the parity structure (Sec. II A 2) but is not the source of Eq. (10).

where the first term identifies the parity-odd vacuum energy with the cosmological constant (a central *assumption*, not a derivation) and $c_\omega \omega^2$ is the vorticity contribution from the 1 + 3 covariant decomposition [25]. CMB isotropy bounds give $(\omega/H)_0 < 5 \times 10^{-11}$ [26], so $|c_\omega| \omega^2/H_0^2 < 2.5 \times 10^{-21}$ —rotation is completely negligible for background expansion. The dark energy scale is set entirely by $\Xi \sim 10^{-123}$ (Sec. XIV A). The expression $\rho_\Lambda = \Xi M_{\text{Pl}}^4$ is a *parameterization* (two free parameters matching one observable), not a controlled derivation; the “fine-tuning reduction from 10^{120} to 10^5 ” reparameterizes the hierarchy as sensitivity to N_{tot} (Sec. XIV A).

1. Inflationary Suppression

The contorsion K_{ab} , sourced by the fermion spin density, dilutes as a^{-3} during inflation. The dilution factor is:

$$D_{\text{inf}} = \exp[-3N_{\text{tot}}] \times \left(\frac{T_{\text{reh}}}{M_{\text{GUT}}} \right)^{3/2}. \quad (13)$$

Matching $\rho_\Lambda \approx (2.3 \text{ meV})^4$ requires $N_{\text{tot}} \approx 92$ (a fitted parameter, not predicted), reducing fine-tuning from 10^{120} to $\sim 10^5$ as sensitivity to $\Delta N_{\text{tot}} \approx 4$ e -folds. This reparameterizes the fine-tuning problem as an initial-condition question (“why ~ 92 e -folds?”) rather than a fundamental-constant question, but does not solve the cosmological constant problem.

We *assume* $w = -1$ at late times; deriving this requires showing that the early-universe spin-torsion operator generates an IR-constant vacuum term, which has not been achieved. Four minimal routes (NJL condensate, one-loop effective action, dynamical Immirzi field, parity-sensitive CMB phenomenology) all yield negative results [27].

2. Galaxy Spin Alignment Mechanism

The galaxy spin asymmetry amplitude evolves as

$$A(z) = A_0(1+z)^{-p} e^{-qz}, \quad A_0 \in [0, 0.02], \quad (14)$$

with A_0 the present-day amplitude. The parity-odd operator introduces systematic chirality into the tidal field, but the framework’s coupling $\alpha/M \sim 10^{-21} \text{ GeV}^{-1}$ underpredicts any plausible spin asymmetry by > 100 orders of magnitude. Our chirality catalog of 8.47 million galaxies [28] confirms the null: $f_{\text{CW}}^{\text{eq}} = 0.5012 \pm 0.0006^3$ (0.4σ from parity conservation). Galaxy spin asymmetry is not a prediction of the theory, and no significant signal exists to explain.

³ The value $f_{\text{CW}}^{\text{eq}} = 0.5012 \pm 0.0006$ is measured on the benchmark-overlap subset where Galaxy Zoo labels are available for validation; the full-catalog equivariant value is $f_{\text{CW}}^{\text{eq}} = 0.4974 \pm 0.0003$ (see Paper 4 [28] for details). Both are consistent with parity conservation.

III. OBSERVATIONAL SIGNATURES AND EVIDENCE

A. CMB E - B Cross-Correlations

The parity-odd effective action generates CMB polarization signatures through cosmic birefringence. The parity-odd operator rotates the CMB polarization plane by a small angle β . For a spatially uniform (isotropic) rotation, the standard result [29] is

$$C_\ell^{EB} \approx \frac{\sin 4\beta}{2} (C_\ell^{EE} - C_\ell^{BB}) \approx 2\beta (C_\ell^{EE} - C_\ell^{BB}), \quad (15)$$

where the small-angle approximation holds for $\beta \ll 1$ rad. This produces nonzero EB power *across all multipoles*, tracking the $EE - BB$ spectrum shape—not localized to any particular ℓ range. Similarly, the induced TB cross-correlation is $C_\ell^{TB} \approx 2\beta C_\ell^{TE}$.

Connecting to Planck.—The Planck measurement reports $\beta \approx 0.30^\circ$ at 2.4 – 2.7σ [29, 30]. Our framework’s parity-odd operator (Eq. 6) is qualitatively consistent with cosmic birefringence, since it violates parity. However, deriving the photon polarization rotation angle β from our gravitational/torsion operator requires an explicit effective coupling between the parity-odd sector and the electromagnetic field—for example, a term of the standard axion-like form

$$\mathcal{L} \supset \frac{\varphi(\tau)}{4f} F_{\mu\nu} \tilde{F}^{\mu\nu}, \quad (16)$$

where φ is a pseudo-scalar field sourced by the spin-torsion sector and f is the associated decay constant, generated by integrating out torsion at one loop with an electromagnetic vertex. This coupling has not yet been derived in this work. We therefore treat the consistency between our framework’s parity violation and the observed $\beta \approx 0.30^\circ$ as suggestive, not as a quantitative prediction.

Relation between φ/f and β .—For a spatially uniform pseudo-scalar rolling over conformal time, the standard result [31] gives a uniform polarization rotation $\beta = \Delta\varphi/(2f)$, where $\Delta\varphi$ is the net field excursion between last scattering and today. This relation assumes three conditions: (i) spatial uniformity of φ over the last-scattering surface (justified if the correlation length exceeds the Hubble radius at decoupling), (ii) photon propagation along FLRW geodesics (valid for the isotropic component), and (iii) $\varphi/f \ll 1$ so that the rotation is in the small-angle regime (consistent with $\beta \approx 0.30^\circ \approx 5 \times 10^{-3}$ rad). Deriving φ and f from the spin-torsion operator $(\alpha/M)\varepsilon^{abcd}K_{ab}R_{cd}$ requires computing the one-loop photon-torsion vertex and matching to the effective Lagrangian (16); this calculation is beyond the scope of the present work. We use the literature values of β as consistency benchmarks throughout.

Anisotropic component.—In addition to the uniform rotation, the cosmic rotation axis defines a preferred direction $\hat{n}$, which introduces an *anisotropic* birefringence

field $\beta(\hat{n})$ with power concentrated at low multipoles. This anisotropic component contributes primarily to off-diagonal ℓ - ℓ' correlations (detectable via quadratic estimators [30]), with additional diagonal power at $\ell \lesssim 10$. A detailed derivation of the anisotropic component's spectrum from the spin-torsion mechanism is deferred to future work (Sec. XV); here we note that alternative models predict different angular spectra:

- *Axion cosmic birefringence*: Scale-invariant power in $\beta(\hat{n})$, producing EB correlations across all ℓ .
- *Chern-Simons gravity*: Peaks at $\ell \sim 100$ –1000.
- *Spin-torsion (this work)*: Isotropic birefringence across all ℓ (rotation angle not derived from the torsion background alone; requires a photon–torsion coupling such as the spectator-ALP channel of Sec. XIV C), plus an anisotropic low- ℓ component whose amplitude and angular spectrum are not derived in this work and are flagged as an open direction in Sec. XIV C.

Observational support comes from Minami & Komatsu [29], who found cosmic birefringence in Planck data at $\beta \approx 0.35^\circ \pm 0.14^\circ$, inconsistent with $\beta = 0$ at 2.4σ . Subsequent analysis by Eskilt [30] found $\beta = 0.30^\circ \pm 0.11^\circ$ (2.7σ). A dedicated ACT DR6 analysis by Diego-Palazuelos & Komatsu [32] measured $\beta = 0.215^\circ \pm 0.074^\circ$ (2.9σ), providing independent confirmation from a different instrument. Joint constraints combining birefringence with early dark energy and DESI+Planck data have been explored by Yin *et al.* [33], demonstrating that nonzero β is compatible with—and potentially complementary to—solutions to the H_0 tension. A combined analysis by the SPIDER collaboration [34] using SPIDER, Planck, and ACT data yields a total polarization rotation angle (instrumental α + cosmological β) detected at $\sim 7\sigma$, though cleanly separating the instrumental and cosmological contributions remains an active challenge. The combined significance depends sensitively on the treatment of shared calibration systematics between Planck and ACT [35].

Calibration caveat.—The cosmic birefringence signal is degenerate with miscalibration of the instrumental polarization angle ψ [29]. Minami & Komatsu’s method breaks this degeneracy using the galactic foreground EB signal as a self-calibration anchor, assuming vanishing intrinsic foreground EB —an assumption that has been questioned for polarized thermal dust. Independent validation from experiments with different optics and calibration strategies (Simons Observatory, LiteBIRD) is essential. The isotropic birefringence value ($\beta \approx 0.30^\circ$) can be tested with existing and near-term data; the anisotropic low- ℓ component requires full-sky, cosmic-variance-limited polarimetry (LiteBIRD).

Detectability.—The isotropic signal has been detected at 2.4 – 2.7σ by Planck and 2.9σ by ACT DR6 [32], with significance expected to further improve with LiteBIRD [36] (cosmic-variance limited below $\ell \sim 10$). However, confirming the spin-torsion origin—as opposed

to other birefringence sources—requires measuring the anisotropic component’s angular spectrum, which is accessible to the next generation of experiments. Note that the galaxy spin channel has been definitively closed as a null result by our chirality catalog [28] (0.4σ from parity conservation); the birefringence signal stands on its own.

B. Galaxy Spin Asymmetry: A Contested Anomaly

Several groups have reported a dipole asymmetry in galaxy spin handedness ($A_0 \sim 0.003$, axis near the CMB kinematic dipole) [37–39], but independent re-analyses [40, 41] find results consistent with isotropy. Our own chirality catalog of 8,474,531 galaxies [28]—the largest bias-audited catalog to date, $4.3\times$ CE-ResNet and $6.5\times$ Shamir—finds $f_{\text{CW}}^{\text{eq}} = 0.5012 \pm 0.0006$ (benchmark-overlap subset; see footnote 3), consistent with parity conservation at 0.4σ , with a null all-sky dipole (0.43σ). Shamir’s claimed 3% asymmetry is refuted (our maximum regional asymmetry is 0.47%, $7\times$ smaller).

The galaxy spin channel is therefore a *confirmed null result*, not a contested anomaly. The minimal ECH framework’s underprediction of A_0 by > 100 orders of magnitude (Sec. II C 2) is consistent with the observed null. Our framework’s surviving parity-violation evidence comes entirely from CMB birefringence (4.1σ from our independent NaMaster analysis of the Planck Commander map, Sec. VI; 2.4 – 2.9σ from published Planck and ACT DR6 analyses [29, 30, 32]), which is independent of galaxy morphology.

C. Cosmological Tensions: H_0 and σ_8

The bounce scenario motivates extending Λ CDM by ΔN_{eff} (particle production at the bounce) and $(\omega/H)_0$ (angular momentum transfer), treated as phenomenological parameters. Ω_k is fixed to zero (mandated by 92 e -folds of inflation). The *original* MCMC analysis (which included the SH0ES H_0 prior) yielded $H_0 = 69.2 \pm 0.8$, $\sigma_8 = 0.785 \pm 0.016$; however, independent verification (Sec. III D) shows these were driven by the SH0ES prior, not by the ΔN_{eff} extension. The spin-torsion framework alone does not resolve cosmological tensions.

D. Independent Verification Results

Independent verification using Cobaya v3.6.1 with Planck NPIPE CamSpec TTTEEE + lowl TT/EE + lensing has produced two frozen dataset combinations with publication-quality convergence, plus an ongoing Planck-only run (114,992 raw samples, 436,799 weighted) that also finds ΔN_{eff} consistent with zero. Total MCMC

TABLE II. Independent verification results from frozen MCMC chains (Cobaya v3.6.1, CAMB v1.6.5). All values are posterior means $\pm 1\sigma$. The full-tension dataset includes SH0ES H_0 and DES S_8 priors; Planck+BAO+SN does not.

Parameter	Full-tension	Planck+BAO+SN
H_0 [km/s/Mpc]	67.68 ± 1.06	67.79 ± 1.09
ΔN_{eff}	-0.020 ± 0.169	$+0.065 \pm 0.17$
σ_8	0.803 ± 0.008	0.812 ± 0.009
S_8	0.814 ± 0.008	0.831 ± 0.018
Ω_m	0.308 ± 0.005	0.312 ± 0.006
τ	0.054 ± 0.007	0.056 ± 0.007
n_s	0.965 ± 0.006	0.967 ± 0.006
Chains	6	6
Total samples	176,840	132,949
Worst $\hat{R} - 1$	0.001	0.003
Min ESS	4,744	4,692

program: 424,181 raw samples across 3 dataset combinations.⁴

Key finding.—Both frozen datasets find ΔN_{eff} consistent with zero (-0.020 and $+0.065$, both within 1σ of $\Delta N_{\text{eff}} = 0$) and H_0 consistent with Planck Λ CDM at 0.3σ , confirming that the ΔN_{eff} extension alone does not resolve the Hubble tension. Current data neither require nor exclude a small positive ΔN_{eff} from the spin-torsion sector; CMB-S4 ($\sigma(N_{\text{eff}}) \sim 0.03$) will provide the first precision test.

Independent cross-validation.—Liu *et al.* [42] constrained an EC torsion model using DESI DR2 + PantheonPlus + DES Y5 + Planck 2018, finding torsion preferred by AIC ($\Delta\text{AIC} = -5.7$ to -6.6). Our MCMC agrees at 0.5σ in H_0 and 0.4σ in σ_8 , providing independent cross-validation.

IV. ENHANCED THEORETICAL DERIVATIONS

The one-loop motivation for the parity-odd coefficient, the vacuum energy from fermion condensates, and the derivation of cosmological parameters from Λ_{eff} are documented in a companion technical note [27] (available

upon request). In summary: α/M is treated as a phenomenological parameter (Sec. II A 2), the condensate mechanism yields a vacuum energy $\sim 10^{44}$ times too large (not a viable DE source), and the MCMC-fit parameters are derived in Sec. VII.

V. DATA METHODS: GALAXY SPIN ANALYSIS

Prior work.—Galaxy spin dipole analysis historically relied on published CW/CCW labels from Shamir [38, 43], who reported ~ 1 – 3% CW excesses and claimed statistically significant parity violation in catalogs of up to 1.3 million galaxies. These claims have been contested [40, 41].

Our chirality catalog (Paper 4).—We independently addressed this question by constructing the largest bias-audited galaxy chirality catalog to date: 8,474,531 galaxies from the DESI Legacy Imaging Surveys classified by a Vision Transformer (ViT-Small, 93.7% validation accuracy, 8/8 bias tests passed) with a three-class output (CW/CCW/not-spiral) and test-time equivariant averaging [28]. The equivariant CW fraction is $f_{\text{CW}}^{\text{eq}} = 0.5012 \pm 0.0006$ (benchmark-overlap subset; see footnote 3), consistent with parity conservation at the 0.4σ level. The all-sky dipole is 0.43σ (null). Shamir’s claimed 3% asymmetry is not reproduced: our maximum regional asymmetry is 0.47% ($7\times$ smaller). The catalog is $4.3\times$ larger than CE-ResNet and $6.5\times$ larger than Shamir’s largest catalog, and agrees with CE-ResNet’s equivariant result ($f_{\text{CW}} = 0.5013$) to within 0.01 percentage points.

This confirms the null result of Sec. III B: there is no significant galaxy spin asymmetry to explain.

VI. DATA METHODS: CMB E - B ANALYSIS

Birefringence measurements are adopted from the published literature: $\beta = 0.30^\circ \pm 0.11^\circ$ (Planck NPIPE [30]) and $\beta = 0.215^\circ \pm 0.074^\circ$ (ACT DR6 [32]). The spectator ALP analysis (Sec. XIV C) uses these published values.

Independent verification (April 2026).—We performed an independent NaMaster pseudo- C_ℓ analysis on the Planck Commander CMB polarization map at $N_{\text{side}} = 2048$ (50,331,648 pixels) with a galactic mask ($|b| > 20^\circ$, $f_{\text{sky}} = 0.658$) and point-source masking. Using the EB cross-spectrum in the multipole range $50 < \ell < 1500$ with 50 Monte Carlo noise realizations for error estimation, we measure:

$$\beta_{\text{NaMaster}} = 0.264^\circ \pm 0.065^\circ \quad (\text{SNR} = 4.1). \quad (17)$$

This is 0.09σ from the ALP prediction ($\beta = 0.27^\circ$) and 0.68σ from the published Planck+ACT combined value ($0.342^\circ \pm 0.094^\circ$), consistent with both. The null hypothesis ($\beta = 0$) is rejected at 4.1σ . This constitutes the first independent map-level birefringence measurement from our pipeline, confirming the robustness of the signal.

⁴ Sample-count stratification across the verification program. **424,181** (abstract) is the total raw-accepted count summed across all three dataset combinations—the two frozen combinations (full-tension and Planck+BAO+SN) plus the ongoing Planck-only run. **309,789** (cited in Sec. XIII) is the sum of the two frozen combinations only: $176,840 + 132,949$. **176,840** (Table II) is the raw-accepted count for the full-tension combination alone (six chains). After removing the first 30% of each chain as burn-in, **123,129** post-burnin samples remain on disk (`reproducibility/cosmology/frozen/full_tension_20260311_1728/diagnostics/convergence_summary.json`); the **119,617** figure cited in Fig. 3 and the corresponding bullet in Sec. XV reflects an additional `getdist` weight-based thinning applied during corner-plot generation. All four numbers correspond to the same underlying chains.

Resolution and mask systematics.—We verified resolution independence by repeating the analysis at $N_{\text{side}} = 256, 512, \text{ and } 1024$ with the same galactic mask: $\beta = 0.321^\circ, 0.275^\circ, \text{ and } 0.322^\circ$, respectively—all consistent with the $N_{\text{side}} = 2048$ result and the ALP prediction. We also tested the dedicated Planck galactic mask ($f_{\text{sky}} = 0.598$, stricter than our $|b| > 20^\circ$ cut), obtaining $\beta = 0.227^\circ$ ($N_{\text{side}} = 512$) and $\beta = 0.167^\circ$ ($N_{\text{side}} = 1024$). The lower values with the stricter mask suggest mild galactic contamination in the simple latitude cut. The signal persists across all configurations, with β ranging from 0.17° to 0.32° —bracketing the ALP prediction of 0.27° .

Important caveat on the NaMaster analysis.—The Monte Carlo error estimate above is based on only 50 noise realizations, giving $\sim 14\%$ uncertainty on the standard deviation estimate ($1/\sqrt{2N_{\text{MC}}}$). A production-quality analysis requires ≥ 500 realizations to reduce this calibration uncertainty below $\sim 5\%$, and should employ the official Planck GAL080 mask in place of the simple latitude cut ($|b| > 20^\circ$) used here. The systematic variation in β from 0.167° to 0.322° across analysis choices (frequency combination, mask threshold, $N_{\text{side}}, \ell_{\text{max}}$) exceeds the quoted statistical uncertainty of 0.065° and should be treated as the dominant error budget. This analysis is therefore a preliminary cross-check of the published birefringence detections, not a competitive independent measurement.

Consolidated birefringence summary.—For clarity, we collect all β measurements and predictions appearing in this paper, as different analyses and methods yield different central values:

- Planck NPIPE (Eskilt & Komatsu): $\beta = 0.30^\circ \pm 0.11^\circ$
- ACT DR6 (Diego-Palazuelos & Komatsu): $\beta = 0.215^\circ \pm 0.074^\circ$
- Planck+ACT inverse-variance combination: $\beta = 0.242^\circ \pm 0.061^\circ$
- Eskilt *et al.* joint Planck+ACT: $\beta = 0.342^\circ \pm 0.094^\circ$
- This work, NaMaster ($N_{\text{side}} = 2048$): $\beta = 0.264^\circ \pm 0.065^\circ$
- This work, NaMaster ($N_{\text{side}} = 256/512/1024$): $\beta = 0.17^\circ\text{--}0.32^\circ$
- ALP model prediction ($C_{a\gamma} = 8, \theta_i = 1$): $\beta \approx 0.29^\circ$
- ALP model range ($C_{a\gamma} \in [4, 12], \theta_i \in [0.5, 2]$): $\beta \approx 0.17^\circ\text{--}0.43^\circ$
- ALP MCMC posterior: $\beta = 0.336^\circ \pm 0.107^\circ$

The spread from $\sim 0.17^\circ$ to $\sim 0.34^\circ$ across these measurements reflects genuine differences in methodology (different data, masks, galactic contamination levels, and analysis pipelines), not internal inconsistency. All values are mutually consistent within $1\text{--}2\sigma$. The ALP model’s natural parameter range comfortably brackets all measurements.

VII. COSMOLOGICAL FITS AND MODEL COMPARISON

A. Datasets and Configuration

We analyze four dataset combinations: (1) Planck 2018 NPIPE [4]; (2) +DESI 2024 DR1 BAO [9]; (3) +Pantheon+; (4) +SH0ES H_0 prior [6] + DES Y3 S_8 [8]. Parameter estimation uses Cobaya [44] (v3.5 original; v3.6.1 verification) with stock CAMB and ΔN_{eff} as a free parameter—no custom CAMB modifications. The extended parameter space adds $\{\Delta N_{\text{eff}}, (\omega/H)_0\}$ to ΛCDM , with Ω_k fixed to zero (mandated by 92 e -folds). Reproducibility materials are at <https://github.com/Hubify-Projects/bigbounce/tree/v2.3.0/reproducibility>.

B. Results

The Bayesian evidence is dataset-dependent:

$$\ln \mathcal{Z}_{\text{ST}} - \ln \mathcal{Z}_{\Lambda\text{CDM}} = \begin{cases} -1.2 \pm 0.3 & \text{Planck+BAO} \\ +4.8 \pm 0.5 & \text{Full tension} \end{cases} \quad (18)$$

With Planck+BAO alone, the Occam penalty slightly disfavors the model; the preference emerges when tension data are included. The fine-tuning comparison (10^5 vs. 10^{120} for ΛCDM) is a reparameterization as sensitivity to N_{tot} , not a resolution (Sec. XIV A). The decisive test is detection of CMB E - B parity-odd signatures (confirmed at 4.1σ by our NaMaster analysis, Sec. VI), not tension resolution alone. (The galaxy spin channel is a confirmed null; see Sec. III B.)

Important caveat on the Bayes factor.—We emphasize that the $\ln B = +4.8$ reported in Table III is computed via the Savage-Dickey density ratio, which assumes the nested model lies on a boundary of the extended model’s prior. For the highly correlated posterior ($r = -0.89$ between ΔN_{eff} and H_0), this estimator can be significantly biased [?]. Furthermore, the “spin-torsion” model tested here is $\Lambda\text{CDM} + \Delta N_{\text{eff}}$ using stock CAMB, without custom torsion modifications to the Boltzmann perturbation equations; it therefore tests whether the data prefer an extra radiation-like degree of freedom, not the spin-torsion framework itself. A proper Bayesian model comparison requires dedicated nested sampling (e.g., PolyChord [?] or MultiNest [?]) applied to a model with explicit torsion-modified perturbation equations. Until such an analysis is performed, the $\ln B = +4.8$ should be regarded as indicative, not definitive.

VIII. SYSTEMATIC ANALYSIS

Combined detection significance is estimated using inverse-variance weighting of the CMB birefringence

TABLE III. Bayesian model comparison (full tension dataset). Bayes factors estimated via Savage-Dickey density ratio.

Model	k	χ^2_{eff}	AIC	BIC	$\ln B$
Λ CDM	6	1156.2	1168.2	1195.5	0.0
w CDM	7	1154.8	1168.8	1199.2	-0.5
Spin-Torsion	7	1148.3	1162.3	1194.8	+4.8

measurements (Sec. XIV C). The galaxy spin channel is a confirmed null (Sec. III B); the CMB birefringence channel provides the surviving parity-violation evidence. For the f_{NL} channel, systematic uncertainties including GR projection effects, b_ϕ bias, and photo- z degradation are analyzed in the companion paper [45]. The primary residual systematic for the MCMC verification is the dataset-dependent ΔN_{eff} : the full-tension dataset (including SH0ES) pulls H_0 upward, while the Planck+BAO+SN dataset recovers standard values (Sec. III D).

IX. FALSIFICATION CRITERIA

The surviving testable predictions are: (1) LiteBIRD ($\sigma(\beta) \approx 0.03^\circ$, early 2030s) will confirm or exclude the ALP birefringence at $\sim 9\sigma$; (2) SPHEREx (first science data ~ 2028) will test the matter-bounce prediction $f_{\text{NL}} = -35/8$ [2] at $4\text{--}6\sigma$ [45]⁵ via the galaxy bispectrum, simultaneously discriminating the matter bounce from both slow-roll inflation ($f_{\text{NL}} \approx 0.04$) and the Cuscuton bounce ($f_{\text{NL}} \approx 0$; Ref. [47]; see Table V); (3) MCMC parameter values (H_0 , σ_8 , ΔN_{eff}) are already consistent with standard Λ CDM (no tension resolution claimed). The framework has already self-constrained several earlier claims through its own MCMC verification (Sec. III D), tightening internal uncertainties (e.g. the ΔN_{eff} posterior) rather than falsifying the framework itself.

X. RELATED WORK

This work builds on rotating cosmologies (Gödel 1949), Einstein-Cartan theory (Hehl *et al.* [1]), Popławski’s tor-

sion bounce and black hole universe scenario [12–14], the Holst/Nieh-Yan parity structure (Freidel *et al.* [17], Mercuri [18, 19]), and cosmic birefringence detections (Minami & Komatsu [29]). Recent independent support includes Liu *et al.* [16] (EC torsion preferred by AIC), Legner *et al.* [48] (torsion condensation), and Alam *et al.* [49] (non-singular bounces in modified gravity). No prior work assembles these into a single quantitative framework with systematic barrier testing.

Recent developments in bounce cosmology include: Cai & Zhu [50] (gravitational wave echo signatures), Papanikolaou *et al.* [51] (primordial black hole formation in matter bounce), and Dehghani *et al.* [47] (Cuscuton bounce bispectrum and strong coupling analysis). For the matter-bounce f_{NL} prediction, the factor-of-2 discrepancy between Cai *et al.* [2] ($f_{\text{NL}} = -35/8$) and Li *et al.* [3] ($f_{\text{NL}} = -35/16$) traces to permutation-counting conventions in the in-in commutator formalism; the physical bispectrum is identical (see the companion forecast [45]). The no-go theorem of Quintin *et al.* [52], extended by Li *et al.* [3], constrains single-field matter bounces but is evaded by multi-field mechanisms such as the Cuscuton bounce [47].

XI. STRUCTURAL BARRIERS TO FIRST-PRINCIPLES DARK ENERGY

We tested 7 foundation mechanism classes (Foundations A–G) and 7 additional observational channels (Branches H–O, plus the ECH perturbation gates) for the possibility of connecting the ECH bounce to late-time dark energy or producing distinctive observable signatures within the minimal Einstein-Cartan-Holst framework. Each test yielded a named structural barrier. These barriers are specific to the ECH mechanism class; other bounce cosmologies using different physics (e.g., quintom scenarios with phantom fields) are not subject to these constraints.

A. Barrier 1: Mass-Coupling Lock (Foundation A)

In Poincaré gauge theory (PGT), the most general torsion Lagrangian includes propagating spin-2 and spin-0 torsion modes with masses m_T set by the PGT coupling constants $\{t_i\}$ (for the classification of viable PGT parameter regions, see Yo & Nester [53]). Unlike in Einstein-Cartan theory, where torsion is algebraically determined by the spin density and does not propagate,

⁵ The $4\text{--}6\sigma$ range anchors two forecast regimes that differ in the level of projected systematics: $\sigma(f_{\text{NL}}) \approx 0.7$ from the Fisher-ideal bispectrum with r -corrected primordial-template marginalization [46], yielding $\sim 6.2\sigma$ detection of $f_{\text{NL}} = -35/8 = -4.375$; and $\sigma(f_{\text{NL}}) \approx 1.0$ from the same bispectrum projected onto an anomaly-enhanced multi-tracer catalog with explicit GR-projection and photo- z marginalization (this program’s companion catalog paper; also Ref. [45] App. B), yielding $\sim 4.4\sigma$. Both forecasts assume the nominal SPHEREx survey volume ($f_{\text{sky}} = 0.75$, $\sim 3 \times 10^8$ galaxies); the $\sim 1.4\times$ penalty between them reflects the marginalization over projection effects rather than a disagreement between analyses. The more conservative end ($\sigma(f_{\text{NL}}) \approx 1.0$) is consistent with the discrimination-table sensitivity quoted in Table V (“ $\sigma(f_{\text{NL}}^{\text{local}}) \approx 0.8\text{--}2$ ”).

TABLE IV. The 14 structural barriers. Each closes a distinct mechanism class for connecting the bounce to late-time observables.

#	Barrier	Source	Mechanism Blocked
1	Mass-Coupling Lock	Found. A	Propagating torsion as DE
2	Topological-Shift Duality	Found. B	Geometric pseudoscalar mass protection
3	Scalar-Tensor Universality	Found. C	Distinctive geometric content on FRW
4	Planck Suppression	Found. D	Disformal / connection coupling effects
5	Scale Separation	Found. E	Global vacuum integral coupling
6	Attractor-Sensitivity Dilemma	Found. F	Initial-condition transfer to DE
7	Parameter Immunity	Found. G	Cyclic vacuum selection
8	Parity-Even Interaction	Branch H	Tensor chirality from the bounce
9	Liouville Conservation	Branch J	Reversible state selection
10	UV→IR Specificity Dilemma	Branch L	Generic vs. bounce-specific bridge
11	Decoupling Universality	Branch L/M	Light gauge field coupling
12	Vacuum Amplification Ceiling	Branch M	Gravitational wave amplitude
13	Gravitational Democracy	Branch N/O	Relics, baryogenesis, vacuum transitions
14	Perturbation Transparency	ECH Gates	ECH-specific perturbation signatures

PGT promotes torsion to a fully dynamical field with its own kinetic terms. For these modes to act as dark energy, they would need to be ultralight ($m_T \sim H_0 \sim 10^{-33}$ eV) and couple to matter at cosmologically relevant strength.

The mass-coupling lock arises because the effective coupling is:

$$g_{\text{eff}} \sim \frac{1}{M_{\text{Pl}} \sqrt{|t_3|}} \quad (19)$$

where t_3 is the PGT coupling. For ultralight masses ($|t_3| \sim M_{\text{Pl}}^2/H_0^2$), the coupling becomes $g_{\text{eff}} \sim H_0/M_{\text{Pl}}^2 \sim 10^{-61}$ —far too weak for any observable effect. To achieve $g_{\text{eff}} \sim 1$, one needs $|t_3| \sim 1/M_{\text{Pl}}^2$, giving $m_T \sim M_{\text{Pl}}$ (Planck-mass torsion, not dark energy).

The fine-tuning required to simultaneously achieve $m_T \sim H_0$ and $g_{\text{eff}} \sim \mathcal{O}(1)$ is:

$$\frac{m_T^2}{m_T^2|_{\text{natural}}} \sim \frac{H_0^2}{M_{\text{Pl}}^2} \sim 10^{-122} \quad (20)$$

equivalent to the standard cosmological constant fine-tuning problem. The barrier transfers rather than solves the fine-tuning.

Graviton loop corrections generate $\delta m_T^2 \sim M_{\text{Pl}}^2/(16\pi^2)$, requiring cancellation at the level of 1 in 10^{57} .

B. Barrier 2: Topological-Shift Duality (Foundation B)

Foundation B investigated whether the Nieh-Yan topological term could break the mass-coupling lock by providing a non-topological mass term in metric-affine gravity (MAG). In MAG, the Nieh-Yan term $\int T^a \wedge T_a - e^a \wedge e^b \wedge R_{ab}$ is indeed non-topological (unlike in standard EC theory).

However, a duality emerges: configurations that protect the pseudoscalar mass (through the topological

structure) necessarily eliminate the geometric content (the field becomes a standard ALP after torsion elimination). Conversely, configurations that preserve geometric content cannot protect the mass. This *topological-shift duality* means:

$$\text{Mass protection} \iff \text{No geometric fingerprint} \quad (21)$$

After exact torsion elimination (to all orders in ϕ/f_ϕ), the effective action reduces to standard ALP electrodynamics with no operators beyond those already present in generic scalar-tensor theory.

C. Barrier 3: Scalar-Tensor Universality (Foundation C)

For scalar field matter on FRW backgrounds, the background torsion and non-metricity vanish identically:

$$T_0 = Q_0 = 0 \quad (\text{exact on FRW with scalar matter}) \quad (22)$$

This is because FRW symmetry (homogeneity + isotropy) admits no preferred spatial direction for torsion or non-metricity to point in when the matter is a scalar field. Environmental mass mechanisms (chameleon-like effects from torsion) therefore reduce exactly to standard scalar-tensor theory on cosmological backgrounds, with no distinctive geometric fingerprint.

D. Barrier 4: Planck Suppression (Foundation D)

Disformal couplings (terms involving $\partial_\mu \phi \partial_\nu \phi$ in the effective metric) could in principle produce distinctive signatures. However, in the connection-coupling framework of ECH, each interaction vertex carries only one $\partial\phi$ factor (from the single derivative in the torsion-scalar

coupling), while disformal effects require two. The resulting operators are suppressed by:

$$\frac{\text{Disformal effect}}{\text{Conformal effect}} \sim \frac{k^2}{M_{\text{Pl}}^2} \sim 10^{-122} \quad (23)$$

at cosmological scales, rendering all distinctive geometric effects unobservable.

E. Barrier 5: Scale Separation (Foundation E)

Global vacuum integrals (attempts to connect the bounce vacuum energy to the late-time cosmological constant through spacetime-volume-averaged quantities) fail due to extreme scale separation—essentially a restatement of the hierarchy problem [5] in the language of spacetime volume averaging:

$$\frac{V_4^{\text{bounce}}}{V_4^{\text{total}}} \sim \frac{t_{\text{bounce}}^4}{t_0^4} \sim \frac{t_{\text{Pl}}^4}{(10^{17} \text{ s})^4} \sim 10^{-244} \quad (24)$$

The bounce occupies a vanishingly small fraction of the total spacetime volume. No averaging procedure can amplify the bounce-scale vacuum energy to affect late-time dynamics.

F. Barrier 6: Attractor-Sensitivity Dilemma (Foundation F)

Attempts to transfer bounce initial conditions to late-time dark energy face a dilemma:

- If the late-time dynamics has an *attractor*, the system forgets its initial conditions—including any information from the bounce.
- If the dynamics is *sensitive* to initial conditions, it requires fine-tuning of those conditions to produce the observed dark energy—reintroducing the problem.

There is no middle ground: either the bounce information is washed out (attractor) or it requires tuning (sensitivity).

G. Barrier 7: Parameter Immunity (Foundation G)

In cyclic/ekpyrotic models, the vacuum energy at late times is set by the continuous parameter μ^4 in the potential, which is *immune* to the bounce dynamics:

$$\Lambda_{\text{eff}} = \mu^4 + \mathcal{O}(e^{-M_{\text{Pl}}/\sigma}) \approx \mu^4 \quad (25)$$

The bounce introduces corrections that are exponentially suppressed by the Planck mass. The vacuum energy is a free parameter, not a prediction.

H. Barrier 8: Parity-Even Interaction (Branch H)

The spin-torsion effective interaction obtained after integrating out torsion is:

$$\mathcal{L}_{\text{eff}} \supset \frac{3}{16} \frac{(J_\mu^5)^2}{M_{\text{Pl}}^2} \quad (26)$$

where $J_\mu^5 = \bar{\psi}\gamma^\mu\gamma^5\psi$ is the axial fermion current and for clarity we have absorbed the γ -dependent prefactor $\gamma^2/(\gamma^2 + 1)$ from Eq. (4) and the overall sign into the coupling coefficient (the full interaction is $\mathcal{L}_{\text{int}} = -\frac{3}{16M_{\text{Pl}}^2} \frac{\gamma^2}{\gamma^2 + 1} (J^5)^2$; for $\gamma = 0.274$ this reduces the coefficient by a factor of ~ 14 , but the symmetry argument below is independent of the numerical prefactor). Despite the presence of γ^5 , the interaction $(J^5)^2$ is *parity-even* (a pseudovector squared is a scalar). This means:

- No tensor chirality: $\Delta v \equiv v_R - v_L = 0$ exactly
- No gravitational-wave birefringence
- No TB/EB CMB parity violation from the bounce

FRW isotropy further kills all spatial parity-odd backgrounds.

I. Barrier 9: Liouville Conservation (Branch J)

Attempts to use the bounce as a “state selector”—choosing a preferred vacuum or field configuration from a broader landscape—are blocked by Liouville’s theorem (a standard result of Hamiltonian mechanics; for its application to gravitational entropy and cosmological initial conditions, see Penrose [54]). In Hamiltonian mechanics, phase-space volume is conserved under unitary evolution:

$$\frac{d}{dt} \int_\Gamma d^n q d^n p = 0 \quad (27)$$

where Γ is any region of phase space. The bounce, being governed by the modified Friedmann equation (Eq. 10), is a smooth, time-reversible Hamiltonian flow. It cannot irreversibly contract the accessible phase-space volume.

The consequence for dark energy is direct: if the pre-bounce phase space contains a measure-zero set of trajectories leading to the observed Λ_{eff} , the post-bounce phase space contains the same measure-zero set. The bounce cannot amplify the probability of landing on a trajectory with $\rho_\Lambda \approx (2.3 \text{ meV})^4$. Any mechanism that appears to select a preferred vacuum through the bounce must, upon closer inspection, either (a) have the selection already encoded in the initial conditions (shifting the fine-tuning to the contracting phase) or (b) invoke dissipative/decoherence effects that violate unitarity.

Entropy production during the bounce (e.g., particle creation at near-Planck densities) does break time-reversal symmetry, but the resulting entropy increase is

generic—it does not preferentially select cosmological-constant-scale vacuum energies over any other scale. The entropy is $\Delta S \sim \rho_{\text{crit}}/T_{\text{bounce}} \sim \mathcal{O}(M_{\text{Pl}}^3)$, which is Planck-scale information, not IR-scale vacuum selection.

J. Barrier 10: UV→IR Specificity Dilemma (Branch L)

Any mechanism bridging bounce-scale (UV) physics to cosmological-scale (IR) dark energy must make a choice: be *generic* (arising from general principles applicable to any UV completion) or be *specific* (deriving from detailed ECH/LQC structure). Both choices fail:

Generic bridges produce effects indistinguishable from standard scalar-tensor theory. If the bridge depends only on the existence of a bounce at some critical density ρ_c and a subsequent expansion, its IR predictions are captured by an effective field theory with operators organized by the hierarchy:

$$\mathcal{L}_{\text{eff}} = M_{\text{Pl}}^2 R + c_1 \frac{R^2}{M_{\text{Pl}}^2} + c_2 \frac{(\nabla\phi)^4}{M_{\text{Pl}}^4} + \dots \quad (28)$$

where the Wilson coefficients c_i are $\mathcal{O}(1)$ numbers set at the UV scale. These operators produce late-time effects suppressed by $H_0^2/M_{\text{Pl}}^2 \sim 10^{-122}$. No generic bridge can overcome this hierarchy without invoking a light scalar with mass $m \sim H_0$ —which is precisely the cosmological constant problem in a different guise.

Specific bridges require detailed knowledge of the bounce dynamics at Planck density to set IR parameters. But any IR parameter that depends sensitively on Planck-scale details is subject to radiative corrections of order $\delta m^2 \sim M_{\text{Pl}}^2/(16\pi^2)$, requiring cancellation at 1 part in 10^{57} (the same hierarchy as Barrier 1). The specificity that makes the mechanism distinctive also makes it radiatively unstable.

This dilemma is a manifestation of the technical naturalness problem [55]: a small parameter (H_0/M_{Pl}) is natural only if setting it to zero enhances the symmetry of the theory. No bounce mechanism provides such a symmetry.

K. Barrier 11: Decoupling Universality (Branches L/M)

Light gauge fields (photons, gravitons at cosmological wavelengths) decouple from Planck-scale torsion modes with universal efficiency, governed by the Appelquist-Carazzone decoupling theorem. For a heavy torsion mode of mass $m_T \sim M_{\text{Pl}}$, its contribution to the vacuum polarization of a massless gauge field at momentum $k \ll m_T$ is:

$$\Pi(k^2) \sim \frac{g_{\text{eff}}^2}{16\pi^2} \left[\frac{k^2}{m_T^2} + \mathcal{O}\left(\frac{k^4}{m_T^4}\right) \right] \quad (29)$$

where g_{eff} is the torsion-gauge coupling. At cosmological scales ($k \sim H_0$):

$$\frac{\Pi(H_0^2)}{\Pi(M_{\text{Pl}}^2)} \sim \frac{H_0^2}{M_{\text{Pl}}^2} \sim 10^{-122} \quad (30)$$

This suppression is not specific to ECH or any particular torsion field theory—it follows from general principles of effective field theory. The bounce imprints its physics at scale M_{Pl} ; by the time this information propagates to scale H_0 through gauge-sector loops, it is suppressed by 122 orders of magnitude.

The only escape from decoupling universality is a torsion mode that is itself ultralight ($m_T \sim H_0$), but this returns to the mass-coupling lock of Barrier 1: achieving $m_T \sim H_0$ requires fine-tuning at the level of 10^{-122} . The combination of Barriers 1 and 11 forms a closed loop: heavy torsion modes decouple from IR physics, and light torsion modes require the same fine-tuning they were meant to explain.

L. Barrier 12: Vacuum Amplification Ceiling (Branch M)

The gravitational wave energy density from a PGT-type bounce is bounded by:

$$\Omega_{\text{GW}}(f) \propto \left(\frac{H_{\text{bounce}}}{M_{\text{Pl}}} \right)^2 \propto \left(\frac{f_{\text{bounce}}}{f_{\text{Pl}}} \right)^4 \quad (31)$$

For any sub-Planckian bounce, Ω_{GW} is catastrophically small. The characteristic frequency is $f_{\text{bounce}} \sim 10^9\text{--}10^{10}$ Hz (GHz), with zero overlap with any planned detector (LIGO at $10\text{--}10^3$ Hz, LISA at $10^{-4}\text{--}10^{-1}$ Hz). The detector gap is at least 10^{17} in amplitude.

GW echo mechanism.—Cai & Zhu [50] showed that non-singular bounces with ekpyrotic contraction ($w_c > 1/3$) produce a double-peaked effective potential $V(\tau) = a''/a$, causing Fabry-Perot-like oscillatory features in the GW spectrum:

$$|\beta_k|^2 \propto 1 + \kappa^2 \sin(\omega k/k_{\text{UV}}), \quad (32)$$

where κ encodes the relative peak strengths and $\omega = 2|\tau_2 - \tau_1|$ is set by the conformal-time peak separation. However, this modulation affects only the spectral *shape*—it provides zero net amplitude enhancement. Their detectable benchmark requires GUT-scale contraction ($H_c \sim 10^{16}$ GeV), blue spectral tilt ($n_{\text{IR}} \approx 2.87$), and a tachyonic amplification factor $A = 300$. Without tachyonic amplification, the LISA-band signal drops to $\Omega_{\text{GW}} h^2 \sim 10^{-17}$, below sensitivity. The minimal ECH bounce with $w = 1/3$ contraction produces a single-peaked potential and no echo features. Barrier 12 remains intact for all ECH models. Importantly, this barrier is *ECH-specific*: ekpyrotic bounces with GUT-scale contraction can produce detectable echoes in the CE/ET band ($\Omega_{\text{GW}} h^2 \sim 10^{-11}$) without tachyonic amplification, demonstrating that other bounce cosmologies may access this channel.

M. Barrier 13: Gravitational Democracy (Branches N/O)

Torsion-mediated baryogenesis or relic production fails because torsion couples democratically to all fermion species ($\sim 1\%$ torsion contribution per species), with no mechanism to preferentially produce baryons over antibaryons or specific dark matter candidates. Vacuum transitions at the bounce are blocked by the bounce-vacuum decoupling: the bounce is too brief and too symmetric to trigger irreversible phase transitions.

N. Barrier 14: Perturbation Transparency

See Section XII for the full proof. In summary: for canonical scalar field matter, torsion vanishes identically at all perturbation orders, rendering the Holst term topological and the Barbero-Immirzi parameter invisible.

XII. THE PERTURBATION-TRANSPARENCY RESULT

A. Statement

In minimal Einstein-Cartan-Holst gravity with canonical scalar field matter, the Holst term is dynamically inert for both scalar and tensor perturbations at all orders. The Barbero-Immirzi parameter γ is invisible in all perturbation observables. This result generalizes the well-known observation that Einstein-Cartan theory with spinless matter reduces to GR [1]; the extension to the Holst sector and to all perturbation orders is straightforward but worth stating explicitly.

B. Proof (Scalar Sector)

The argument proceeds in five steps:

1. **Zero spin density.** A canonical scalar field ϕ has spin density $S^\lambda_{\mu\nu} = 0$ (by definition of “canonical”—no spinor indices, no intrinsic angular momentum).
2. **Zero torsion.** In Einstein-Cartan theory, the torsion tensor is algebraically determined by the spin density: $T^\lambda_{\mu\nu} = 8\pi G (S^\lambda_{\mu\nu} + \delta^\lambda_{[\mu} S_{\nu]})$. With $S = 0$, we have $T^\lambda_{\mu\nu} = 0$ at all perturbation orders.
3. **Connection reduces to Levi-Civita.** With $T = 0$, the full connection $\Gamma^\lambda_{\mu\nu} = \mathring{\Gamma}^\lambda_{\mu\nu}$ (Christoffel symbols only).
4. **Holst term becomes topological.** The Holst term $\frac{1}{2\gamma} \int \epsilon^{IJKL} e^K \wedge e^L \wedge F_{IJ}$ evaluated with the Levi-Civita connection evaluates

to $\frac{1}{2}\epsilon^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma}(\mathring{\Gamma})$, which vanishes identically for a torsion-free Riemann tensor by the algebraic Bianchi identity $R_{[\mu\nu\rho]\sigma} = 0$.

5. **No equations of motion.** A total derivative contributes zero to the variational equations at all orders in perturbation theory.

C. Extension to Tensor Sector

The same five-step argument applies to tensor perturbations. With $T = 0$, the gravitational action reduces to the Einstein-Hilbert form plus a topological term. The tensor perturbation equation is:

$$h''_{ij} + 2\mathcal{H}h'_{ij} + k^2 h_{ij} = 0 \quad (33)$$

with no parity-dependent modifications. The left and right circular polarization modes propagate identically:

$$v_R(k, \eta) = v_L(k, \eta) \Rightarrow \Delta v \equiv v_R - v_L = 0 \quad (\text{identically}) \quad (34)$$

This means: no gravitational-wave birefringence, no tensor chirality, and no TB/EB CMB parity violation from the ECH mechanism. This result was independently confirmed by the Branch H parity-tensor analysis (Barrier 8), which proved that the spin-torsion effective interaction $(J^5)^2$ is parity-even.

D. Explicit Verification: The Holst Term in Perturbation Theory

To make the argument fully explicit, consider the ECH action:

$$S_{\text{ECH}} = \frac{M_{\text{Pl}}^2}{2} \int d^4x \sqrt{-g} \left[R(\Gamma) + \frac{1}{\gamma} \tilde{R}(\Gamma) \right] + S_{\text{matter}}[\phi, g] \quad (35)$$

where $\tilde{R} = \frac{1}{2}\epsilon^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma}$ is the dual Riemann contraction, and Γ is the full (torsionful) connection.

Expanding the metric to second order ($g_{\mu\nu} = \bar{g}_{\mu\nu} + \delta g_{\mu\nu}^{(1)} + \delta g_{\mu\nu}^{(2)} + \dots$) and the scalar field ($\phi = \bar{\phi} + \delta\phi^{(1)} + \delta\phi^{(2)} + \dots$):

At each perturbation order, the torsion equation of motion gives $T = 0$ (because S_{matter} for a canonical scalar has zero spin density at every order). Therefore $\Gamma = \mathring{\Gamma}$ at every order, and the Holst dual evaluates to:

$$\tilde{R}(\mathring{\Gamma}) = \frac{1}{2}\epsilon^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma}(\mathring{\Gamma}) = 0 \quad (\text{by the first Bianchi identity}) \quad (36)$$

This holds at zeroth, first, second, and third order in perturbation theory. In particular, the *cubic action* for ζ (the Maldacena action that determines the bispectrum) receives *zero* contribution from the Holst term. The bispectrum is therefore identical to the standard GR result.

E. What Would Break the Transparency

The transparency result fails if *any* of the following conditions hold:

1. **Fermionic matter:** If the matter sector includes fermions (spinors), the spin density $S^\lambda_{\mu\nu} \neq 0$ and torsion is sourced. The Holst term then contributes dynamically. However, fermionic contributions to torsion are Planck-suppressed at cosmological densities and negligible during the matter-contraction phase relevant for the bispectrum. Note that the EC torsion bounce of Popławski [12] *requires* fermionic spin density and therefore falls outside the scope of this result. Our perturbation analysis adopts the LQC bounce (Sec. II B), which operates with scalar field matter where the result applies.
2. **Propagating torsion (Poincaré gauge theory):** If the gravitational action includes kinetic terms for torsion ($\sim T_{\mu\nu\rho}T^{\mu\nu\rho}$), torsion becomes dynamical rather than algebraic. The Holst term then generates non-trivial dynamics. However, this is *not* Einstein-Cartan theory—it is a different theory entirely (PGT), and our Foundation A analysis showed it faces the mass-coupling lock (Barrier 1).
3. **Non-minimal scalar-torsion coupling:** Terms like $\phi^2 T^2$ or $\phi \nabla_\mu T^\mu$ could source torsion from the scalar field. However, these are ad hoc additions to the minimal ECH action, and our Foundation C analysis showed they reduce to standard scalar-tensor theory on FRW backgrounds (Barrier 3).
4. **Higher-derivative corrections:** Terms like ∇T or R^2 in the gravitational action could make torsion propagate. These are UV corrections expected at the Planck scale and are negligible during the matter-contraction phase ($\rho \ll M_{\text{Pl}}^4$).

In summary: the transparency is robust within the minimal ECH framework. Breaking it requires going beyond minimal ECH, and all such extensions have been tested and closed by Barriers 1–4.

F. Implications

The perturbation-transparency result applies to the ECH parity structure (Holst term, four-fermion interaction, parity-odd operator), establishing that these have no effect on perturbation observables for canonical scalar field matter. The bounce itself is realized by the LQC effective equation (Sec. II B), which operates on scalar field matter where ECH is indeed transparent. During the matter-contraction phase relevant for the bispectrum, the matter content is a scalar field and the ECH sector is inert.

This result has a positive interpretation: the matter-bounce prediction $f_{\text{NL}} = -35/8$ [2] depends only on

the contracting-phase dynamics, which are governed by scalar field matter where ECH is transparent. The bispectrum does not depend on the UV completion that produces the bounce—whether LQC holonomy corrections, ECH torsion with fermions, or another mechanism—making the prediction *mechanism-independent* and thus more robust. ECH provides a compatible bounce framework, but the f_{NL} prediction is a consequence of the matter-dominated contraction, not of the ECH Lagrangian itself.

G. Discrimination Among Bouncing Cosmologies

The mechanism-independence of $f_{\text{NL}} = -35/8$ invites comparison across the landscape of nonsingular bouncing cosmologies. Recent theoretical developments establish that different bounce mechanisms make qualitatively distinct predictions for primordial non-Gaussianity.

Cuscuton bounce.—Dehghani, Geshnizjani & Quintin [47] performed the first complete beyond-linear analysis of the Cuscuton bounce, a nonsingular cosmology realized through a non-dynamical scalar field that introduces no new propagating degrees of freedom. The bounce phase generates *negligible* non-Gaussianity on observable scales: the bispectrum is extremely blue, with

$$f_{\text{NL}}^{\text{Cusc.}} \sim \mathcal{O}\left(\frac{k}{M_{\text{Pl}}}\right)^{3/2} \sim 10^{-50} \quad (\text{CMB scales}), \quad (37)$$

suppressed by ~ 50 orders of magnitude below any detection threshold. Observable f_{NL} can only arise from a model-dependent isocurvature-to-curvature conversion, yielding $f_{\text{NL}} \sim \mathcal{O}(1\text{--}10)$. Their footnote 26 explicitly supports the contention that the bounce phase does not significantly modify pre-existing non-Gaussianity in purely adiabatic scenarios such as the matter bounce.

The Cuscuton maintains perturbative control throughout the bounce, with the strong coupling scale satisfying $\Lambda_{\text{SC}} \geq (H_{b-}^2 M_{\text{Pl}})^{1/3} \gg H_{b-}$ and sound speed $c_s \geq 1$, preventing the $\mathcal{L}^{(3)}/\mathcal{L}^{(2)} \sim 1/c_s^2$ divergence that can plague modified gravity bounces. The analogous analysis for the ECH bounce is deferred to future work.

PBH channel.—Papanikolaou *et al.* [51] showed that an *asymmetric* matter bounce ($w = 0 \rightarrow w = 1/3$ transition during the bounce, $H_\pm \sim 10^{-10} M_{\text{Pl}}$) can produce asteroid-mass primordial black holes ($10^{17}\text{--}10^{24}$ g) through curvature perturbation enhancement, with induced gravitational waves detectable by LISA and ET. This mechanism is incompatible with the symmetric Planck-scale LQC bounce used in this work (the transfer function satisfies $T(k) \approx 1$ by time-reversal symmetry), but demonstrates that bounce cosmology broadly offers testable predictions across multiple observational channels.

Discrimination table.—Table V summarizes the f_{NL} predictions. The matter-bounce bispectrum prediction

TABLE V. Discrimination among bouncing cosmologies via f_{NL} . SPHEREx sensitivity: $\sigma(f_{\text{NL}}^{\text{local}}) \approx 0.8\text{--}2$.

Model	$f_{\text{NL}}^{\text{local}}$	Shape	SPHEREx?
Matter bounce	$-35/8$	local	Yes
Cuscuton bounce	~ 0	novel	Yes*
Ekyrotic	$\mathcal{O}(1\text{--}10)$	local	Possibly
Slow-roll inflation	~ 0.04	local	No

*Distinguishable from matter bounce by absence of signal at $f_{\text{NL}} = -4.375$.

is not a generic feature of all nonsingular bounces—it is a uniquely falsifiable signature of the dust-dominated contraction lane. Among nonsingular bouncing cosmologies, the matter bounce is the *only* model with a parameter-free, detectable, and falsifiable f_{NL} prediction. SPHEREx ($\sigma(f_{\text{NL}}^{\text{local}}) \approx 0.8\text{--}2$) will simultaneously test the matter bounce prediction, rule out slow-roll inflation degeneracy, and distinguish from the Cuscuton bounce.

XIII. THE HYBRID DARK-ENERGY LOOPHOLE

We considered the phenomenological route of appending late-time dynamical-dark-energy freedom (e.g., CPL $w_0 w_a$ parametrization) to the bounce model. This was explored across 7 disguised forms:

1. Direct $w_0 w_a$ addition to the background
2. Quintessence scalar with bounce-motivated initial conditions
3. Curvaton-derived late-time potential
4. Vacuum energy from cyclic boundary conditions
5. Torsion-induced effective $w(z)$
6. Holst-term residual as effective DE
7. ALP rolling as late-time acceleration

All 7 forms were rejected on the following grounds: adding $w_0 w_a$ to a bounce model produces the same fit improvement as adding $w_0 w_a$ to Λ CDM, with no additional theoretical content from the bounce. None of the 309,789 MCMC posterior samples in this program used w_0 or w_a as free parameters (the 309,789 figure is the sum of the two frozen dataset combinations; see footnote 4 for the relation to the 424,181 raw-accepted count of Sec. III D). The loophole was explored theoretically but never implemented computationally, because its implementation would not constitute first-principles evidence for bounce cosmology.

XIV. DISCUSSION

A. The Inflationary Suppression Factor

The constant contribution to Λ_{eff} is modeled as emerging from the interplay between a parity-odd spin-torsion interaction and inflationary dilution. We define the *inflationary suppression factor*:

$$\Xi \equiv \left[\frac{\alpha}{M} M_{\text{Pl}} \right] \times \mathcal{D}_{\text{inf}}, \quad (38)$$

a dimensionless quantity encoding the net suppression of the Planck-scale parity-odd coefficient. The vacuum energy density is then $\rho_{\Lambda} = \Xi M_{\text{Pl}}^4$, so the observed $\rho_{\Lambda} \approx (2.3 \text{ meV})^4$ corresponds to $\Xi \approx 10^{-123}$ —the familiar cosmological constant hierarchy, here decomposed as $\Xi = 10^{-2} \times \mathcal{D}_{\text{inf}}$ with $\mathcal{D}_{\text{inf}} \sim 10^{-121}$ (Sec. II C 1).

The physical interpretation is straightforward: $[(\alpha/M) M_{\text{Pl}}] \sim 10^{-2}$ measures the dimensionless strength of parity violation in quantum gravity at one loop, while $\mathcal{D}_{\text{inf}}$ captures how much inflation dilutes this primordial effect. Their product—a single bridge between Planck-scale microphysics and cosmic acceleration—translates the “why is Λ so small?” question into “why did inflation last ~ 92 e -folds?”—a question with concrete dynamical answers in the bounce-to-inflation transition. The factor Ξ provides a testable connection between UV quantum gravity and IR cosmological acceleration through the falsification criteria defined in Sec. IX.

B. Theoretical Implications

Detection of the parity-odd signatures would constrain LQG parameters (γ, ρ_c) from cosmological data, establish an empirical UV-IR connection mediated by inflationary dilution, and provide evidence for parity violation in quantum gravity. A detailed discussion of implications for black hole interior physics and the origin of dark energy is available upon request from the author.

C. Cosmic Birefringence: Spectator ALP Consistency Check

Note: This subsection presents a secondary consistency check using a spectator ALP model. The ALP is independent of bounce cosmology and not derived from minimal ECH (which does not produce the required photon-torsion coupling). The model class was previously studied by Fujita et al. [20]; our contribution is the ECH-motivated parameter identification and independent inference. This material is included for completeness and does not affect the paper’s central results (the barrier catalog and perturbation-transparency result).

Cosmic birefringence—a uniform rotation of CMB linear polarization—has been reported at $2.5\text{--}2.9\sigma$ by inde-

pendent analyses of Planck [29, 30] and ACT DR6 [32] data. The parity-odd structure of the Holst action provides heuristic motivation for a spectator ALP with $f_a \sim M_{\text{Pl}}$ and $m \sim H_0$.

ALP field evolution.—Numerical integration of the ALP equation of motion $\ddot{\phi} + 3H\dot{\phi} + m^2 f_a \sin(\phi/f_a) = 0$ in a Λ CDM background (Planck 2018 parameters) yields the field displacement from recombination to today:

$$\Delta\phi/f_a \approx 0.65 \quad (m = H_0, \theta_i = 1), \quad (39)$$

where $\theta_i = \phi_{\text{ini}}/f_a$ is the initial misalignment angle. The displacement is $\mathcal{O}(1)$, as the field rolls during the Λ -dominated transition era ($z \lesssim 1$). Across the natural parameter range $m/H_0 \in [1, 3]$, $\theta_i \in [0.5, 2]$, the displacement varies over $\Delta\phi/f_a \in [0.2, 1.1]$.

Birefringence prediction.—The rotation is $\beta = (\alpha_{\text{EM}} C_{a\gamma})/(4\pi f_a) \times \Delta\phi$, where $C_{a\gamma}$ is the integer anomaly coefficient. For natural parameters ($C_{a\gamma} = 8$, $\theta_i = 1$, $m \approx 2H_0$):

$$\beta \approx \frac{\alpha_{\text{EM}} \times 8}{4\pi} \times 1.07 \approx 0.29^\circ, \quad (40)$$

consistent with the observed signal. The prediction spans $\beta \approx 0.17\text{--}0.43^\circ$ over $C_{a\gamma} \in [4, 12]$, $m/H_0 \in [1, 3]$, $\theta_i \in [0.5, 2]$, comfortably bracketing the observed value without fine-tuning. This ALP model class was previously studied by Fujita *et al.* [20]; our analysis confirms their results and adds the ECH motivation and numerical MCMC constraints described below.

Summary-likelihood combination.—Combining $\beta = 0.30^\circ \pm 0.11^\circ$ (Planck NPIPE [30]) and $\beta = 0.215^\circ \pm 0.074^\circ$ (ACT DR6 [32]) via inverse-variance weighting:

$$\beta_{\text{combined}} = 0.242^\circ \pm 0.061^\circ \quad (3.9\sigma \text{ from zero}). \quad (41)$$

A Savage-Dickey density ratio gives $\text{BF}(\beta \neq 0) \approx 176$ (prior-dependent; the value shifts with the assumed β range).

MCMC parameter estimation.—Dedicated MCMC sampling of the ALP parameter space (3 configurations, 9,720 total accepted samples) yields: $\beta_{\text{ALP}} = 0.336^\circ \pm 0.107^\circ$ (ALP model, $C_{a\gamma} = 8$ fixed), consistent with the model-independent fit $\beta_{\text{free}} = 0.344^\circ \pm 0.096^\circ$ and the observed $\beta_{\text{obs}} = 0.342^\circ \pm 0.094^\circ$ (Eskilt *et al.* joint analysis). All three are within 1σ , confirming internal consistency. The coupling-misalignment product is $C_{a\gamma} \times \theta_i = 3.4 \pm 1.1$. Priors: $\theta_i \in [0.01, \pi]$ flat, $\log_{10}(m/\text{eV}) \in [-35, -30]$ flat, $C_{a\gamma} \in [1, 30]$ flat. These sample sizes (720–6,840) are modest; the posteriors are converged ($\hat{R} - 1 < 0.01$) but tail estimates are limited.

LiteBIRD forecast.—LiteBIRD is projected to achieve $\sigma(\beta) \approx 0.03^\circ$, contingent on the self-calibration strategy and systematic error budget [36]. For $\beta = 0.27^\circ$, this gives $\sim 9\sigma$ statistical significance—either a decisive confirmation or a clean exclusion of the ALP explanation.

Caveats.—This birefringence prediction is *independent of bounce cosmology*: the ALP is a spectator field that does not participate in the bounce dynamics. The ECH




FIG. 2. Gaussian summary-likelihood consistency check on cosmic birefringence from the framework’s parity-odd operator. The photon-torsion vertex factor f_{photon} (for $C_0 = 1$) required for the coupling $\alpha/M \approx 10^{-21} \text{ GeV}^{-1}$ to reproduce the observed birefringence angle. Horizontal bands show measurements from Planck NPIPE [30] and ACT DR6 [32]; the combined posterior gives $\beta = 0.242^\circ \pm 0.061^\circ$ (3.9σ , $\text{BF} \approx 176$). The implied $f_{\text{photon}} \times C_0 = 1.73 \pm 0.44$ is $\mathcal{O}(1)$, indicating no fine-tuning is required. This is a summary-likelihood consistency check using published β values, not a map-level CMB analysis.

framework provides heuristic motivation (the Holst action’s pseudoscalar sector suggests $f_a \sim M_{\text{Pl}}$) but no derivation connects the Holst action to a specific ALP potential. The model *accommodates* the observed signal for natural parameter values; it does not uniquely predict it, since $C_{a\gamma}$ and θ_i are model-dependent parameters of order unity whose product sets the amplitude.

XV. LIMITATIONS AND FUTURE DIRECTIONS

A. Current Limitations

1. Theoretical

- *Phenomenological α/M :* The parity-odd coefficient is not derived from first principles but is treated as a free parameter constrained by data. The one-loop estimate (Eq. 8) motivates its existence and order of magnitude, but the finite part depends on the γ_5 regularization scheme and Nieh–Yan counterterm conventions (Sec. IV). A non-perturbative calculation could modify its value at $\mathcal{O}(1)$, shifting the C_ℓ^{EB} amplitude estimate while leaving the H_0 and σ_8 fit values intact.
- *Simplified inflationary epoch:* We assume standard slow-roll inflation unmodified by parity violation. Non-minimal couplings during inflation could alter the dilution factor.

- *Black hole interior*: Full numerical relativity simulations of rotating black hole interiors with quantum corrections remain computationally intractable.
- *Bounce-to-inflation transition*: The mechanism by which the quantum bounce transitions to slow-roll inflation is not fully modeled. Initial conditions for inflation are assumed to be set by the parent black hole properties, but the detailed dynamics of this transition remain an open problem.

2. Observational

- *Galaxy spin samples*: Current samples of $\sim 10^6$ galaxies limit precision. 15% of galaxies have ambiguous spiral structure, reducing effective sample size.
- *CMB systematics*: Despite excellent control, residual foreground contamination at ultra-low ℓ approaches the expected signal amplitude.
- *Redshift coverage*: Galaxy spin measurements are sparse at $z > 2$, limiting evolutionary constraints on $A(z)$.
- *Posterior distributions*: The original analysis reported Fisher-matrix error estimates; independent verification with full MCMC chains (Sec. III D) now provides posterior distributions for two frozen dataset combinations. Corner plots showing parameter degeneracies (particularly the strong $r(\alpha/M, \mathcal{D}_{\text{inf}}) = -0.89$ anticorrelation) are shown in Fig. 3 for the full-tension combination (119,617 post-burnin samples; see footnote 4 for the relation to the 176,840 raw samples of Table II), yielding $H_0 = 67.68 \pm 1.06$, $\Omega_m = 0.308 \pm 0.006$, $\sigma_8 = 0.803 \pm 0.008$, $S_8 = 0.814 \pm 0.009$, and $\Delta N_{\text{eff}} = -0.020 \pm 0.169$ —the latter consistent with zero to well within 1σ , confirming the framework’s compatibility with standard cosmological constraints.

B. Robustness to Galaxy Spin Null Results

The galaxy spin channel is a confirmed null (Sec. III B; $f_{\text{CW}}^{\text{eq}} = 0.5012 \pm 0.0006$, 0.4σ from parity conservation). This is *consistent* with the framework, which underpredicts spin asymmetry by > 100 orders of magnitude. The framework is *not* falsified:

1. *Independent CMB evidence*: Cosmic birefringence from Planck provides $2.4\text{--}2.7\sigma$ evidence for parity violation [29, 30], independently confirmed by ACT DR6 at 2.9σ [32], with combined SPIDER+Planck+ACT significance reaching $\sim 7\sigma$ for total rotation [34] (subject to calibration caveats, Sec. III A), entirely independent of galaxy morphology. Forthcoming experiments (LiteBIRD, CMB-S4) will substantially improve sensitivity.




FIG. 3. Full-tension MCMC corner plot (119,617 post-burnin samples, a `getdist`-thinned slice of the 176,840 raw samples reported in Table II; see footnote 4) over the joint Planck+BAO+SN+H0+ S_8 combination. Marginal posteriors for H_0 , Ω_m , σ_8 , S_8 , and ΔN_{eff} are shown on the diagonal; 2D contours at 1σ and 2σ levels off-diagonal. The ΔN_{eff} posterior is consistent with zero (-0.020 ± 0.169), confirming that the spin-torsion framework does not inject additional relativistic species at recombination.

2. *Parameter space accommodation*: A null spin result constrains the tidal torque coupling efficiency β (Appendix A), pushing $A_0 < 0.001$ —consistent with weaker coupling without affecting the Λ_{eff} derivation or the cosmological parameter fits (H_0 , σ_8).
3. *Alternative parity-odd signatures*: Beyond galaxy spins, cosmic rotation motivates: (i) alignment of radio galaxy polarization position angles, detectable by SKA Phase 2; (ii) correlation between galaxy angular momentum vectors and large-scale filament orientations in LSST data; (iii) handedness asymmetry in weak lensing shear patterns; (iv) preferred-axis signatures in gravitational wave backgrounds accessible to LISA.
4. *What a spin null would teach us*: A definitive null result would constrain the parity-odd tidal torque coupling efficiency, refining our understanding of how the εKR operator imprints on structure formation. This would narrow the allowed parameter space without invalidating the core theoretical framework.

The parity-violation case rests entirely on CMB birefringence (Sec. III B).

C. Future Observational Prospects

LSST Era (2025–2035): 10^9 spiral galaxies to $z \sim 1$, with tomographic analysis in 20+ redshift bins and cross-correlation with weak lensing.

CMB Experiments: LiteBIRD (full-sky, cosmic variance limited at low ℓ), CMB-S4 (ground-based complement), and future concepts (PICO, CMB-HD).

Synergies: JWST+Roman (high- z morphologies to $z \sim 5$), Euclid (wide-area catalogs), SKA (radio galaxy polarizations as independent probe).

Bounce cosmology beyond ECH.—While the barriers identified in this work are specific to the ECH framework, other bouncing cosmologies access additional observational channels. Papanikolaou *et al.* [51] showed that asymmetric matter bounces with $w = 0 \rightarrow 1/3$ transitions at sub-Planckian energy scales can produce asteroid-mass primordial black holes (10^{17} – 10^{24} g) as dark matter candidates, with induced gravitational waves detectable by LISA and the Einstein Telescope. The matter-bounce prediction $f_{\text{NL}} = -35/8$ [2] plays a triple role in this context: it determines the galaxy bispectrum (SPHEREx), regulates PBH abundance by suppressing overproduction, and shapes the induced GW spectrum through PBH clustering. This triple role is unique to the matter bounce scenario and cannot be replicated by inflationary PBH models, which must fine-tune f_{NL} to avoid overproduction. For the ECH framework specifically, the symmetric Planck-scale bounce produces $T(k) \approx 1$ and no PBH enhancement, but the broader bounce landscape offers this channel as a complementary probe.

NANOGrav spectral fit.—The matter-bounce induced GW spectrum has a universal f^2 infrared scaling, corresponding to spectral index $\gamma = 3$ in the characteristic-strain parameterization $h_c \propto f^{(3-\gamma)/2}$. *Important caveat:* the comparison that follows uses synthetic data points reconstructed from the published NANOGrav 15-year power-law fit, not the raw free-spectrum posteriors. As a consequence, the bounce-model fit achieves $\chi^2/\text{dof} = 0.012$ —a trivially good fit that reflects the synthetic data closely matching the model template by construction, not a meaningful goodness-of-fit test. The Bayes factors derived from these reconstructed data carry correspondingly reduced statistical weight and should be read as illustrative template-level preferences, not posterior-level model evidence. A proper analysis using the full posterior distributions would provide a more realistic model comparison. With that caveat, a spectral template fit to the NANOGrav 15-year synthetic data points yields an illustrative Bayes factor $B(\text{bounce}/\text{SMBH}) \approx 302$ in favor of the matter-bounce prediction ($\gamma = 3$) over the standard SMBH binary merger spectrum ($\gamma = 13/3$), and $B(\text{bounce}/\text{strings}) \approx 8.1 \times 10^4$ over cosmic strings ($\gamma = 5/3$). The reduced chi-squared values are $\chi^2/\text{dof} = 0.012$ (bounce), 1.04 (SMBH), and 1.90 (strings), with $\Delta\text{BIC} = +10.7$ and $+21.9$ for SMBH and strings respectively relative to the bounce model. The best-fit peak frequency $f_{\text{peak}} = 45.6$ nHz lies above the PTA band, plac-

ing the observed frequencies in the pure f^2 regime where the prediction is parameter-free. Within this caveat, the best-fit is consistent with matter-bounce $\gamma = 3$; a posterior-level analysis using the raw NANOGrav 15-year free-spectrum data is deferred to future work (see also the independent GPU MCMC $\gamma = 3.20 \pm 0.42$ result reported in the companion anomaly-catalog paper [56]).

D. Theoretical Research Program

The framework presented here opens several concrete lines of investigation, each constituting an independent research effort that would strengthen or constrain the model:

1. Higher-Loop and Non-Perturbative Verification

The most pressing theoretical need is a first-principles determination of α/M , which is currently a phenomenological parameter motivated by the one-loop estimate (Eq. 8). Three complementary approaches are feasible:

1. *Two-loop calculation in dimensional regularization:* Extending Sec. IV to two loops requires evaluating fermion-loop diagrams with two axial-torsion vertex insertions and one graviton vertex. The technical challenge is the γ^5 treatment in $d = 4 - \epsilon$ dimensions (the ‘t Hooft-Veltman vs. Breitenlohner-Maison schemes give different finite parts). The Nieh-Yan theorem suppresses the topological contribution at two loops, but the non-topological finite part—arising from the dressed torsion propagator—could modify the one-loop result at the 10–30% level. This calculation would also reveal whether the RG running (Eq. 9) receives threshold corrections near the QCD scale.
2. *Spin-foam vertex amplitudes:* The EPRL-FK spin-foam model provides a non-perturbative definition of the graviton propagator in LQG. Computing the parity-odd coefficient from spin-foam amplitudes in a torsionful background would constitute a fully non-perturbative verification. The key technical step is evaluating the asymptotic expansion of the vertex amplitude with an axial-current insertion, which has been partially achieved for the parity-even sector [11]. Extension to the parity-odd channel requires tracking the γ -dependent phases in the Lorentzian vertex.
3. *Lattice LQG:* Numerical evaluation of the path integral on a simplicial lattice with dynamical torsion provides a scheme-independent check. The parity-odd coefficient can be extracted from the correlation function $\langle \varepsilon^{abcd} K_{ab} R_{cd} \rangle$ measured on lattice configurations. This approach directly addresses the scheme-dependence limitation identified in Sec. XV.

Any of these approaches yielding α/M within an order of magnitude of the phenomenological best-fit value

would confirm the framework’s self-consistency; a discrepancy by more than $\mathcal{O}(10)$ would require re-evaluation of the C_ℓ^{EB} amplitude prediction while leaving the tension resolution (which depends on bounce parameters, not α/M) intact.

2. Galaxy Spin Dipole: Resolved

The galaxy spin dipole has been definitively resolved as a null result (Sec. III B), constraining $|\delta| < 0.24\%$ at 95% CL—the tightest galaxy chirality constraint to date, and $7\times$ smaller than Shamir’s claimed 3% excess. A detailed derivation of A_0 from the $\varepsilon^{abcd}K_{ab}R_{cd}$ operator is no longer needed: the framework’s > 100 order-of-magnitude underprediction is consistent with the observed null. Future work on this channel shifts to redshift-dependent parity tests and cross-correlation with spectroscopic surveys.

3. Bounce-to-Inflation Transition Dynamics

The current framework assumes that the quantum bounce connects smoothly to slow-roll inflation, with initial conditions set by the parent black hole. A detailed treatment requires:

- Numerical simulation of the transition from torsion-dominated bounce ($\rho \sim \rho_{\text{crit}}$) through reheating to standard slow-roll, tracking the vorticity ω^a and torsion T^{abc} fields throughout.
- Determination of the total inflationary e -folding number N_{tot} as a function of parent black hole mass and spin, which would eliminate the residual 10^5 fine-tuning if N_{tot} turns out to be dynamically fixed.
- Computation of ΔN_{eff} from a controlled calculation of particle production at the bounce. Independent MCMC verification (Sec. III D) constrains $\Delta N_{\text{eff}} = -0.020 \pm 0.169$ (full-tension) and $+0.065 \pm 0.17$ (Planck+BAO+SN), both consistent with zero within 1σ . The original phenomenological range 0.1–0.5 is not supported by the full posterior analysis; a non-perturbative calculation is needed to determine whether the bounce mechanism produces a detectable ΔN_{eff} signal.
- Determination of the primordial scalar power spectrum through the bounce. The modified Friedmann equation ($H^2 \propto \rho[1 - \rho/\rho_c]$) alters perturbation evolution at near-Planck densities, potentially imprinting features on the primordial power spectrum $\mathcal{P}_{\mathcal{R}}(k)$. With $N_{\text{tot}} = 92$, such features would appear at comoving scales $k \sim 10^{15} \text{ Mpc}^{-1}$, corresponding to sub-asteroid-mass horizons ($M \sim 10^{-16} M_\odot$) where primordial black hole constraints are currently weakest. Whether the spin-torsion variant produces such features, and at

what amplitude, remains an open question requiring a full perturbation calculation through the bounce that has not yet been performed.

4. Parity-Odd Primordial Gravitational Waves

[Status update: closed by perturbation-transparency result.] The parity-odd operator (Eq. 6) was initially expected to affect tensor perturbations during inflation, producing a chiral gravitational wave background. However, the perturbation-transparency result (Sec. XV F) establishes that minimal ECH gravity produces *no* tensor parity splitting: $\Delta v \equiv v_R - v_L = 0$ exactly for canonical scalar field matter, because the Holst term reduces to a topological invariant when torsion vanishes. The effective spin-torsion interaction $(J^5)^2$ is parity-even (Barrier 8), independently confirming the absence of gravitational-wave chirality. This research direction is therefore closed within the minimal ECH framework. GW chirality would require fermionic matter contributions to torsion, which are Planck-suppressed at inflationary densities ($\rho_{\text{inf}}/\rho_{\text{Pl}} \sim 10^{-12}$).

5. Black Hole Interior Numerical Relativity

Full numerical relativity simulations of rotating black hole interiors with dynamical torsion have not been attempted, primarily due to the additional degrees of freedom in the torsion field. Such simulations would:

- Test whether the smooth-bounce approximation (Eq. 10) holds for realistic rotating collapse with finite angular momentum.
- Compute the efficiency of angular momentum transfer through the bounce as a function of parent Kerr spin parameter a_* .
- Determine whether baby universe formation is generic or requires fine-tuned initial conditions.

6. Connection to Other Quantum Gravity Approaches

The parity-odd operator $\varepsilon^{abcd}K_{ab}R_{cd}$ could potentially arise in quantum gravity frameworks beyond LQG:

- In string theory, parity-violating terms in the low-energy effective action arise from compactifications with topological flux (e.g., Calabi-Yau manifolds with non-trivial torsion cohomology). Identifying whether these reproduce the structure of Eq. (6) would connect our framework to string phenomenology.
- The asymptotic safety program for quantum gravity predicts specific values for gravitational couplings at the UV fixed point. Computing the parity-odd coefficient in this framework would provide an independent

determination of α/M , potentially constraining γ from a non-LQG perspective.

E. Structural Tension: Dark Energy vs. Bounce f_{NL}

An important internal tension exists between the two main physical mechanisms in this framework, and it should be stated prominently rather than treated as a minor caveat. The inflationary suppression mechanism (Sec. II C 1) requires $N_{\text{tot}} \approx 92$ e -folds of post-bounce inflation to produce the observed dark energy scale. However, 92 e -folds would push the bounce-imprinted perturbation modes far beyond the observable horizon ($\sim e^{32}$ times larger), erasing the matter-bounce f_{NL} signature from the CMB and LSS and replacing it with the standard slow-roll value $f_{\text{NL}} \approx 0.015$. This means the geometric dark energy mechanism and the bounce f_{NL} prediction *cannot both be correct*: if the DE suppression works as modeled, the bounce signal is invisible; if the bounce signal is observable, the DE mechanism requires a different realization.

This is not a secondary technicality but rather an *open structural question* that any future development of this framework must resolve. Three possible resolutions exist: (i) the DE mechanism is incorrect and the bounce cosmology contributes only through f_{NL} and other perturbation signatures, with dark energy having a separate origin; (ii) the DE mechanism operates through a different realization that does not require $N_{\text{tot}} \approx 92$; or (iii) the matter-bounce f_{NL} prediction arises from a contracting phase that precedes the $N_{\text{tot}} \approx 92$ inflationary epoch, preserving both. Since the 14 structural barriers independently close the DE derivation routes (Sec. XI), option (i) is currently favored by the evidence: bounce cosmology and dark energy are independent problems.

F. Structural Closure

The systematic analysis presented in Secs. XI–XIII established that minimal ECH gravity is perturbation-transparent and cannot produce late-time dark energy through any standard mechanism. See those sections for the full 14-barrier catalog, the perturbation-transparency result, and the hybrid dark-energy loophole analysis.

G. Open Questions

1. *Origin of γ* : Why is $\gamma \approx 0.274$? The Barbero-Immirzi parameter is currently fixed by black hole entropy counting. A derivation from a more fundamental principle—perhaps a symmetry argument or fixed-point condition—would place the entire framework on firmer ground and potentially suggest deviations from the entropy-counting value that could be tested through our observables.

2. *UV cutoff identification*: The loop calculation (Eq. 8) contains a logarithmic dependence on Λ_{UV}/μ . We identify Λ_{UV} with the LQG area-gap mass $\sim \sqrt{\gamma} M_{\text{Pl}}$, but this choice affects the coefficient at the $\mathcal{O}(\ln)$ level. A first-principles derivation of the UV scale from spin-foam dynamics would remove this ambiguity.
3. *Dynamical dark energy evolution*: Our framework implies a strictly constant Λ_{eff} in the late universe (since $\omega^2/H_0^2 < 10^{-20}$). The question of whether the parity-odd coefficient could run with the Hubble scale remains open in principle, but subsequent analysis (see the perturbation-transparency result, Sec. XV F above) establishes that minimal ECH cannot produce distinctive late-time dynamics through scalar perturbation channels. Any connection to dynamical dark energy would require additional phenomenological freedom (e.g., CPL $w_0 w_a$ parametrization) that does not derive from the bounce physics itself.
4. *Multi-messenger parity tests*: Gravitational wave detectors could in principle test parity-odd signatures. However, the perturbation-transparency result (Sec. XV F) establishes that minimal ECH produces no tensor parity splitting ($\Delta v = v_R - v_L = 0$ exactly) for canonical scalar matter. The effective spin-torsion interaction $(J^5)^2$ is parity-even (Barrier 8). GW chirality from ECH would require fermionic matter contributions, which are Planck-suppressed at cosmological densities.
5. *Information paradox*: If the universe originated inside a black hole through a torsion-regulated bounce, does information cross the bounce? The smoothness of our bounce solution ($H^2 \rightarrow 0$ continuously) suggests information preservation, but a rigorous treatment using the Page curve or holographic entanglement entropy framework remains an open problem with implications for the black hole information paradox.
6. *Cosmic hierarchy of bounces*: If every black hole spawns a baby universe, does every universe contain black holes that spawn further universes? This recursive structure implies a cosmic hierarchy of bounces, each with potentially different values of γ (and hence different physics). Whether γ is truly universal or varies across the hierarchy is a deep question with implications for the landscape/swampland program.

XVI. CONCLUSIONS

We have investigated whether Einstein-Cartan-Holst spin-torsion gravity can produce late-time dark energy or distinctive cosmological signatures. The answer is largely negative, but the investigation yields substantive structural results and identifies the surviving science case for the matter bounce scenario within an ECH-compatible cosmology.

Central result: structural closure.—Through systematic analysis of 7 foundation studies and 17 research branches, we establish 14 independent structural barriers (Sec. XI) that close all standard routes from the nonsingular bounce to dark energy within the minimal ECH framework. The central theoretical finding is the perturbation-transparency result (Sec. XII): for canonical scalar field matter, torsion vanishes at all perturbation orders, the Holst term reduces to a topological invariant, and the Barbero-Immirzi parameter γ is invisible in all scalar and tensor perturbation observables. The dark energy scale $\Lambda_{\text{eff}} = c_\omega \omega^2 + \Xi M_{\text{Pl}}^2$ is a phenomenological parameterization, not a first-principles derivation (Sec. XV).

A structural tension.—As detailed in Sec. XVE, the DE suppression mechanism and the bounce f_{NL} prediction cannot both be correct. Since the 14 structural barriers independently close the DE derivation routes, this tension reinforces the conclusion that bounce cosmology and dark energy are independent problems.

The framework.—The parity-odd effective action is constructed from the Holst term in first-order variables (Eq. 6), with α/M treated as a phenomenological parameter. The LQC bounce at $\rho_{\text{crit}} \simeq 0.27\text{--}0.41 \rho_{\text{Pl}}$ (depending on the Barbero-Immirzi entropy-counting scheme; Sec. IIB) provides a nonsingular origin with no free parameters.

Observational context.—Planck cosmic birefringence at $2.4\text{--}2.7\sigma$ [29, 30], independently confirmed by ACT DR6 at 2.9σ [32], with combined SPIDER+Planck+ACT total rotation at $\sim 7\sigma$ [34] (subject to calibration caveats); DESI 2024–2025 dynamical dark energy evidence at $3.1\text{--}4.2\sigma$ [9, 10]; EC torsion preferred by AIC in multiple independent analyses [16, 42]. Independent Cobaya v3.6.1 verification across two frozen dataset combinations (Table II) finds ΔN_{eff} consistent with zero in both: -0.020 ± 0.169 (full-tension, 176,840 samples) and $+0.065 \pm 0.17$ (Planck+BAO+SN, 132,949 samples). Current data neither require nor exclude the predicted spin-torsion ΔN_{eff} contribution; the framework is observationally viable but not yet confirmed. The Hubble tension reduction reported in the original analysis ($4.9\sigma \rightarrow 2.9\sigma$) is driven by the SH0ES prior, not by the ΔN_{eff} extension alone.

Observational signatures.—The framework’s parity-odd operator is qualitatively consistent with the observed isotropic cosmic birefringence (Planck $2.4\text{--}2.7\sigma$, ACT DR6 2.9σ), though deriving β quantitatively from the torsion operator requires a photon-torsion coupling. The most natural one-loop coupling via the chiral anomaly triangle diagram gives $g_{a\gamma}^{\text{torsion}} \sim \alpha_{\text{EM}}^2 (\alpha/M) C_{\text{anomaly}} / (2\pi) \approx 7 \times 10^{-26} \text{ GeV}^{-1}$, which is $\sim 10^5$ times too small to produce the observed signal; the suppression arises from the α_{EM}^2 anomaly factor. The spectator ALP model (Sec. XIV C), which bypasses this suppression through a direct $\phi F \tilde{F}$ coupling, successfully accommodates the data. A Gaussian summary-likelihood consistency check (Sec. XIV C) combining Planck NPIPE and ACT DR6 measurements gives

$\beta = 0.242^\circ \pm 0.061^\circ$ (3.9σ , $\text{BF} \approx 176$); matching this requires only $f_{\text{photon}} \times C_0 = 1.73 \pm 0.44$, an $\mathcal{O}(1)$ product establishing compatibility without fine-tuning. An anisotropic low- ℓ birefringence component aligned with the cosmic rotation axis is expected but its amplitude is not yet derived. The galaxy spin dipole has phenomenological form $A(z) = A_0(1+z)^{-p}e^{-qz}$ with empirical fit $A_0 \sim 0.003$; the coupling α/M underpredicts the amplitude by ~ 125 orders of magnitude when the post-Newtonian parity-odd tidal correction, thermal spin polarization ($T/m_p \sim 10^{-13}$), and weak-field curvature are accounted for (Sec. IIC 2). No amplification mechanism is known within minimal ECH. Correlated cosmic anomaly axes are expected from the shared rotation axis.

Fine-tuning.—The inflationary suppression mechanism translates the cosmological constant hierarchy (10^{120}) into a constraint on total inflationary e -folds ($N_{\text{tot}} \approx 92$), reducing the apparent tuning to $\sim 10^5$. A 100,000-sample Monte Carlo sensitivity scan (Fig. VIIB) confirms this quantitatively: 2.2% of the physically motivated parameter space produces a viable vacuum energy, with N_{tot} identified as the single controlling parameter (Spearman $|\rho_s| = 0.996$; all other parameters $|\rho_s| < 0.08$). The viable range $N_{\text{tot}} \in [79, 95]$ is physically reasonable. This reparameterizes the fine-tuning problem as an initial-condition question (“how long did inflation last?”) rather than a fundamental-constant question (“why is Λ_{bare} so small?”), but does not solve it; the residual 10^5 tuning reflects sensitivity to N_{tot} , which is fitted, not predicted.

Falsifiability.—Explicit criteria are defined for CMB E - B spectral shape, galaxy spin dipole axis/amplitude/evolution, and cosmological parameter ranges. The model can be tested with forthcoming data from CMB-S4, LiteBIRD, Euclid, and LSST.

Known limitations.—This work does not derive the IR effective vacuum term from first principles: the $w = -1$ behavior at late times is assumed, not derived from an explicit IR effective action calculation (Sec. IIC 1); the birefringence consistency lacks a derived photon-torsion coupling (the rotation angle β cannot be predicted without one); H_0 and σ_8 values are MCMC fits within an extended parameter space, not first-principles predictions; ΔN_{eff} is a phenomenological parameter whose bounce origin is qualitative, not quantitative; Ω_k is fixed to zero as mandated by 92 e -folds of inflation; the galaxy spin amplitude A_0 is empirical, with a large order-of-magnitude gap between the α/M coupling and the observed value (Sec. IIC 2); and the cosmological MCMC uses stock CAMB with N_{eff} as a free parameter, not a bespoke spin-torsion theory module (see Data and Code Availability).

First-principles derivation status.—We have separately investigated whether the phenomenological vacuum term ρ_Λ with $w = -1$ can be derived from first principles within the minimal Einstein–Cartan–Holst–Dirac framework, and whether the framework produces distinctive observational signatures. Four routes were

tested: (i) a Nambu–Jona-Lasinio condensate mechanism exploiting the gravitationally induced four-fermion interaction, (ii) the strict one-loop fermion effective action after exact torsion elimination, (iii) promoting the Barbero-Immirzi parameter to a dynamical pseudoscalar field $\gamma \rightarrow \theta(x)$, and (iv) assessing whether the parity-odd operator maps to distinctive birefringence observables. All four yield clean negative results. The condensate route fails because the scalar/pseudoscalar channel is repulsive at $\gamma = 0.274$ and subcritical by a factor of ~ 175 even when attractive. The one-loop route fails because all Barbero-Immirzi dependence resides in the four-fermion vertex, which does not contribute at one loop. The dynamical Immirzi field reduces to a standard axion-like particle after torsion elimination, with no novel gravitational content and cosmological dynamics yielding $w = +1$ (stiff matter). The parity assessment finds no photon coupling in the minimal framework; any assumed coupling produces constant- β birefringence indistinguishable from generic axion-like-particle models. These negative results, documented in a companion technical note [27], establish that the most natural minimal-model candidates for a first-principles dark-energy mechanism fail at the tested approximation orders. The framework survives as a motivated phenomenological model; progress toward a first-principles derivation requires physics beyond the minimal model class.

This framework motivates dark energy from quantum gravitational effects in spin-torsion cosmology. While it partially reduces cosmological tensions and provides a distinctive set of correlated parity-odd signatures, the significant gaps identified above—particularly the missing photon-torsion coupling, the empirical A_0 , and the assumed $w = -1$ —must be addressed before the model can claim predictive power beyond phenomenological fitting.

Structural barriers and perturbation transparency.—Systematic analysis across 7 foundations and 17+ branches established 14 independent structural barriers that close all standard routes from the bounce to dark energy *within the minimal ECH framework* (Sec. XI). These barriers are ECH-specific: other bouncing cosmologies—notably the quintom scenario, which uses dynamical phantom and quintessence fields rather than geometric torsion—can in principle unify the bounce with late-time dark energy through mechanisms that bypass all 14 barriers. DESI DR2 evidence for equation-of-state crossing at $3.1\text{--}4.2\sigma$ (dataset-dependent [10]) lends empirical support to such models. The central result of this work is the perturbation-transparency result (Sec. XII): minimal ECH gravity is dynamically inert for scalar and tensor perturbations when the matter content is a canonical scalar field. The observable science case therefore rests on predictions of the matter bounce scenario [2], which is compatible with the ECH framework but not derived from it—principally $f_{\text{NL}} = -35/8$, testable by SPHEREx at $4\text{--}6\sigma$ significance through the multi-tracer galaxy bispectrum [46]. This prediction discriminates the matter

bounce from both slow-roll inflation ($f_{\text{NL}} \approx 0.04$) and other bouncing cosmologies such as the Cuscuton bounce ($f_{\text{NL}} \approx 0$; Ref. [47]; Table V). See the focused forecast paper [45] for the full Bayesian discrimination analysis.

Supplementary materials.—Interactive data visualizations and observational comparison charts are available at <https://bigbounce.hubify.app> [57].

Data and Code Availability

All materials necessary to reproduce the cosmological and galaxy spin results are publicly available at:

<https://github.com/Hubify-Projects/bigbounce/tree/v2.3.0/reproducibility>

(pinned to tag v2.3.0).

The repository includes:

- Four Cobaya v3.5 YAML configurations: `cobaya_planck.yaml`, `cobaya_planck_bao.yaml`, `cobaya_planck_bao_sn.yaml`, and `cobaya_full_tension.yaml`—one per dataset combination in Table III. All use stock CAMB with N_{eff} as a free parameter; no custom CAMB modifications are required.
- `galaxy_spins/spin_fit_stan.py` — Hierarchical Bayesian model (CmdStanPy) fitting the phenomenological $A(z)$ to published aggregate CW/CCW galaxy counts.
- `data_build/build_galaxy_spin_dataset.py` — Reproducible pipeline that downloads the Galaxy Zoo DECaLS catalog [58] from Zenodo (DOI: 10.5281/zenodo.4573248, CC-BY-4.0) and extracts an object-level spiral galaxy catalog. CW/CCW aggregate counts for the $A(z)$ fit are from Shamir [43].
- `docs/IMPLEMENTATION_MAP.md` — Mapping from each “This work” result to the code artifact that produces it.
- `docs/KNOWN_GAPS.md` — Honest disclosure of what cannot currently be reproduced.

What is NOT included.—MCMC chains are not pre-computed (regenerate via `reproduce_cosmology.sh`, $\sim 4\text{--}12$ h per configuration on 4 CPU cores). Bayes factors were estimated via the Savage-Dickey density ratio from MCMC posteriors, which is biased for the correlated $\Delta N_{\text{eff}}\text{--}H_0$ posterior (see the caveat in Sec. VII); dedicated nested sampling via PolyChord or MultiNest is needed for robust evidence but is not provided. No CNN galaxy classifier is included; the hierarchical fit uses published catalog labels. No CMB polarization map analysis code is provided; all birefringence values are literature citations (Sec. VI).

ACKNOWLEDGMENTS

We thank the Planck, CMB-S4, LiteBIRD, LSST, and DESI collaborations for providing the observational foundation for this work. We particularly acknowledge the foundational contributions of Nikodem Popławski, whose pioneering work on Einstein-Cartan cosmology, torsion-induced bounces, and rotating black hole universes provided essential theoretical groundwork. We thank Simone Mercuri, Laurent Freidel, Djordje Minic, and Tatsu Takeuchi for their fundamental derivations connecting the Barbero-Immirzi parameter to parity-violating interactions in Loop Quantum Gravity. We acknowledge Lior Shamir for his persistent observational documentation of large-scale galaxy spin asymmetries. We thank Yuto Minami and Eiichiro Komatsu for establishing the observational evidence for cosmic birefringence. We acknowledge the theoretical and observational cosmology communities for the frameworks and precision measurements that enable these tests. Computational resources were self-funded by the author (RunPod H200 and H100 instances).

The author acknowledges the use of Claude (Anthropic) as an AI research assistant during the systematic barrier-cataloging, perturbation-gate verification, and manuscript preparation phases of this work, specifically for literature survey, cross-paper consistency checking, LaTeX formatting, and structured-dossier maintenance. All scientific claims, derivations, numerical results, and bibliographic attributions were independently verified by the author. No external funding was received for this research.

Appendix A: Complete Parameter Summary

^aVerified values from the full-tension frozen MCMC chains (Cobaya v3.6.1, Table II). The Planck+BAO+SN dataset gives consistent results: $H_0 = 67.79 \pm 1.09$, $\sigma_8 = 0.812 \pm 0.009$, $\Delta N_{\text{eff}} = +0.065 \pm 0.17$. The original analysis reported $H_0 = 69.2 \pm 0.8$ and $\Delta N_{\text{eff}} = 0.3 \pm 0.2$, but these were driven by the SH0ES prior.

^bThe Ω_m values differ between Table II (0.308 ± 0.005 , full-tension) and this table (0.310 ± 0.008) because this table reports the average across both frozen dataset combinations (full-tension and Planck+BAO+SN, the latter giving $\Omega_m = 0.312 \pm 0.006$), rounded to reflect the broader uncertainty. The values are consistent within 1σ .

Key parameter correlations from the Fisher matrix analysis: $r(H_0, \sigma_8) = -0.45$; $r(A_0, p) = -0.62$; $r(\alpha/M, \mathcal{D}_{\text{inf}}) = -0.89$ (strong anticorrelation means only their product Ξ is well-constrained). Total χ^2 per degree of freedom: $\chi^2/\text{dof} = 1148.3/1142 = 1.006$.

Appendix B: Dimensional Analysis

The parity-odd operator (Eq. 7) has off-shell mass dimension +1, not the +4 required for a Lagrangian density. Explicitly: $[\alpha/M] = -1$ and $[\varepsilon^{\mu\nu\rho\sigma} e_\mu^I e_\nu^J \mathcal{F}_{IJ\rho\sigma}] = +2$, giving $[\mathcal{L}_{\text{odd}}] = +1$. The missing three powers of mass arise through on-shell evaluation at Planck-scale bounce densities: contorsion evaluates to $K \sim M_{\text{Pl}}$ and curvature to $R \sim M_{\text{Pl}}^2$, giving $\rho_\Lambda^{\text{bounce}} \sim (\alpha/M) M_{\text{Pl}}^3 = [(\alpha/M) M_{\text{Pl}}] M_{\text{Pl}}^4 \sim 10^{-2} M_{\text{Pl}}^4$. Inflationary dilution ($\mathcal{D}_{\text{inf}} \sim e^{-3N_{\text{tot}}}$) then yields $\rho_\Lambda = \Xi M_{\text{Pl}}^4$ with $[\rho_\Lambda] = +4$. This is a *scaling ansatz*—dimensionally correct on-shell at the bounce—not a derivation from a renormalizable off-shell EFT. A rigorous resolution would require either identifying the correctly dimensioned operator from the complete set of dimension-4 parity-odd curvature invariants, or proving that the on-shell evaluation is protected by a symmetry of the torsionful sector. No quantitative conclusion in this paper depends on resolving this gap: the 14 structural barriers (Secs. XI–XIII) close the bounce-to-dark-energy route on independent grounds. The full dimensional audit is available upon request from the author.

Appendix C: Reproducibility Materials

All MCMC results in this paper were obtained using the Cobaya framework (v3.5 for the original analysis; v3.6.1 with CAMB v1.6.5 for independent verification) [44] with stock CAMB as the theory engine. The cosmological model is Λ CDM extended by ΔN_{eff} as a free parameter; no custom CAMB modifications are required. The full reproducibility package is available at:

<https://github.com/Hubify-Projects/bigbounce/tree/v2.3.0/reproducibility>

(pinned to tag v2.3.0).
The package includes:

- Four Cobaya YAML configurations (one per dataset combination): `cobaya_planck.yaml`, `cobaya_planck_bao.yaml`, `cobaya_planck_bao_sn.yaml`, `cobaya_full_tension.yaml`.
- `reproduce_cosmology.sh` — One-command script to regenerate all MCMC chains (~ 4 – 12 h per configuration on 4 CPU cores).
- `galaxy_spins/spin_fit_stan.py` — Hierarchical Bayesian model fitting $A(z) = A_0(1+z)^{-p}e^{-qz}$ to published aggregate CW/CCW counts [43], using CmdStanPy.
- `data_build/build_galaxy_spin_dataset.py` — Downloads Galaxy Zoo DECaLS [58] from Zenodo and builds an object-level spiral galaxy catalog.

TABLE VI. Complete parameter summary with priors, verified values, and physical interpretations.

Parameter	Description	Prior	Verified Value	Notes
<i>Fundamental theory parameters</i>				
γ	Barbero-Immirzi	Fixed: 0.274	0.274 ± 0.020	LQG area spectrum
α/M	Parity-odd coeff.	Log-uniform $[10^{-40}, 10^{-20}]$	$\sim 10^{-21} \text{ GeV}^{-1}$	One-loop calculation
$\mathcal{D}_{\text{inf}}$	Inflation dilution	Derived from N_{tot}	$\sim e^{-3N_{\text{tot}}}$	$N_{\text{tot}} \sim 92$ e -folds (fitted to ρ_{Λ})
$\rho_{\text{crit}}/\rho_{\text{Pl}}$	Bounce density	Derived from γ	$0.27^{+0.07}_{-0.05}$	Modified Friedmann
Ξ	Infl. suppression factor	Derived	$\sim 10^{-123}$	$= [(\alpha/M)M_{\text{Pl}}] \times \mathcal{D}_{\text{inf}}$
<i>Galaxy spin model parameters</i>				
A_0	Amplitude at $z = 0$	Uniform $[0, 0.02]$	< 0.001 (95% CL)	Null result (Sec. III B)
p	Power-law index	Uniform $[0, 2]$	0.5 ± 0.3	Redshift evolution
q	Exponential decay	Uniform $[0, 2]$	0.5 ± 0.3	High- z suppression
$\delta^{(s)}$	Survey offset	$\mathcal{N}(0, 0.005^2)$	$\lesssim 0.002$	Instrument/PSF bias
$b^{(s)}(z)$	Selection efficiency	Fixed per survey	0.3–0.9	From simulations
<i>Standard cosmological parameters</i>				
H_0	Hubble constant	Flat $[60, 80]$	$67.68 \pm 1.06^{\text{a}}$	km/s/Mpc
σ_8	Clustering amplitude	Flat $[0.7, 0.9]$	$0.803 \pm 0.008^{\text{a}}$	
Ω_m	Matter density	Flat $[0.1, 0.5]$	$0.310 \pm 0.008^{\text{b}}$	
$\Omega_b h^2$	Baryon density	Flat $[0.019, 0.025]$	0.02237 ± 0.00015	
n_s	Scalar spectral index	Flat $[0.9, 1.1]$	0.9649 ± 0.0042	
τ	Optical depth	Flat $[0.01, 0.1]$	0.054 ± 0.007	
<i>Extended parameters (our model)</i>				
Ω_k	Curvature	Fixed: 0	—	Mandated by 92 e -folds
ΔN_{eff}	Extra species	Flat $[-1, 2]$ (via $N_{\text{eff}} \in [2.046, 5.046]$)	$-0.020 \pm 0.169^{\text{a}}$	Consistent with zero
$(\omega/H)_0$	Vorticity	$[0, 5 \times 10^{-11}]$	$< 2.1 \times 10^{-11}$	CMB isotropy (95% CL)

- `docs/IMPLEMENTATION_MAP.md` — Maps every “This work” numerical result to the code and configuration that produces it.
- `docs/KNOWN_GAPS.md` — Honest disclosure of reproducibility gaps.
- `results/mcmc_posterior_summary.txt` — Auto-generated posterior summary table (produced by `reproduce_cosmology.sh`).

Pre-computed MCMC chains are not included due to file size (~ 1 GB per run). Bayes factors (Table III) were estimated via the Savage-Dickey density ratio, which is biased for the correlated posterior (see Sec. VII caveat);

dedicated nested sampling would provide more robust estimates but is not included. The NaMaster birefringence analysis uses only 50 Monte Carlo realizations (see Sec. VI caveat). No CNN galaxy classifier or CMB polarization map analysis code is included; see the Data and Code Availability statement (Sec. XVI) for details.

Appendix D: Claims Classification

Table VII classifies every major claim in this paper as *Derived* (follows from the formalism with explicit calculation shown), *Assumed* (input assumption not derived in this work), or *Fit/Inferred* (value determined by fitting to data).

- [1] F. W. Hehl, P. von der Heyde, G. D. Kerlick, and J. M. Nester, General relativity with spin and torsion: Foundations and prospects, *Reviews of Modern Physics* **48**, 393 (1976).
- [2] Y.-F. Cai, W. Xue, R. Brandenberger, and X. Zhang, Non-gaussianity in a matter bounce, *JCAP* **0905**, 011, [arXiv:0903.0631](#).
- [3] Y.-K. Li, J. Quintin, D.-G. Wang, and Y.-F. Cai, Matter bounce cosmology with a generalized single field: non-Gaussianity and an extended no-go theorem, *JCAP* **03**, 031, [arXiv:1612.02036 \[astro-ph.CO\]](#).
- [4] Planck Collaboration, N. Aghanim, *et al.*, Planck 2018 results. VI. cosmological parameters, *Astronomy & As-*

trophysics **641**, A6 (2020), [arXiv:1807.06209 \[astro-ph.CO\]](#).

- [5] S. Weinberg, The cosmological constant problem, *Reviews of Modern Physics* **61**, 1 (1989).
- [6] A. G. Riess, W. Yuan, L. M. Macri, *et al.*, A comprehensive measurement of the local value of the Hubble constant with 1 km/s/Mpc uncertainty from the Hubble Space Telescope and the SH0ES team, *The Astrophysical Journal Letters* **934**, L7 (2022), [arXiv:2112.04510 \[astro-ph.CO\]](#).
- [7] M. Asgari *et al.*, KiDS-1000 cosmology: Cosmic shear constraints on the amplitude of matter fluctuations, *Astronomy & Astrophysics* **645**, A104 (2021),

TABLE VII. Classification of claims: Derived vs. Assumed vs. Fit/Inferred. This table is intended to prevent overclaiming and to make the epistemic status of each result transparent.

Claim	Status	Where	Notes
<i>Derived (explicit calculation shown)</i>			
LQC bounce at $\rho_{\text{crit}} \simeq 0.27\text{--}0.41 \rho_{\text{Pl}}$	Derived	Eq. (10)	From LQC; range reflects ABCK ($\gamma = 0.274$) vs. DLM ($\gamma =$
Four-fermion interaction from torsion	Derived	Eq. (4)	Standard EC result (Hehl 1976)
Parity-odd operator structure	Derived	Eq. (6)	From Holst term + fermion loop
$\mathcal{D}_{\text{inf}} = e^{-3N_{\text{tot}}} (T_{\text{reh}}/M_{\text{GUT}})^{3/2}$	Derived	Eq. (13)	Dilution of $K_{ab} \propto a^{-3}$
$C_\ell^{EB} \approx 2\beta(C_\ell^{EE} - C_\ell^{BB})$	Derived	Eq. (15)	Standard birefringence formula
$\Lambda_{\text{eff}} = \Xi M_{\text{Pl}}^2 + c_\omega \omega^2$	Derived	Eq. (12)	1 + 3 covariant decomposition
Dimensional chain: $\rho_\Lambda = \Xi M_{\text{Pl}}^4$	Derived	App. B	On-shell evaluation
<i>Assumed (input, not derived in this work)</i>			
$w = -1$ at late times	Assumed	Sec. IIC 1	IR effective action not computed
$\gamma = 0.274 \pm 0.020$	Assumed	Eq. (2)	LQG black hole entropy
Parent rotating black hole origin	Assumed	Sec. IIB	Popławski scenario
Smooth bounce $\rightarrow$ slow-roll transition	Assumed	Sec. XVD	Not numerically simulated
$A(z) = A_0(1+z)^{-p}e^{-qz}$ functional form	Assumed	Eq. (14)	Phenomenological ansatz
<i>Fit / Inferred from data</i>			
$H_0 = 69.2 \pm 0.8$ km/s/Mpc	MCMC fit	Eq. (IIIC)	Original (superseded; see Sec. IIID)
$H_0 = 67.68 \pm 1.06$ km/s/Mpc	Verification	Table II	Full-tension, frozen chains
$\sigma_8 = 0.785 \pm 0.016$	MCMC fit	Eq. (IIIC)	Original (tension dataset)
$\sigma_8 = 0.803 \pm 0.008$	Verification	Table II	Full-tension, frozen chains
$\Delta N_{\text{eff}} = -0.020 \pm 0.169$	Verification	Table II	Full-tension; consistent with zero
$\Delta N_{\text{eff}} = +0.065 \pm 0.17$	Verification	Table II	Planck+BAO+SN; consistent with zero
$\alpha/M \sim 10^{-21}$ GeV $^{-1}$	Fit	Table VI	One-loop motivates order of magnitude
$A_0 < 0.001$ (95% CL)	Upper bound	Sec. IIIB	Null result from 8.47M-galaxy chirality catalog [28]
$\beta = 0.242^\circ \pm 0.061^\circ$	Literature combination	Sec. XIV C	Summary likelihood of Planck + ACT
$f_{\text{photon}} \times C_0 = 1.73 \pm 0.44$	Consistency check	Sec. XIV C	Derived from combined β ; $\mathcal{O}(1)$
$\ln B = +4.8$ (tension dataset)	Inferred	Eq. (VII B)	Dataset-dependent; Savage-Dickey estimator, biased at $r =$

arXiv:2007.15633 [astro-ph.CO].

- [8] DES Collaboration, T. M. C. Abbott, *et al.*, Dark Energy Survey Year 3 results: Cosmological constraints from galaxy clustering and weak lensing, *Physical Review D* **105**, 023520 (2022), arXiv:2105.13549 [astro-ph.CO].
- [9] DESI Collaboration, A. G. Adame, *et al.*, DESI 2024 VI: cosmological constraints from the measurements of baryon acoustic oscillations, arXiv preprint (2024), arXiv:2404.03002 [astro-ph.CO].
- [10] DESI Collaboration, M. Abdul-Karim, *et al.*, DESI DR2 results II: Measurements of baryon acoustic oscillations and cosmological constraints, *Physical Review D* **112**, 083515 (2025), arXiv:2503.14738 [astro-ph.CO].
- [11] A. Ashtekar and P. Singh, Loop quantum cosmology: A status report, *Classical and Quantum Gravity* **28**, 213001 (2011), arXiv:1108.0893 [gr-qc].
- [12] N. J. Popławski, Cosmological constant from quarks and torsion, *Annalen der Physik* **523**, 291 (2011), arXiv:1005.0893 [gr-qc].
- [13] N. J. Popławski, Universe in a black hole in Einstein-Cartan gravity, *The Astrophysical Journal* **832**, 96 (2016), arXiv:1410.3881 [gr-qc].
- [14] N. J. Popławski, Cosmology with torsion: An alternative to cosmic inflation, *Physics Letters B* **694**, 181 (2010), arXiv:1007.0587 [astro-ph.CO].
- [15] N. J. Popławski, Four-fermion interaction from torsion as dark energy, *General Relativity and Gravitation* **44**, 491 (2012), arXiv:1102.5667 [gr-qc].
- [16] R.-Y. Liu *et al.*, Einstein-Cartan torsion and the S_8 tension, *European Physical Journal C* **85**, 112 (2025).
- [17] L. Freidel, D. Minic, and T. Takeuchi, Quantum gravity, torsion, parity violation and all that, *Physical Review D* **72**, 104002 (2005), arXiv:hep-th/0507253 [hep-th].
- [18] S. Mercuri, Fermion coupling to the Holst action, *Physical Review D* **73**, 084016 (2006), arXiv:gr-qc/0601013 [gr-qc].
- [19] S. Mercuri, Peccei-quinn mechanism in gravity and the nature of the Barbero-Immirzi parameter, *Physical Review Letters* **103**, 081302 (2009), arXiv:0902.2764 [gr-qc].
- [20] T. Fujita, K. Murai, H. Nakatsuka, and S. Tsujikawa, Detection of isotropic cosmic birefringence and its implications for axionlike particles including dark energy, *Physical Review D* **103**, 043509 (2021), arXiv:2011.11894 [astro-ph.CO].
- [21] A. Ashtekar, J. C. Baez, A. Corichi, and K. Krasnov, Quantum geometry and black hole entropy, *Physical Review Letters* **80**, 904 (1998), arXiv:gr-qc/9710007.
- [22] M. Domagała and J. Lewandowski, Black-hole entropy from quantum geometry, *Classical and Quantum Gravity* **21**, 5233 (2004), arXiv:gr-qc/0407051.
- [23] K. A. Meissner, Black-hole entropy in loop quantum gravity, *Classical and Quantum Gravity* **21**, 5245 (2004), arXiv:gr-qc/0407052.
- [24] I. L. Shapiro and P. M. Teixeira, Quantum Einstein-Cartan theory with the Holst term, *Classical and Quantum Gravity* **31**, 185002 (2014), arXiv:1402.4854 [gr-qc].
- [25] G. F. R. Ellis and M. Bruni, Covariant and gauge-invariant approach to cosmological density fluctuations, *Physical Review D* **40**, 1804 (1989).

- [26] D. Saadeh, S. M. Feeney, A. Pontzen, H. V. Peiris, and J. D. McEwen, How isotropic is the universe?, *Physical Review Letters* **117**, 131302 (2016), [arXiv:1605.07178 \[astro-ph.CO\]](#).
- [27] H. Golden, Systematic closure of minimal first-principles routes to dark energy in Einstein-Cartan-Holst gravity (2026), companion technical note, available upon request from the author.
- [28] H. Golden, 8.47 million galaxies classified for spiral handedness: A bias-audited chirality catalog from the DESI legacy imaging surveys (2026), forthcoming. Catalog and model available at <https://huggingface.co/bamfai/galaxy-chirality-v2>.
- [29] Y. Minami and E. Komatsu, New extraction of the cosmic birefringence from the Planck 2018 polarization data, *Physical Review Letters* **125**, 221301 (2020), [arXiv:2011.11254 \[astro-ph.CO\]](#).
- [30] J. R. ESKILT and E. Komatsu, Improved constraints on cosmic birefringence from the WMAP and Planck cosmic microwave background polarization data, *Physical Review D* **106**, 063503 (2022).
- [31] S. M. Carroll, Quintessence and the rest of the world: Suppressing long-range interactions, *Physical Review Letters* **81**, 3067 (1998), [arXiv:astro-ph/9806099 \[astro-ph\]](#).
- [32] P. Diego-Palazuelos and E. Komatsu, Cosmic birefringence from the Atacama Cosmology Telescope data release 6, *arXiv preprint* (2025), [arXiv:2509.13654 \[astro-ph.CO\]](#).
- [33] L. Yin, G.-H. Du, T.-N. Li, and X. Zhang, Joint constraints on cosmic birefringence and early dark energy from ACT, Planck, DESI, and PantheonPlus, *arXiv preprint* (2026), [arXiv:2601.13624 \[astro-ph.CO\]](#).
- [34] SPIDER Collaboration, Constraints on cosmic birefringence from SPIDER, Planck, and ACT observations, *arXiv e-prints* (2025), [arXiv:2510.25489 \[astro-ph.CO\]](#).
- [35] ACT Collaboration, F. J. Qu, *et al.*, The Atacama Cosmology Telescope: DR6 gravitational lensing map and cosmological parameters, *The Astrophysical Journal* **962**, 112 (2024), [arXiv:2304.05203 \[astro-ph.CO\]](#).
- [36] LiteBIRD Collaboration, E. Allys, *et al.*, Probing cosmic inflation with the LiteBIRD cosmic microwave background polarization survey, *Progress of Theoretical and Experimental Physics* **2023**, 042F01 (2023), [arXiv:2202.02773 \[astro-ph.IM\]](#).
- [37] L. Shamir, Handedness asymmetry of spiral galaxies with $z < 0.3$ shows cosmic parity violation and a dipole axis, *Physics Letters B* **715**, 25 (2012), [arXiv:1207.3674 \[astro-ph.CO\]](#).
- [38] L. Shamir, Analysis of the alignment of non-random patterns of spin directions in populations of spiral galaxies, *The Astrophysical Journal* **938**, 77 (2022).
- [39] M. J. Longo, Detection of a dipole in the handedness of spiral galaxies with redshifts $z \sim 0.04$, *Physics Letters B* **699**, 224 (2011), [arXiv:1104.2815 \[astro-ph.CO\]](#).
- [40] D. Patel and H. Desmond, A critical assessment of galaxy spin asymmetry studies, *Monthly Notices of the Royal Astronomical Society* **528**, 2553 (2024).
- [41] O. H. E. Philcox and J. Ereza, Testing cosmic parity violation with galaxy spins, *Physical Review D* **111**, 023501 (2025), [arXiv:2410.18185 \[astro-ph.CO\]](#).
- [42] T. Liu, X. Li, T. Xu, M. Biesiada, and J. Wang, Torsion cosmology in the light of DESI, supernovae and CMB observational constraints, *European Physical Journal C* (2025), [arXiv:2507.04265 \[gr-qc\]](#).
- [43] L. Shamir, Asymmetry in galaxy spin directions in JWST JADES data, *arXiv preprint* (2024), [arXiv:2401.09450 \[astro-ph.GA\]](#).
- [44] J. Torrado and A. Lewis, Cobaya: Code for Bayesian analysis of hierarchical physical models, *Journal of Cosmology and Astroparticle Physics* **05** (057), 057, [arXiv:2005.05290 \[astro-ph.IM\]](#).
- [45] H. Golden, Testing the matter bounce with primordial non-gaussianity: Forecasts for spherex and megamapper (2026), forthcoming.
- [46] C. Heinrich, O. Dore, and E. Krause, Measuring f_{nl} with the spherex multi-tracer redshift space bispectrum, (2023), [arXiv:2311.13082](#).
- [47] S. Dehghani, G. Geshnizjani, and J. Quintin, Cuscuton Bounce Beyond the Linear Regime: Bispectrum and Strong Coupling, (2025), [arXiv:2503.01992 \[gr-qc\]](#).
- [48] S. Legner, W. Handley, and W. Barker, Alleviating the Hubble tension with torsion condensation (TorC), *arXiv e-prints* (2025), [arXiv:2507.09228 \[astro-ph.CO\]](#).
- [49] S. Alam, S. Sen, and S. Sengupta, Bouncing cosmologies in modified gravity with space time torsion, *Eur. Phys. J. C* (2025), [arXiv:2509.03508 \[gr-qc\]](#).
- [50] Y.-F. Cai and J.-H. Zhu, Smoking-gun signatures of bounce cosmology from echoes of relic gravitational waves, (2026), [arXiv:2603.13924 \[astro-ph.CO\]](#).
- [51] T. Papanikolaou, S. Banerjee, Y.-F. Cai, S. Capozziello, and E. N. Saridakis, Primordial black holes and induced gravitational waves in non-singular matter bouncing cosmology, *JCAP* **06**, 066, [arXiv:2404.03779 \[gr-qc\]](#).
- [52] J. Quintin, Z. Sherkatghanad, Y.-F. Cai, and R. H. Brandenberger, Evolution of cosmological perturbations and the production of non-Gaussianities through a nonsingular bounce: Indications for a no-go theorem in single field matter bounce cosmologies, *Phys. Rev. D* **92**, 063532 (2015), [arXiv:1508.04141 \[hep-th\]](#).
- [53] H.-J. Yo and J. M. Nester, Hamiltonian analysis of Poincaré gauge theory scalar modes, *International Journal of Modern Physics D* **11**, 747 (2002), [arXiv:gr-qc/0112030](#).
- [54] R. Penrose, Singularities and time-asymmetry, in *General Relativity: An Einstein Centenary Survey*, edited by S. W. Hawking and W. Israel (Cambridge University Press, 1979) pp. 581–638.
- [55] G. 't Hooft, Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking, *NATO Sci. Ser. B* **59**, 135 (1980).
- [56] H. Golden, Multi-survey spectral anomaly detection: 319,443 uncataloged objects from 37 million sources across eight astronomical archives (2026), companion paper, forthcoming. Preprint number HUBIFY-2026-003; includes an independent GPU MCMC analysis of the NANOGrav 15-year data yielding $\gamma = 3.20 \pm 0.42$, consistent with the matter-bounce prediction $\gamma = 3$ at 0.48σ .
- [57] H. Golden, Spin-torsion cosmology and the search for geometric dark energy: Interactive data and supplementary materials (2026), interactive data visualizations, observational comparison charts, and complete paper available online.
- [58] M. Walmsley, C. Lintott, T. Gérón, *et al.*, Galaxy Zoo DECaLS: Detailed visual morphology measurements from volunteers and deep learning for 314 000 galaxies, *Monthly Notices of the Royal Astronomical Society* **509**, 3966 (2022), [arXiv:2102.08414 \[astro-ph.GA\]](#).