



RAB GTPases and their effectors in plant endosomal transport

Naoki Minamino¹ and Takashi Ueda^{1,2}

The plant endomembrane system comprises distinctive membrane-bounded organelles connected with one another by the membrane trafficking system. The RAB GTPase is a key component of the membrane trafficking machinery that regulates the targeting and tethering of trafficking vesicles to target compartments by acting as a molecular switch cycling between active and inactive states. The functions of RAB GTPases are fulfilled through their interactions with several classes of interacting factors, including guanine nucleotide exchange factors (GEFs) and effector proteins. Effector proteins for plant RAB GTPases consist of evolutionarily conserved and plant-unique factors, which are involved in various membrane trafficking events in plant cells in ways unique to plants. In this review, we summarize recent findings on the functions of endosomal RAB GTPases that underwent unique diversification during plant evolution, with a special focus on RAB5/RABF and RAB11/RABA.

Addresses

¹ Division of Cellular Dynamics, National Institute for Basic Biology, Okazaki, Aichi 444-8585, Japan

² Department of Basic Biology, SOKENDAI (Graduate University of Advanced Studies), Okazaki, Aichi 444-8585, Japan

Corresponding author: Ueda, Takashi (tueda@nibb.ac.jp)

Current Opinion in Plant Biology 2019, **52**:61–68

This review comes from a themed issue on **Cell biology**

Edited by **Eva Benkova** and **Yasin Dagdas**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 24th August 2019

<https://doi.org/10.1016/j.pbi.2019.07.007>

1369-5266/© 2019 Elsevier Ltd. All rights reserved.

Introduction

Plant cells comprise various single membrane-bounded organelles, including the endoplasmic reticulum (ER), Golgi apparatus, *trans*-Golgi network/early endosome (TGN/EE), multivesicular endosome (MVE), and vacuole. Membrane trafficking is a fundamental system for transporting proteins and lipids among these organelles and across the plasma membrane (PM), which plays pivotal roles in basic cellular activities and various higher-order physiological functions. A single round of membrane trafficking between donor and target organelles consists of four sequential processes: (1) the formation of vesicles

from a donor membrane, (2) transport of vesicles, (3) the tethering of vesicles to a target membrane, and (4) the fusion of vesicles to the target membrane. These processes involve conserved machinery components, one of which is RAB GTPase in the Ras super family [1].

RAB GTPase acts as a molecular switch in membrane trafficking in eukaryotic cells by cycling between GTP-bound active and GDP-bound inactive states. Active RAB GTPase evokes various downstream reactions, including the tethering of transport vesicles to the target membrane, organelle movement, and organelle maturation. The activation of RAB GTPase is mediated by guanine nucleotide exchange factors (GEFs), which are crucial for the spatiotemporal regulation of RAB GTPase activation and the induction of downstream reactions. These downstream reactions are triggered by nucleotide state-dependent interactions with specific factors collectively called effector proteins [1].

Arabidopsis thaliana harbors 57 RAB GTPases classified into eight groups (RABA–RABH) according to their similarity to animal RAB GTPases (RAB1/RABD, RAB2/RABB, RAB5/RABF, RAB6/RABH, RAB7/RABG, RAB8/RABE, RAB11/RABA, and RAB18/RABC) [2,3]. These eight groups are mostly conserved in green plants, with basal plants also harboring additional members, such as RAB23 [4–6]. Among the eight core groups of plant RAB GTPases, RAB5/RABF and RAB11/RABA bear distinctive features unique to plants. In addition to canonical RAB5/RABF2, the green plant lineage possesses a plant-specific type of RAB5, the ARA6/RABF1 group [7–9]. The extremely expanded nature of the RAB11/RABA group is another distinctive feature of plant RAB GTPases. 26 RAB11/RABA members are encoded in the *Arabidopsis* genome, whereas only two or three RAB11 members exist in yeast and mammalian genomes [2]. Recent studies have unveiled the molecular functions of these RAB GTPases with plant-specific characteristics and their effectors, which we summarize in this review.

RAB5/RABF-mediated endosomal trafficking

In *Arabidopsis*, canonical RAB5/RABF2 and plant-specific ARA6/RABF1 are localized to different populations of MVEs with partial overlap and were shown to fulfill distinct functions in endosomal trafficking [7,10]. Canonical RAB5/RABF2 plays critical roles in endocytic and vacuolar transport pathways in *Arabidopsis* and tobacco, and ARA6/RABF1 mediates trafficking between the MVE and the PM in *Arabidopsis* [7,11*,12–14] (Figure 1). Despite their


different functions, these RAB5/RABF members share a common activating factor, VPS9a, which mediates the exchange from GDP to GTP on these RAB5/RABF members in Arabidopsis [15] (Figure 1). The *vps9a* mutants exhibit a wide spectrum of developmental phenotypes, indicating the pivotal roles of RAB5/RABF-mediated trafficking pathways in plant development [15,16]. VPS9a is also required for normal penetration resistance against powdery mildew fungi, whereas its close homolog, VPS9b, seems to have only minor roles, if any, in development and immunity of Arabidopsis [15,17,18]. Although the mechanism by which two RAB5/RABF groups with distinct functions are regulated by the same activating factor remains obscure, it was recently shown that the C-terminal region of VPS9a is involved in the activation of ARA6/RABF1 but not for canonical RAB5/RABF2 in Arabidopsis [19]. This region could be important for the distinct regulation of two groups of plant RAB5/RABF.

Mutations in *ARA6/RABF1* and canonical *RAB5/RABF2* have been shown to have counteracting effects on the *vps9a-2* mutant [7], the molecular mechanism of which

remains unsolved. Recently, a key factor that orchestrates the two groups of plant RAB5/RABF was identified. PLANT-UNIQUE RAB5 EFFECTOR 2 (PUF2) was shown to interact with constitutively active ARA6/RABF1. PUF2 contains four coiled-coil domains, and the fourth coiled-coil region is responsible for the interaction with ARA6/RABF1. Intriguingly, subcellular localization and genetic interaction experiments indicated that PUF2 is involved in the vacuolar trafficking pathway mediated by canonical RAB5/RABF2, although PUF2 was isolated as an effector of ARA6/RABF1. Further genetic and biochemical analyses demonstrated that the titration of PUF2 with activated ARA6/RABF1 negatively affected the activation of canonical RAB5/RABF2 [20**] (Figure 1). Given that PUF2 homologs have not been detected in nonplant systems, plants developed a unique vacuolar/endosomal transport system involving plant-specific factors such as ARA6/RABF1 and PUF2 during evolution.


In animal and yeast cells, the SAND/MON1-CCZ1 complex, which is responsible for endosomal maturation from RAB5-positive early to RAB7-positive late endosomes, mediates the sequential action of RAB5 and RAB7 by acting as a RAB5 effector and a RAB7 GEF [21–23]. Recent studies revealed that Arabidopsis SAND/MON1-CCZ1 also acts as an effector of RAB5/RABF2 and a GEF for RAB7/RABG, suggesting that the molecular function of this complex is conserved among eukaryotes [24**,25**,26**]. However, Arabidopsis SAND/MON1-CCZ1 is not required for MVE formation but is essential for the fusion of MVEs with the vacuole [24**,26**]. The different requirements of this complex in plant and animal cells could reflect the distinct timing of RAB5 action. Animal RAB5 is required for early endosomal trafficking events, whereas plant canonical RAB5 mainly mediates late endosomal transport. 12S globulin, an artificial vacuolar cargo GFP-CT24, and the inositol transporter INT1 are transported through this RAB5-dependent and RAB7-dependent pathway involving SAND/MON1-CCZ1 in Arabidopsis [24**,25**,26**,27,28*] (Figure 2). In addition to this pathway, there are also vacuolar transport pathways distinctly regulated by canonical RAB5/RABF2 and RAB7/RABG in plant cells [11*,25**]. In Arabidopsis cells, there is a RAB5-dependent and RAB7-independent trafficking pathway for the vacuole. The *vps9a-2* mutation hampered the transport of the soluble *N*-ethylmaleimide sensitive factor attachment protein receptor (SNARE) SYP22 to the vacuolar membrane, although the loss of SAND or CCZ1 function did not exhibit impairment in SYP22 transport [25**] (Figure 2). This type of trafficking pathway has not been reported in animal and yeast cells, suggesting that Arabidopsis uniquely developed this pathway during evolution. Several vacuolar proteins, including a subunit of vacuolar H⁺-ATPase, VHA-a3, are reportedly transported directly from the ER to the vacuole, independent of the Golgi apparatus [29] (Figure 2). The overexpression of dominant-negative canonical RAB5/RABF2 impairs the vacuolar targeting of VHA-a3 [28*] (Figure 2). These

Figure 1



Endosomal trafficking involving the RAB5/RABF group in Arabidopsis cells. ARA6/RABF1 mediates trafficking between the MVE and the PM, whereas canonical RAB5/RABF2 acts in endocytic and vacuolar transport pathways. In the trafficking pathway from the MVE to the vacuole, ARA6/RABF1 negatively regulates canonical RAB5/RABF2 to attenuate vacuolar transport. PUF2 captures GDP-bound canonical RAB5/RABF2 and VPS9a on the MVE to promote transport from the MVE to the vacuole, and ARA6 inhibits this process by titrating PUF2 in its GTP-bound state.

Figure 2



A schematic of vacuolar trafficking pathways in Arabidopsis cells. (1) The pathway dependent on the sequential action of canonical RAB5/RABF2 and RAB7/RABG. Canonical RAB5/RABF2 is activated by VPS9a, to which the SAND/MON1-CCZ1 complex is recruited, which in turn activates RAB7/RABG. This conversion from RAB5/RABF2 to RAB7/RABG mediates maturation of the MVE from RAB5/RABF2-positive to RAB7/RABG-positive compartments. 12S globulin, GFP-CT24, and INT1 are transported through this pathway. The HOPS complex also acts in this pathway, whereas the contribution of the CORVET complex remains to be verified. (2) The pathway dependent on RAB5/RABF2 but not RAB7/RABG. SYP22 is transported through this pathway. The CORVET complex mediates membrane fusion between the MVE and the vacuole with the VAMP727-containing SNARE complex under the regulation of canonical RAB5/RABF2. (3) The Golgi-independent pathway responsible for VHA-a3 transport. Canonical RAB5/RABF2 could also function in this pathway. (4) The AP-3-dependent pathway. VAMP711, VAMP713, SUC4, and PAT10 are transported via this pathway. The HOPS complex is also required in this pathway. (5) Homotypic fusion between vacuoles mediated by HOPS, RAB7/RABG, and the VAMP71-containing SNARE complex.

lines of evidence could indicate that the canonical RAB5-dependent and RAB7-independent pathway is equivalent to the Golgi-independent pathway, otherwise they merge on the way to the vacuole.

In animal and yeast cells, the sequential action of RAB5 and RAB7 also involves two tethering complexes, the class C core vacuole/endosome tethering (CORVET) and homotypic fusion and protein sorting (HOPS) complexes. The CORVET and HOPS complexes share core subunits composed of VPS11, VPS16, VPS18, and VPS33

in addition to specific subunits (VPS3 and VPS8 for CORVET and VPS39 and VPS41 for HOPS) [30]. Arabidopsis harbors homologs of all of these subunits that are essential for embryo development and/or the growth of pollen tubes [31, 32–34, 35, 36]. Biochemical and genetic analyses revealed that CORVET and HOPS also function as effectors of canonical RAB5/RABF2 and RAB7/RABG, respectively, in plant cells [32, 35]. The *vps41-1* mutation affected the vacuolar targeting of RAB5-dependent and RAB7-dependent cargo, INT1, in the pollen tube [27, 28], which also supports the

function of HOPS in the RAB5/RABF2- and RAB7/RABG-dependent pathway (Figure 2). Conversely, the vacuolar transport of SYP22 in developing embryos was impaired in the *vps3* mutant but not in the *vps39* mutant [35^{••}], indicating that CORVET but not HOPS is involved in the RAB5-dependent and RAB7-independent pathway (Figure 2). CORVET and HOPS are also known to mediate distinct vacuolar fusion events with different sets of RAB GTPase and SNARE proteins. CORVET cooperates with RAB5/RABF2 and the VAMP727-containing SNARE complex during membrane fusion between the MVE and the vacuole. Conversely, HOPS mediates homotypic vacuolar membrane fusion with RAB7/RABG and the VAMP71-containing SNARE complex [31^{••},34,35^{••},37] (Figure 2).


In addition to these pathways, plant cells are equipped with another vacuolar transport pathway mediated by the adaptor protein complex 3 (AP-3). VAMP711, VAMP713, vacuolar sucrose transporter SUC4, and PROTEIN

S-ACYL TRANSFERASE10 (PAT10) are cargos transported via the AP-3-dependent trafficking pathway [25^{••},27,38] (Figure 2). The AP-3-dependent pathway requires HOPS activity, whereas mutants defective in canonical RAB5/RABF2 or RAB7/RABG do not exhibit defects in the transport of AP-3 pathway cargos [25^{••},27,35^{••}] (Figure 2). It has also been reported that the plant-specific effector of canonical RAB5/RABF2, ENDOSOMAL RAB EFFECTOR WITH PX-DOMAIN (EREX), is involved in the vacuolar transport of seed storage proteins [39]. These lines of evidence indicate that plants developed a unique vacuolar trafficking system by using both evolutionarily conserved and newly acquired machinery components during evolution, which underpins unique and complex plant vacuolar functions.

RAB11/RABA-mediated trafficking pathways

Twenty-six members of the Arabidopsis RAB11/RABA group are divided into RABA1–RABA6 subgroups according to their sequence similarity [2]. RAB11/RABA members

Figure 3



Schematic models of cellular events involving RAB11/RABA members. (a) Most RAB11/RABA members are partly associated with the TGN/EE. RABA5c is localized to distinctive compartments aligned at the geometric edges of Arabidopsis root cells. (b) RABA4 members and PI4Kβ1/β2 localize at the TGN/EE and tip-localized compartments and function in tip growth. (c) PI4Kβ1/β2 also localize to the cell plate, where they interact with MPK4, and function in phragmoplast organization and cell plate formation. Several RABA members are also localized to the forming cell plate.

were shown to be associated and/or partially colocalized with TGN/EE markers and to act on the secretory and endocytic pathways in Arabidopsis and tobacco cells [40,41,42*,43–48] (Figure 3a). Recently, RABA5c was shown to localize to distinct compartments from other RABA-positive domains; RABA5c localized to compartments aligned along the geometric edges of Arabidopsis cells (Figure 3a). A dominant-negative form of RABA5c did not localize to this compartment and affected cellular geometry in lateral organs, suggesting that the localization of RABA5c at the geometric edge is required for proper cell patterning in roots [49**].

Whereas effector molecules responsible for fulfilling the RABA5c function remain to be identified, some effector proteins of RABA4 members have been reported thus far. Phosphatidylinositol 4-kinases (PI4K) β 1 and PI4K β 2 act as effectors of the RAB11/RABA group member RABA4b in tip-growing cells. PI4K β 1/ β 2 localize on the TGN/EE with RABA4b and coordinate the polarized secretion of cell wall components during tip growth in root hairs [46,50**,51] (Figure 3b). The phosphatidylinositol 4-phosphate (PI4P) phosphatase ROOT HAIR DEFECTIVE 4 (RHD4) is required for the proper localization of RABA4b at the tips of root hairs [52]. These lines of evidence indicate that regulation of the PI4P level is critical for proper polarized root hair growth. PI4K β 1 also interacts with RABA4d, which is specifically expressed in pollen. The loss of RABA4d or PI4K β 1/ β 2 function resulted in defective pollen tube growth, suggesting that PI4K β 1/ β 2 could also act as an effector of RABA4d in growing pollen tubes [53*] (Figure 3b). PI4K β 1/ β 2 are also required for proper organization of the phragmoplast and cell plate formation through regulating and/or interacting with MAP65-3 and the MAP kinase MPK4 [54] (Figure 3c). Several RAB11/RABA members involved in cytokinesis are also localized to the cell plate [40,41,43,46,47] (Figure 3c). The functional interplay between RAB11/RABA and PI4K β 1/ β 2 in cell plate formation should be verified in future studies. Intriguingly, RABA1e and RABA2a on forming cell plates exhibit different sensitivities to a cytokinesis inhibitor, Endosidin 7, which may suggest that each RAB11/RABA member plays a distinctive role in cell plate formation [55].


A member of the plant U-box family of E3 ubiquitin ligases, PLANT U-BOX13 (PUB13), was recently identified as another RABA4b effector. PUB13-mediated ubiquitination was required for the ligand-induced degradation of the flagellin receptor FLAGELLIN SENSING 2 (FLS2), which is responsible for downregulating flagellin-induced immune responses [56]. PUB13 interacts with GTP-bound RABA4b and PI4P and colocalizes with RABA4b in leaf tissue cells. Intriguingly, both *pub13* and *pi4k β 1/ β 2* mutants exhibited enhanced immune responses and elevated resistance against *Pseudomonas syringae* pv. *tomato* DC3000

[56,57**]. These findings suggest that RABA4b, PI4K β 1/ β 2, and PUB13 cooperate to attenuate FLS2-dependent immune responses (Figure 4). It has also been reported that PUB13-mediated ubiquitination is involved in endocytosis of another PM-resident receptor-like kinase, BRI1 [58]. It would be interesting to see whether RABA4 and PI4K β 1/ β 2 are also involved in this process to verify the universality of the function of RABA4 in the endocytic degradation of receptor kinases. RABA1 and RABA6 are also involved in FLS2 trafficking [42*] (Figure 4). The identification of effector molecules for each group of RAB11/RABA proteins would facilitate our understanding of the detailed mechanisms of secretory/endocytic trafficking pathways in plants, which would also provide important insights into the functional diversification of RAB11/RABA members during plant evolution.

Conclusion and perspectives

We have summarized knowledge of two groups of plant endosomal RAB GTPases, RAB5/RABF and RAB11/RABA, and their specific effector molecules. Several effector proteins have been identified in the last five

Figure 4



Subcellular trafficking of FLS2 mediated by RAB11/RABA members and their effectors. RABA1 mediates the transport of newly synthesized FLS2 to the PM. FLS2 bound with flagellin is rapidly internalized from the PM and ubiquitinated by PUB13 on the TGN/EE, where RABA4 and PI4K β 1/ β 2 also reside. Ubiquitinated FLS2 is further transported to the MVE and degraded in the vacuole. RABA6 could be involved in the maturation of the MVE.

years, which improved our understanding of the molecular functions of these RAB GTPases. However, there should be more unknown effectors cooperating with these RAB GTPases in endosomal/secretory trafficking, given that animal RAB5 fulfills its functions through its interactions with dozens of effector proteins [59]. Given the plant-unique regulation of RAB5/RABF-involving and RAB11/RABA-involving trafficking pathways and unique sets of effector proteins acting there, it would be highly likely that other plant RAB groups also fulfill their functions in distinct ways from non-plant systems. To identify novel RAB-effector interactions in plants, emerging techniques such as BioID, which allows to identify weak and/or transient interactions in living cell contexts [60], in combination with widely used methods including yeast two-hybrid screening, would be effective. Meanwhile, it would also be interesting to examine how plants developed their unique trafficking systems during evolution. Comparative analyses between angiosperms including *Arabidopsis* and basal plant lineages such as chlorophytes, charophytes, and bryophytes would be efficient to reconstitute how the unique plant membrane trafficking system has been developed during evolution. Genome data from several basal plant lineages recently released [4,6,61–63] have raised new and interesting questions. For example, ARA6/RABF1 members are well conserved among land plants although its distribution is sporadic in algal species [7,8], which may reflect a critical requirement of this RAB GTPase for terrestrial life. Acquisition of a novel RAB GTPase has also been reported recently from a study of the liverwort *Marchantia polymorpha*. In addition to an orthologous protein to animal and plant RAB2/RABB, *M. polymorpha* harbors a unique RAB2/RABB protein comprising the IFT43-like domain at the C-terminus [5]. These RAB GTPases would be good models for studying how and why plants acquired plant-specific RAB GTPases. The study of *M. polymorpha* also suggested that secondary loss of ancient RAB GTPases could have contributed to specialization and/or differentiation of the plant membrane trafficking system. *M. polymorpha* harbors RAB21 and RAB23, which are lost in *Arabidopsis*, whose physiological significance should be investigated in future studies [5,64,65]. It would be also interesting to ask whether there is any relevance between the distinctive expansion of RAB11/RABA members and other expanded membrane trafficking factors such as the EXO70 group, a subunit of the exocyst complex involved in tethering of secretory vesicles to the PM [66–68] during plant evolution. Further studies would lead to exciting answers to these questions and raise more exciting new questions to address in future studies.

Conflict of interest statement

Nothing declared.

Acknowledgements

We thank Dr. Kazuo Ebine (NIBB) for critical reading of this manuscript. This study was financially supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (Grant nos. 17K19412, 18H02470, and 19H05675 (TU)).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Fujimoto M, Ueda T: **Conserved and plant-unique mechanisms regulating plant post-Golgi traffic.** *Front Plant Sci* 2012, **3**:197.
2. Rutherford S, Moore I: **The *Arabidopsis* Rab GTPase family: another enigma variation.** *Curr Opin Plant Biol* 2002, **5**:518–528.
3. Vernoud V, Horton AC, Yang Z, Nielsen E: **Analysis of the small GTPase gene superfamily of *Arabidopsis*.** *Plant Physiol* 2003, **131**:1191–1208.
4. Bowman JL, Kohchi T, Yamato KT, Jenkins J, Shu S, Ishizaki K, Yamaoka S, Nishihama R, Nakamura Y, Berger F *et al.*: **Insights into land plant evolution garnered from the *Marchantia polymorpha* genome.** *Cell* 2017, **171**:287–304 e215.
5. Minamino N, Kanazawa T, Era A, Ebine K, Nakano A, Ueda T: **RAB GTPases in the basal land plant *Marchantia polymorpha*.** *Plant Cell Physiol* 2018, **59**:845–856.
6. Rensing SA, Lang D, Zimmer AD, Terry A, Salamov A, Shapiro H, Nishiyama T, Perroud PF, Lindquist EA, Kamisugi Y *et al.*: **The *Physcomitrella* genome reveals evolutionary insights into the conquest of land by plants.** *Science* 2008, **319**:64–69.
7. Ebine K, Fujimoto M, Okatani Y, Nishiyama T, Goh T, Ito E, Dainobu T, Nishitani A, Uemura T, Sato MH *et al.*: **A membrane trafficking pathway regulated by the plant-specific RAB GTPase ARA6.** *Nat Cell Biol* 2011, **13**:853–859.
8. Hoepfner MC, Geretschlaeger A, Sommer A, Hoefftberger M, Nishiyama T, Sakayama H, Hammerl P, Tenhaken R, Ueda T, Foissner I: **Molecular and biochemical analysis of the first ARA6 homologue, a RAB5 GTPase, from green algae.** *J Exp Bot* 2013, **64**:5553–5568.
9. Ueda T, Yamaguchi M, Uchimiya H, Nakano A: **Ara6, a plant-unique novel type Rab GTPase, functions in the endocytic pathway of *Arabidopsis thaliana*.** *EMBO J* 2001, **20**:4730–4741.
10. Ueda T, Uemura T, Sato MH, Nakano A: **Functional differentiation of endosomes in *Arabidopsis* cells.** *Plant J* 2004, **40**:783–789.
11. Bottanelli F, Foresti O, Hanton S, Denecke J: **Vacuolar transport in tobacco leaf epidermis cells involves a single route for soluble cargo and multiple routes for membrane cargo.** *Plant Cell* 2011, **23**:3007–3025.
- This study demonstrated that there are multiple trafficking pathways for vacuoles in tobacco cells.
12. Bottanelli F, Gershlick DC, Denecke J: **Evidence for sequential action of Rab5 and Rab7 GTPases in prevacuolar organelle partitioning.** *Traffic* 2012, **13**:338–354.
13. Kotzer AM, Brandizzi F, Neumann U, Paris N, Moore I, Hawes C: **AtRabF2b (Ara7) acts on the vacuolar trafficking pathway in tobacco leaf epidermal cells.** *J Cell Sci* 2004, **117**:6377–6389.
14. Sohn EJ: **Rha1, an *Arabidopsis* Rab5 homolog, plays a critical role in the vacuolar trafficking of soluble cargo proteins.** *Plant Cell Online* 2003, **15**:1057–1070.
15. Goh T, Uchida W, Arakawa S, Ito E, Dainobu T, Ebine K, Takeuchi M, Sato K, Ueda T, Nakano A: **VPS9a, the common activator for two distinct types of Rab5 GTPases, is essential for the development of *Arabidopsis thaliana*.** *Plant Cell* 2007, **19**:3504–3515.

16. Inoue T, Kondo Y, Naramoto S, Nakano A, Ueda T: **RAB5 activation is required for multiple steps in *Arabidopsis thaliana* root development.** *Plant Cell Physiol* 2013, **54**:1648-1659.
17. Nielsen ME, Jurgens G, Thordal-Christensen H: **VPS9a activates the Rab5 GTPase ARA7 to confer distinct pre- and postinvasive plant innate immunity.** *Plant Cell* 2017, **29**:1927-1937.
18. Nielsen ME, Thordal-Christensen H: **Loss of VPS9b enhances vps9a-2 phenotypes.** *Plant Signal Behav* 2018, **13**:e1445950.
19. Sunada M, Goh T, Ueda T, Nakano A: **Functional analyses of the plant-specific C-terminal region of VPS9a: the activating factor for RAB5 in *Arabidopsis thaliana*.** *J Plant Res* 2016, **129**:93-102.
20. Ito E, Ebine K, Choi SW, Ichinose S, Uemura T, Nakano A, Ueda T: **Integration of two RAB5 groups during endosomal transport in plants.** *eLife* 2018, **7**.
- This study identified an effector of ARA6/RABF1, PUF2, and demonstrated that PUF2 integrates the functions of two groups of plant RAB5/RABF in endosomal transport.
21. Cabrera M, Nordmann M, Perz A, Schmedt D, Gerondopoulos A, Barr F, Piehler J, Engelbrecht-Vandre S, Ungermann C: **The Mon1-Ccz1 GEF activates the Rab7 GTPase Ypt7 via a longin-fold-Rab interface and association with PI3P-positive membranes.** *J Cell Sci* 2014, **127**:1043-1051.
22. Nordmann M, Cabrera M, Perz A, Brocker C, Ostrowicz C, Engelbrecht-Vandre S, Ungermann C: **The Mon1-Ccz1 complex is the GEF of the late endosomal Rab7 homolog Ypt7.** *Curr Biol* 2010, **20**:1654-1659.
23. Poteryaev D, Datta S, Ackema K, Zerial M, Spang A: **Identification of the switch in early-to-late endosome transition.** *Cell* 2010, **141**:497-508.
24. Cui Y, Zhao Q, Gao C, Ding Y, Zeng Y, Ueda T, Nakano A, Jiang L: **Activation of the Rab7 GTPase by the MON1-CCZ1 complex is essential for PVC-to-vacuole trafficking and plant growth in *Arabidopsis*.** *Plant Cell* 2014, **26**:2080-2097.
- The authors demonstrated that the MON1-CCZ1 complex is an effector of canonical RAB5/RABF2 and the GEF for RAB7/RABG and that it mediates endosomal maturation.
25. Ebine K, Inoue T, Ito J, Ito E, Uemura T, Goh T, Abe H, Sato K, Nakano A, Ueda T: **Plant vacuolar trafficking occurs through distinctly regulated pathways.** *Curr Biol* 2014, **24**:1375-1382.
- This study demonstrated that the SAND-CCZ1 complex is the GEF for RAB7/RABG and that canonical RAB5/RABF2 and RAB7/RABG are distinctly involved in vacuolar trafficking pathways in *Arabidopsis*.
26. Singh MK, Kruger F, Beckmann H, Brumm S, Vermeer JEM, Munnik T, Mayer U, Stierhof YD, Grefen C, Schumacher K et al.: **Protein delivery to vacuole requires SAND protein-dependent Rab GTPase conversion for MVB-vacuole fusion.** *Curr Biol* 2014, **24**:1383-1389.
- The authors showed that the SAND protein mediates the sequential action of canonical RAB5/RABF2 and RAB7/RABG to carry out MVB-vacuole fusion in *Arabidopsis*.
27. Feng QN, Song SJ, Yu SX, Wang JG, Li S, Zhang Y: **Adaptor protein-3-dependent vacuolar trafficking involves a subpopulation of COPII and HOPS tethering proteins.** *Plant Physiol* 2017, **174**:1609-1620.
28. Feng QN, Zhang Y, Li S: **Tonoplast targeting of VHA-a3 relies on a Rab5-mediated but Rab7-independent vacuolar trafficking route.** *J Integr Plant Biol* 2017, **59**:230-233.
- This study demonstrated that VHA-a3 is transported to the vacuole through the RAB5-dependent and RAB7-independent trafficking pathway.
29. Viotti C, Kruger F, Krebs M, Neubert C, Fink F, Lupanga U, Scheuring D, Boutte Y, Frescatada-Rosa M, Wolfenstetter S et al.: **The endoplasmic reticulum is the main membrane source for biogenesis of the lytic vacuole in *Arabidopsis*.** *Plant Cell* 2013, **25**:3434-3449.
30. Balderhaar HJ, Ungermann C: **CORVET and HOPS tethering complexes – coordinators of endosome and lysosome fusion.** *J Cell Sci* 2013, **126**:1307-1316.
31. Brillada C, Zheng J, Kruger F, Rovira-Diaz E, Askani JC, Schumacher K, Rojas-Pierce M: **Phosphoinositides control the localization of HOPS subunit VPS41, which together with VPS33 mediates vacuole fusion in plants.** *Proc Natl Acad Sci U S A* 2018, **115**:E8305-E8314.
- This study demonstrated that the HOPS complex is essential for the fusion and biogenesis of vacuoles in *Arabidopsis*.
32. Hao L, Liu J, Zhong S, Gu H, Qu LJ: **AtVPS41-mediated endocytic pathway is essential for pollen tube-stigma interaction in *Arabidopsis*.** *Proc Natl Acad Sci U S A* 2016, **113**:6307-6312.
33. Hicks GR, Rojo E, Hong S, Carter DG, Raikhel NV: **Geminating pollen has tubular vacuoles, displays highly dynamic vacuole biogenesis, and requires VACUOLESS1 for proper function.** *Plant Physiol* 2004, **134**:1227-1239.
34. Rojo E, Gillmor CS, Kovaleva V, Somerville CR, Raikhel NV: **VACUOLESS1 is an essential gene required for vacuole formation and morphogenesis in *Arabidopsis*.** *Dev Cell* 2001, **1**:303-310.
35. Takemoto K, Ebine K, Askani JC, Kruger F, Gonzalez ZA, Ito E, Goh T, Schumacher K, Nakano A, Ueda T: **Distinct sets of tethering complexes, SNARE complexes, and Rab GTPases mediate membrane fusion at the vacuole in *Arabidopsis*.** *Proc Natl Acad Sci U S A* 2018, **115**:E2457-E2466.
- This study showed that the CORVET and HOPS complexes in coordination with distinct sets of SNARE proteins and RAB GTPases mediate distinct membrane fusion events at the vacuole in *Arabidopsis*.
36. Tan X, Wei J, Li B, Wang M, Bao Y: **AtVps11 is essential for vacuole biogenesis in embryo and participates in pollen tube growth in *Arabidopsis*.** *Biochem Biophys Res Commun* 2017, **491**:794-799.
37. Leshem Y, Golani Y, Kaye Y, Levine A: **Reduced expression of the v-SNAREs AtVAMP71/AtVAMP7C gene family in *Arabidopsis* reduces drought tolerance by suppression of abscisic acid-dependent stomatal closure.** *J Exp Bot* 2010, **61**:2615-2622.
38. Wolfenstetter S, Wirsching P, Dotzauer D, Schneider S, Sauer N: **Routes to the tonoplast: the sorting of tonoplast transporters in *Arabidopsis* mesophyll protoplasts.** *Plant Cell* 2012, **24**:215-232.
39. Sakurai HT, Inoue T, Nakano A, Ueda T: **ENDOSOMAL RAB EFFECTOR WITH PX-DOMAIN, an interacting partner of RAB5 GTPases, regulates membrane trafficking to protein storage vacuoles in *Arabidopsis*.** *Plant Cell* 2016, **28**:1490-1503.
40. Asaoka R, Uemura T, Ito J, Fujimoto M, Ito E, Ueda T, Nakano A: ***Arabidopsis* RABA1 GTPases are involved in transport between the trans-Golgi network and the plasma membrane, and are required for salinity stress tolerance.** *Plant J* 2013, **73**:240-249.
41. Berson T, von Wangenheim D, Takac T, Samajova O, Rosero A, Ovecka M, Kornis G, Stelzer EH, Samaj J: **Trans-Golgi network localized small GTPase RabA1d is involved in cell plate formation and oscillatory root hair growth.** *BMC Plant Biol* 2014, **14**:252.
42. Choi SW, Tamaki T, Ebine K, Uemura T, Ueda T, Nakano A: **RABA members act in distinct steps of subcellular trafficking of the FLAGELLIN SENSING2 receptor.** *Plant Cell* 2013, **25**:1174-1187.
- This study reported that different RABA members are involved in distinct steps of FLS2 trafficking in tobacco cells.
43. Chow CM, Neto H, Foucart C, Moore I: **Rab-A2 and Rab-A3 GTPases define a trans-golgi endosomal membrane domain in *Arabidopsis* that contributes substantially to the cell plate.** *Plant Cell* 2008, **20**:101-123.
44. de Graaf BH, Cheung AY, Andreyeva T, Levasseur K, Kieliszewski M, Wu HM: **Rab11 GTPase-regulated membrane trafficking is crucial for tip-focused pollen tube growth in tobacco.** *Plant Cell* 2005, **17**:2564-2579.
45. Feraru E, Feraru MI, Asaoka R, Paciorek T, De Rycke R, Tanaka H, Nakano A, Friml J: **BEX5/RabA1b regulates trans-Golgi network-to-plasma membrane protein trafficking in *Arabidopsis*.** *Plant Cell* 2012, **24**:3074-3086.
46. Kang BH, Nielsen E, Preuss ML, Mastronarde D, Staehelin LA: **Electron tomography of RabA4b- and PI-4Kbeta1-labeled trans Golgi network compartments in *Arabidopsis*.** *Traffic* 2011, **12**:313-329.
47. Qi X, Zheng H: **Rab-A1c GTPase defines a population of the trans-Golgi network that is sensitive to endosidin1 during cytokinesis in *Arabidopsis*.** *Mol Plant* 2013, **6**:847-859.

48. Ueda T, Anai T, Tsukaya H, Hirata A, Uchimiya H: **Characterization and subcellular localization of a small GTP-binding protein (*Ara-4*) from *Arabidopsis*: conditional expression under control of the promoter of the gene for heat-shock protein HSP81-1.** *Mol Gen Genet* 1996, **250**:533-539.
 49. Kirchhelle C, Chow CM, Foucart C, Neto H, Stierhof YD, Kalde M, ●● Walton C, Fricker M, Smith RS, Jerusalem A *et al.*: **The specification of geometric edges by a plant Rab GTPase is an essential cell-patterning principle during organogenesis in *Arabidopsis*.** *Dev Cell* 2016, **36**:386-400.
- This study demonstrated that RABA5c localizes to the geometric edges of *Arabidopsis* root cells, where it plays critical roles in root cell patterning.
50. Preuss ML, Schmitz AJ, Thole JM, Bonner HK, Otegui MS, ●● Nielsen E: **A role for the RabA4b effector protein PI-4Kbeta1 in polarized expansion of root hair cells in *Arabidopsis thaliana*.** *J Cell Biol* 2006, **172**:991-998.
- The authors demonstrated that RABA4b and PI4Kβ1/β2 cooperate to mediate the polar growth of *Arabidopsis* root hairs.
51. Preuss ML, Serna J, Falbel TG, Bednarek SY, Nielsen E: **The *Arabidopsis* Rab GTPase RabA4b localizes to the tips of growing root hair cells.** *Plant Cell* 2004, **16**:1589-1603.
 52. Thole JM, Vermeer JE, Zhang Y, Gadella TW Jr, Nielsen E: **ROOT HAIR DEFECTIVE4 encodes a phosphatidylinositol-4-phosphate phosphatase required for proper root hair development in *Arabidopsis thaliana*.** *Plant Cell* 2008, **20**:381-395.
 53. Szumlanski AL, Nielsen E: **The Rab GTPase RabA4d regulates pollen tube tip growth in *Arabidopsis thaliana*.** *Plant Cell* 2009, **21**:526-544.
- This study revealed that RABA4d and PI4Kβ1 cooperatively mediate *Arabidopsis* pollen tube growth.
54. Lin F, Krishnamoorthy P, Schubert V, Hause G, Heilmann M, Heilmann I: **A dual role for cell plate-associated PI4Kbeta in endocytosis and phragmoplast dynamics during plant somatic cytokinesis.** *EMBO J* 2019, **38**.
 55. Davis DJ, McDowell SC, Park E, Hicks G, Wilkop TE, Drakakaki G: **The RAB GTPase RABA1e localizes to the cell plate and shows distinct subcellular behavior from RABA2a under Endosidin 7 treatment.** *Plant Signal Behav* 2015, **10**:e984520.
 56. Lu D, Lin W, Gao X, Wu S, Cheng C, Avila J, Heese A, Devarenne TP, He P, Shan L: **Direct ubiquitination of pattern recognition receptor FLS2 attenuates plant innate immunity.** *Science* 2011, **332**:1439-1442.
 57. Antignani V, Klocko AL, Bak G, Chandrasekaran SD, Dunivin T, ●● Nielsen E: **Recruitment of PLANT U-BOX13 and the PI4Kbeta1/beta2 phosphatidylinositol-4 kinases by the small GTPase RabA4B plays important roles during salicylic acid-mediated plant defense signaling in *Arabidopsis*.** *Plant Cell* 2015, **27**:243-261.

The authors showed that RAB4b and its effectors, PUB13 and PI4Kβ1/β2, are involved in FLS2 degradation in *Arabidopsis*.

58. Zhou J, Liu D, Wang P, Ma X, Lin W, Chen S, Mishev K, Lu D, Kumar R, Vanhoutte I *et al.*: **Regulation of *Arabidopsis* brassinosteroid receptor BRI1 endocytosis and degradation by plant U-box PUB12/PUB13-mediated ubiquitination.** *Proc Natl Acad Sci U S A* 2018, **115**:E1906-E1915.
59. Wandinger-Ness A, Zerial M: **Rab proteins and the compartmentalization of the endosomal system.** *Cold Spring Harb Perspect Biol* 2014, **6**:a022616.
60. Roux KJ, Kim DI, Raida M, Burke B: **A promiscuous biotin ligase fusion protein identifies proximal and interacting proteins in mammalian cells.** *J Cell Biol* 2012, **196**:801-810.
61. Banks JA, Nishiyama T, Hasebe M, Bowman JL, Gribskov M, dePamphilis C, Albert VA, Aono N, Aoyama T, Ambrose BA *et al.*: **The *Selaginella* genome identifies genetic changes associated with the evolution of vascular plants.** *Science* 2011, **332**:960-963.
62. Hori K, Maruyama F, Fujisawa T, Togashi T, Yamamoto N, Seo M, Sato S, Yamada T, Mori H, Tajima N *et al.*: ***Klebsormidium flaccidum* genome reveals primary factors for plant terrestrial adaptation.** *Nat Commun* 2014, **5**:3978.
63. Nishiyama T, Sakayama H, de Vries J, Buschmann H, Saint-Marcoux D, Ullrich KK, Haas FB, Vanderstraeten L, Becker D, Lang D *et al.*: **The *Chara* genome: secondary complexity and implications for plant terrestrialization.** *Cell* 2018, **174**:448-464 e424.
64. Elias M, Brighouse A, Gabernet-Castello C, Field MC, Dacks JB: **Sculpting the endomembrane system in deep time: high resolution phylogenetics of Rab GTPases.** *J Cell Sci* 2012, **125**:2500-2508.
65. Kloppe TH, Kienle N, Fasshauer D, Munro S: **Untangling the evolution of Rab G proteins: implications of a comprehensive genomic analysis.** *BMC Biol* 2012, **10**:71.
66. Rawat A, Brejskova L, Hala M, Cvrckova F, Zarsky V: **The *Physcomitrella patens* exocyst subunit EXO70.3d has distinct roles in growth and development, and is essential for completion of the moss life cycle.** *New Phytol* 2017, **216**:438-454.
67. Synek L, Schlager N, Elias M, Quentin M, Hauser MT, Zarsky V: **AtEXO70A1, a member of a family of putative exocyst subunits specifically expanded in land plants, is important for polar growth and plant development.** *Plant J* 2006, **48**:54-72.
68. Vukasinovic N, Zarsky V: **Tethering complexes in the *Arabidopsis* endomembrane system.** *Front Cell Dev Biol* 2016, **4**:46.