

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/301841230>

Characterizing the reproduction number of epidemics with early sub-exponential growth dynamics

Article in *Journal of The Royal Society Interface* · March 2016

DOI: 10.6084/M9.FIGSHARE.3980118

CITATIONS

15

READS

141

4 authors, including:



Gerardo Chowell

Georgia State University

361 PUBLICATIONS 8,977 CITATIONS

[SEE PROFILE](#)



Lone Simonsen

Roskilde University

318 PUBLICATIONS 17,109 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Book reviews [View project](#)



Long-term Impact of Childhood Vaccination Programs: Using Historical data to Understand Modern Disease Patterns [View project](#)

Characterizing the reproduction number of epidemics with early sub-exponential growth dynamics

Gerardo Chowell^{1,2}, Cécile Viboud², Lone Simonsen^{3,4}, Seyed M. Moghadas⁵

¹ School of Public Health, Georgia State University, Atlanta, GA, USA

² Division of International Epidemiology and Population Studies, Fogarty International Center, National Institutes of Health, Bethesda, MD, USA

³ Department of Public health, University of Copenhagen, Copenhagen, Denmark

⁴ Department of Global Health, George Washington University, Washington DC, USA

⁵ Agent Based Modelling Laboratory, York University, Toronto, Canada.

Corresponding author:

Gerardo Chowell, PhD

School of Public Health, Georgia State University, Atlanta, GA, USA

Division of International Epidemiology and Population Studies, Fogarty International Center, National Institutes of Health, Bethesda, MD, USA

Email: gchowell@gsu.edu

Body word count: 4,531

Abstract word count: 314

Abstract

Early estimates of the transmission potential of emerging and re-emerging infections are increasingly used to inform public health authorities on the level of risk posed by outbreaks. Existing methods to estimate the reproduction number generally assume exponential growth in case incidence in the first few disease generations, before susceptible depletion sets in. In reality, outbreaks can display sub-exponential (i.e., polynomial) growth in the first few disease generations, owing to clustering in contact patterns, spatial effects, inhomogeneous mixing, reactive behavior changes, or other mechanisms. Here, we introduce the *generalized* growth model to characterize the early growth profile of outbreaks and estimate the effective reproduction number, with no need for explicit assumptions about the shape of epidemic growth. We demonstrate this phenomenologic approach using analytical results and simulations from mechanistic models, and provide validation against a range of empirical disease datasets. Our results suggest that sub-exponential growth in the early phase of an epidemic is the rule rather the exception. Mechanistic simulations show that slight modifications to the classical SIR model, result in sub-exponential growth, and in turn a rapid decline in the reproduction number within 3-5 disease generations. For empirical outbreaks, the generalized-growth model consistently outperforms the exponential model for a variety of directly and indirectly transmitted diseases datasets (pandemic influenza, measles, smallpox, bubonic plague, cholera, foot-and-mouth disease, HIV/AIDS, and Ebola) with model estimates supporting sub-exponential growth dynamics. The rapid decline in effective reproduction number predicted by analytical results and observed in real and synthetic datasets within 3-5 disease generations contrasts with the expectation of invariant reproduction number in epidemics obeying exponential growth. The generalized-growth concept also provides a compelling argument for the unexpected extinction of certain emerging disease outbreaks during the early ascending phase. Overall, our approach promotes a more reliable and data-driven characterization of the early epidemic phase, which is important for accurate estimation of the reproduction number and prediction of disease impact.

Keywords: Effective reproduction number; phenomenological model; generalized-growth model; epidemics; exponential growth; sub-exponential growth.

Introduction

There is a long and successful history of using compartmental transmission models to study epidemic dynamics, often calibrated using time series data describing the progression of the epidemic [1-6]. A fundamental tenet of the classic epidemic theory is that the initial growth phase should be exponential in the absence of susceptible depletion or interventions measures. However, early sub-exponential (e.g., polynomial) growth patterns have been observed in outbreaks of HIV/AIDS [7-9], Ebola [10], and foot-and-mouth disease [11]. Potential mechanisms remain debated but include spatial heterogeneity, perhaps mediated by the route of transmission (i.e., airborne vs. close contact) [10-12], clustering of contacts [8], and reactive population behavioral changes that can gradually mitigate the transmission rate [10, 11]. Accordingly, a range of mechanistic models can reproduce sub-exponential growth dynamics before susceptible depletion sets in, including models with gradually declining contact rate over time [13] and spatially-structured models such as household-community networks [12], regular lattice static contact networks, or small-world networks with weak global coupling [13, 14]. For real epidemics however, the underlying mechanisms governing sub-exponential growth can be difficult to disentangle and hence to model [13].

Given the relatively common occurrence of sub-exponential growth dynamics in empirical data and the variety of mechanisms at play, a flexible phenomenological model has been proposed to reproduce a variety of growth profiles. In the Generalized Growth Model (GGM), a tuning parameter (called the deceleration of growth, p) can reproduce a range of dynamics from constant incidence ($p = 0$) to exponential growth ($p = 1$) [11].

Application of this generalized-growth model to empirical data supports a notably slow spread ($p < 1$) of the 2014 Ebola outbreaks at district-level in parts of West Africa, intermediate spread profiles for historical plague and smallpox outbreaks ($p = 0.8$), and near exponential dynamics for pandemic influenza ($p \approx 1$) [15]. Departure from standard epidemic theory may be more common than previously thought since transmission

heterogeneities are the rule rather than the exception [11]. In this paper, we build on the generalized-growth model concept [11, 13] to show that faithful characterization of such departures is important for accurate estimation of the reproduction number.

The basic reproduction number, commonly denoted by R_0 , is a key parameter that characterizes the early epidemic spread in a fully susceptible population, and can be used to inform public health authorities on the level of risk posed by an infectious disease and the potential effects of intervention strategies [16]. According to the classical theory of epidemics, largely based on compartmental modeling (e.g., [1, 2, 17, 18]), R_0 is expected to remain invariant during the early phase of an epidemic that grows exponentially and as long as susceptible depletion remains negligible [2]. More generally, temporal variation in the transmission potential of infectious diseases are monitored via the effective reproduction number, R_t , defined as the average number of secondary cases per primary case at calendar time t [19]. If $R_t < 1$ then the epidemic declines while $R_t > 1$ indicates widespread transmission.

Here, we expand on the generalized-growth method [11] to characterize and estimate the effective reproduction number R_t during the early growth phase and before susceptible depletion sets in. We illustrate our phenomenologic approach using a combination of analytical results and synthetic datasets derived from mechanistic models with spatial or temporal effects yielding sub-exponential growth, and apply the approach to empirical data reflecting a variety historic and contemporary outbreaks. We show that sub-exponential growth dynamics is both common and important for accurate assessment of the early transmission potential.

Materials and Methods

Our study is organized in four sections. First, we describe the phenomenological ‘generalized-growth’ model and the method to estimate the effective reproduction number in this model. Second, we simulate from this phenomenologic model to gauge the expected magnitude of temporal variation in reproduction number and test our analytic predictions.

Third we simulate epidemic datasets based on mechanistic transmission models that reproduce sub-exponential growth dynamics, and apply the generalized-growth estimation approach to these synthetic data. Fourth, we test a range of empirical outbreak datasets for presence of sub-exponential growth and estimation of the effective reproduction number.

The effective reproduction number during the early epidemic growth phase

We extend a previously established generalized-growth model (GGM) [11] to estimate the effective reproduction number R_g according to disease generations g . Briefly, the GGM is a useful phenomenological model that relaxes the assumption of exponential growth in the early ascending phase of an outbreak, taking the form:

$$C' = rC^p \quad (1)$$

where $C'(t)$ describes the incidence at time t , the solution $C(t)$ describes the cumulative number of cases at time t , r is a positive parameter denoting the growth rate (with units of (people) $^{1-p}$ per time), and $p \in [0,1]$ is a ‘deceleration of growth’ parameter (dimensionless). If $p = 0$, this equation describes constant incidence over time and the cumulative number of cases grows linearly, whereas $p = 1$ describes exponential growth in the Malthus equation and the solution is given by $C(t) = C_0 e^{rt}$, where C_0 is the initial number of cases. An equivalent approach to model sub-exponential growth would be to modulate the growth rate rather than the cumulative number of cases (see Supplementary Information).

For the exponential growth model, the average number of secondary cases generated by initial cases during the first generation interval T_g (assumed to be fixed) is estimated by [20, 21]:

$$R_0^{\text{exp}} = \frac{C'(T_g)}{C'(0)} = \frac{rC_0 e^{rT_g}}{rC_0} = e^{rT_g} \quad (2)$$

The expression for R_0^{exp} depends only on r and T_g . Moreover, during the exponential growth phase, R_0^{exp} remains invariant at e^{rT_g} . This can be shown by analyzing R_g^{exp} , the ratio of case incidences over consecutive generation intervals, which is given by:

$$R_g^{\text{exp}} = \frac{C'[(g+1)T_g]}{C'[gT_g]} = e^{r(g+1)T_g - rgT_g} = e^{rT_g} \quad (3)$$

In the case of sub-exponential growth, i.e. when $p < 1$, we can characterize the effective reproduction number R_g^{subexp} over disease generations, g . For such polynomial epidemics, equation (1) exhibits an explicit solution that describes the cumulative number of cases over time, $C_{\text{subexp}}(t)$, in the form of [22]:

$$C_{\text{subexp}}(t) = [r(1-p)t + A]^{1/(1-p)} \quad (4)$$

where $A = C_0^{1-p}$. Hence, the corresponding incidence equation is given by:

$$C'_{\text{subexp}}(t) = r[r(1-p)t + A]^{p/(1-p)} \quad (5)$$

The analytic expression for the effective reproduction number by disease generation, R_g^{subexp} , captures the ratio of case incidences over consecutive disease generations:

$$R_g^{\text{subexp}} = \frac{C'_{\text{subexp}}[(g+1)T_{g+1}]}{C'_{\text{subexp}}[gT_g]} = \left[1 + \frac{r(1-p)T_g}{r(1-p)gT_g + A} \right]^{p/(1-p)} \quad (6)$$

In contrast to the exponential growth model ($p = 1$) where R_g^{exp} was independent of disease generation throughout the early growth phase, we observe that R_g^{subexp} in generation g varies as a function of g . Given that A is fixed in Equation (6), the ratio $\frac{r(1-p)T_g}{r(1-p)gT_g + A}$ declines to zero as g increases, and thus R_g^{subexp} approaches 1.0 asymptotically. Moreover, $R_g^{\text{subexp}} \rightarrow e^{rT_g}$ as $p \rightarrow 1^-$ (see Supplementary Information).

Numerical estimation of the effective reproduction number

The effective reproduction number can be estimated from case incidence data simulated from the generalized-growth model and using information about the distribution of the disease generation interval (Table 1). Specifically, based on the incidence at calendar time t_i denoted by I_i , and the discretized probability distribution of the generation interval denoted by ρ_i , the effective reproduction number can be estimated using the renewal equation [19, 23]:

$$R_{t_i} = \frac{I_i}{\sum_{j=0}^i I_{i-j} \rho_j} \quad (7)$$

where the denominator represents the total number of cases that contribute (as primary cases) to generating the number of new cases I_i (as secondary cases) at calendar time t_i [19].

Trends in effective reproduction number based on simulations from the phenomenological ‘generalized growth’ model

To gauge the expected temporal variation in the effective reproduction number for a range of growth profiles and test analytic predictions, we simulate incidence data using the generalized-growth model (Equation 1). We fix the growth rate parameter r , but assume different distributions of the disease generation interval (e.g., exponential, gamma, uniform, delta), and vary the ‘deceleration of growth’ parameter p between 0 and 1 [11]. We analyze the outbreak trajectory in the first 5 disease generations to estimate the effective reproduction number using Equation (7) (assuming a fixed generation interval) and compare with estimates obtained from the analytic expressions in Equations 3 and 6.

Trends in early growth dynamics based on mechanistic models simulations

Next, we develop three specific examples of mechanistic transmission models that support early sub-exponential growth dynamics. These include: (1) SIR (susceptible-infectious-removed) dynamics on a spatially structured model, as substantial levels of clustering have been hypothesized to yield early polynomial epidemic growth [8, 11, 12] (2) SIR compartmental model with reactive behavioural changes *via* a time-dependent transmission rate [10, 11, 24-26]; and (3) an SIR compartmental model with inhomogeneous mixing [27, 28]. We briefly describe these models below.

SIR epidemics on a spatially structured model

One of the putative mechanisms leading to early polynomial epidemic growth dynamics is clustering [8, 11], a network property that quantifies the extent to which the contacts of one individual are also contacts of each other [14]. Contact networks are particularly useful to explore the impact of clustering; here we use a network-based transmission model with household-community structure, which has been previously applied to study the transmission dynamics of Ebola [12, 29]. In this model, individuals are organized within households of size H (each household contains H individuals) and households are organized within communities of size \bar{C} households (each community contains $\bar{C} \times H$

individuals). Network connectivity is identical for every individual. The household reproduction number R_{0H} was set at 2.0 and the community reproduction number R_{0c} was set at 0.7 based on previous study [12]. For a fixed household size ($H = 5$) and different values of the community size parameter, we analyze the temporal profile in case incidence and the effective reproduction number during the first few disease generations from 200 independent stochastic realizations.

SIR compartmental model with reactive behavioural changes

In addition to contact clustering, rapid onset of behavior changes is another mechanism that has been hypothesized to lead to sub-exponential growth dynamics, as it would result in an early decline in effective reproduction number. For instance, during the 2014-15 Ebola epidemic, some areas of West Africa exhibited early sub-exponential growth even before control interventions were put in place [10, 30].

To model behavior change, we consider a classical SIR epidemic model [1, 3] with time-dependent contact rate following:

$$\begin{aligned}\frac{dS(t)}{dt} &= -\beta(t)S \frac{I}{N} \\ \frac{dI(t)}{dt} &= \beta(t)S \frac{I}{N} - \gamma I \\ \frac{dR(t)}{dt} &= \gamma I\end{aligned}\tag{8}$$

where $S(t)$, $I(t)$, and $R(t)$ denote the number of susceptible, infectious and removed (recovered) hosts in a randomly-mixed population of size N , $\beta(t)$ is the time-dependent transmission rate, the probability that a susceptible individual encounters an infectious individual is given by $I(t)/N$, and $1/\gamma$ is the mean infectious period.

In the classical SIR model with constant transmission rate β , in a completely susceptible population, $S(0) \approx N$, and $I(t)$ grows exponentially during the early epidemic phase, e.g., $I(t) \approx I_0 e^{\gamma(R_0-1)t}$, where $R_0 = \beta/\gamma$ is the average number of secondary cases generated by a

primary case during the infectious period. When susceptible depletion kicks in ($S(t) < S(0)$), the effective reproduction number, R_t , declines following $R_t = \frac{S(t)}{N} R_0$. During the first few disease generations, where $S(0)/N \approx 1$, the classical SIR model supports a reproduction number that is nearly invariant, i.e., $R_t \approx R_0$. Here, to capture behavior change, we model an exponential decline in the transmission rate $\beta(t)$ from an initial value β_0 towards $\phi\beta_0$ at rate $q > 0$ following:

$$\beta(t) = \beta_0 \left((1 - \phi)e^{-qt} + \phi \right)$$

Here $\beta(t)$ leads to early sub-exponential growth dynamics whenever $R_0 > 1$ and $q > 0$. Assuming that $R_0 > 1$ in a sufficiently large susceptible population so that the effect of susceptible depletion is negligible in the early epidemic phase, the quantity $1 - \phi$ models the proportionate reduction in β_0 that is needed for the effective reproduction number to asymptotically reach 1.0. Hence, ϕ can be estimated as $1/R_0$. If $q = 0$, $\beta(t) = \beta_0$ and we recover the classic SIR transmission model with early exponential growth dynamics. In general, a faster decline of the effective reproduction number towards 1.0 occurs for higher values of q , even without susceptible depletion. It is worth noting that prior HIV/AIDS models (e.g., [31]) have incorporated exponential decay in the transmission rate in a similar manner as described here, albeit the rate of decay was assumed to be a time-dependent function of HIV/AIDS prevalence.

To examine the behaviour of the effective reproduction number R_g over disease generations in the above model, we analyze the temporal progression in the number of cases at generation g based on the following discrete equations [32]:

$$I_{g+1} = S_g \left(1 - e^{-R_0((1-\phi)e^{-qg}+\phi)\frac{I_g}{N}} \right) \quad (9)$$

$$S_{g+1} = S_g - I_{g+1}$$

where $\phi = \frac{1}{R_0}$, I_g is the number of new cases at generation g and S_g is the number of remaining susceptibles at generation g . We initialize simulations with $I_0 = 1$ and $S_0 = N$ where N is set to 10^8 individuals.

SIR compartmental model with nonlinear incidences

Beyond contact clustering and decay in transmission rate, a third mechanism potentially accounting for sub-exponential growth is departure from mass action, which may arise due to spatial structures or other forms of nonhomogenous mixing. These effects can be incorporated in SIR models using nonlinear incidence rates (see e.g., [33, 34]). For instance, the incidence rate can take the form: $\beta_0 S(t) I^\alpha(t) / N$ where α is a phenomenological scaling mixing parameter; $\alpha = 1$ models homogeneous mixing while $\alpha < 1$ reflects contact patterns that deviate from random mixing and lead to slower epidemic growth [35]. A related version of this model is the TSIR model [28], which has found applications in various infectious disease systems including measles [28, 36], rubella [37], and dengue [38].

Here we consider an SIR model with non-homogeneous mixing, with constant transmission rate β_0 and mixing parameter α , following:

$$\begin{aligned} \frac{dS(t)}{dt} &= -\beta_0 S \frac{I^\alpha}{N} \\ \frac{dI(t)}{dt} &= \beta_0 S \frac{I^\alpha}{N} - \gamma I \\ \frac{dR(t)}{dt} &= \gamma I \end{aligned} \tag{10}$$

To analyze the progression of the reproduction number R_g over disease generations in the above model, we use the following discrete equations describing the number of cases at generation g [13, 32]:

$$\begin{aligned} I_{g+1} &= S_g \left(1 - e^{-R_0 \frac{I_g^\alpha}{N}}\right) \\ S_{g+1} &= S_g - I_{g+1} \end{aligned} \quad (11)$$

where I_g is the number of new cases at generation g and S_g is the number of remaining susceptible individuals at generation g . We initialize simulations with $I_0 = 1$ and $S_0 = N$ where N is set to 10^8 individuals.

Application to real outbreak data

Lastly, we analyze a variety of empirical outbreak datasets to test the importance of sub-exponential growth in observed disease dynamics and the resulting impact on the effective reproduction number estimates. We rely on a convenience sample representing a variety of pathogens, geographic contexts, and time periods, and include outbreaks of pandemic influenza, measles, smallpox, bubonic plague, cholera, foot-and-mouth disease (FMD), HIV/AIDS, and Ebola (Table 1 and Supplementary Information for time series). The temporal resolution of the datasets varies from daily to annual. For each outbreak, the onset week corresponds to the first observation associated with a monotonic increase in the case incidence, up to the peak incidence.

We focus on the first 3-5 disease generations, depending on the length of the available empirical time series. We estimate the effective reproduction number using the 2-step approach. In the first step, we use nonlinear least-squares to fit the generalized growth model to the synthetic mechanistic data, and estimate parameters r and P (Equation 1, [11]). The initial number of cases C_0 is fixed according to the first observation. Nominal

95% confidence intervals for parameter estimates r and p are constructed by simulations of 200 best-fit curves $C'(t)$ using parametric bootstrap with a Poisson error structure, as in prior studies [39]. In the second step, we simulate epidemic curves using the generalized-growth model with estimated r and p , and apply equation (7) to the simulated incidence data. We assume a gamma distribution for the generation interval, with means and standard deviations as in Table 1 [40-46]. Also, for each outbreak, we compare the goodness of fit of the phenomenological generalized-growth model vs. the exponential growth models.

Results

Trends in effective reproduction number based on simulations from the phenomenological ‘generalized growth’ model

We first analyze simulations of epidemic growth under the generalized growth model in the first 5 disease generations of the outbreak, for different values of r and p and a fixed generation interval (Figure 1 and Figure S2). Our simulations confirm the analytical results described in Equations (3) and (6) in relation to changes in the effective reproduction number under early exponential ($p = 1$) and sub-exponential growth dynamics ($p < 1$).

As expected, the greater the departure from exponential growth (p close to 0), the lower the effective reproduction number R_g^{subexp} . But more importantly, in the case of sub-exponential growth, and for a given growth rate r , the effective reproduction number R_g^{subexp} is a dynamic quantity that approaches 1.0 asymptotically with increasing disease generations. In contrast, for exponential growth ($p = 1$), the effective reproduction number remains invariant during the early epidemic growth phase.

We also run simulations under different assumptions regarding the distribution of the generation interval and vary P in the range $0 < p \leq 1$ (Figure 2). The declining trend in the effective reproduction number associated with the sub-exponential growth regime ($p < 1$) persists independently of the generation interval distribution. Moreover, as P decreases, estimates of the effective reproduction number become less dependent on the

generation interval distribution (Figure 2). This indicates that for a sufficiently small $p < 1$, the mean of the generation interval distribution provides sufficient information to estimate the reproduction number, without the need to specify a full distribution.

Trends in case incidence and the effective reproduction number based on mechanistic models simulations

SIR epidemics on a spatially structured epidemic model

Figure 3 shows simulations of case incidence and the effective reproduction number R_g derived from the household-community transmission model for different levels of community mixing, tuned by \bar{C} . As expected, the lower the community mixing, the greater the departure from homogeneous mixing, and hence the greater the departure from early exponential growth dynamics. Early sub-exponential growth dynamics are observed in all community mixing scenarios tested ($\bar{C}=25, 45$ and 65 households), which is consistent with a declining trend in the effective reproduction number R_g (Figure 3).

SIR compartmental model with reactive behavioural changes

Representative profiles of R_g for the SIR model with time-dependent transmission rate $\beta(t)$ are shown in Figure 4 for different values of the speed of transmission decline, tuned by parameter q . The decline in effective reproduction number $R_g = I_{g+1} / I_g$ ($g = 0 \dots n$) is more pronounced as the decline in transmission rate is faster (i.e., $q > 0$). Early sub-exponential growth dynamics is seen in all simulations where $q > 0$ (Figure 4).

SIR compartmental model with nonlinear incidence rates

Simulations for this model display concave down incidence curves in semi-logarithmic scale, supporting the presence of early sub-exponential growth, even for values of α slightly below the homogenous mixing regime (i.e., α just below 1, Figure S3).

Accordingly, the effective reproduction number R_g exhibits a declining trend during the

first few disease generations (Figure 5). By contrast, the reproduction number remains invariant at $R_g = R_0 = 2$ when $\alpha = 1$ (Figure 5(B), black curve).

Application to real outbreak data

Lastly, we apply the concepts of sub-exponential growth dynamics to a variety of empirical outbreak datasets. Figures S4-S5 provide a comparative analysis of the goodness of fit provided by the generalized-growth and the exponential growth models across outbreaks. Our results indicate that the generalized-growth model consistently outperforms the exponential growth model in the early ascending phase of the outbreak, even when p is only slightly below 1.0 (i.e., departure from the exponential model is slight). Across outbreaks, we find variability in the deceleration of growth parameter, even for a given pathogen (median $p = 0.57$, IQR: 0.46–0.84; Figure 6). Not surprisingly, parameter uncertainty declines with increasing length of the early epidemic phase used for estimation (Figures 6-7). On the other hand, mean estimates of p (Table 1) are stable during the first 3-5 disease generations (ANOVA, P=0.9).

When we use the generalized-growth model to estimate the effective reproduction number, we find a declining trend in the effective reproduction number with increasing disease generation intervals and variability in estimates of the reproduction number across 21 outbreaks representing 8 different pathogens (Figure 7). Further, estimates of the effective reproduction number are sensitive to small changes in the deceleration of growth parameter across outbreaks (Spearman rho>0.62, P<0.002; Figure S6).

Model fits to empirical data illustrates a variety of exponential and sub-exponential growth profiles across pathogens (Figures S7-S11). For instance, the autumn 1918 influenza pandemic in San Francisco is characterized by near exponential growth, with $p \sim 0.8 - 0.9$ and a relatively stable reproduction number in the range 1.7–1.8 (Figure S7). In contrast, the foot-and-mouth disease outbreak in Uruguay at the farm level displays slower initial growth with mean $p \sim 0.4 - 0.5$ and a more variable reproduction number in the range 1.6–2.8 (Figure S8). For the HIV/AIDS epidemic in Japan (1985-2012), we estimated the mean effective reproduction number in the range 1.3–1.6 with $p \sim 0.5$.

assuming a mean generation interval of 4 years, consistent with pronounced departure from exponential growth (Figure S9).

The wealth of district-level Ebola data available for the 2014 epidemic in West Africa provides a good opportunity to gauge geographic variations in the growth profiles, and in the resulting effective reproduction numbers. Indeed, we find variability across geographic locations in the effective reproduction number (median=1.46, IQR: 1.26–1.83) and deceleration parameter p (median=0.58, IQR: 0.46–0.72), and correlation between these parameter estimates (Spearman rho=0.81, P<0.001). For comparison, at the 5th disease generation interval, the highest estimate of the effective reproduction number was at 2.5 (95% CI: 2.0 – 2.7) for the 2014 Ebola outbreak in Montserrado, Liberia while the lowest estimate was at 1.03 (95% CI: 1 – 1.1) for the outbreak in Bomi, Liberia (Figure 7).

Discussion

In this study, we introduce a quantitative ‘generalized growth’ framework to characterize the transmission potential of pathogens in the early phase of an outbreak, when susceptible depletion remains negligible, without making explicit assumptions about the epidemic growth profile. The phenomenological ‘generalized growth’ model reproduces a range of growth dynamics from polynomial to exponential [11] and is agnostic of the mechanisms affecting growth, which may include contact patterns, spatial effects, non-homogeneous mixing, and/or behavior changes. A phenomenological model can be particularly useful when biological mechanisms are difficult to identify. Using a combination of analytical results, simulations from mechanistic models, and analyses of empirical outbreak data, we demonstrate that the effective reproduction number typically displays a downward trend within the first 3-5 disease generations. Evidence of sub-exponential growth, and associated reproduction number decline, is found in disease systems as varied as Ebola, pandemic influenza, smallpox, plague, cholera, measles, foot-and-mouth disease, and HIV/AIDS. Our results indicate that the concept of sub-exponential growth is both widespread and important to consider for accurate assessment of the reproduction number.

For epidemics that truly depart from exponential growth theory, traditional estimation methods relying on the assumption of exponential growth are expected to inflate reproduction number estimates. The bias between theoretical values and estimates increases as departure from exponential growth becomes more pronounced, i.e., when p decreases towards 0, representing slower epidemic spread compared to the exponential case where $p = 1$. For instance, our estimate of the reproduction number for the 1972 smallpox epidemic in Khulna, Bangladesh (~ 2 (95% CI: 1.6–2.6)) is significantly lower than earlier historic estimates of smallpox based on an exponential growth assumption (range 3.5 – 6.0) [47]. In contrast, when p is near 1.0, indicating near exponential growth, our estimates of the reproduction number remain consistent with those of compartmental models. This is the case for the 1905 bubonic plague epidemic in Bombay, India [48] or the 1918 influenza pandemic in San Francisco [15]. Overall, our estimates for Ebola outbreaks tend to be slightly lower than those reported in prior studies, possibly because of sub-exponential growth at the district levels [12, 26, 43, 49-54]. It is also worth noting that the incorporation of generalized-growth in a phenomenological logistic-type model can substantially increase the performance of the model for short-term forecasting and prediction of the final epidemic size as recently illustrated in the context of the Zika epidemic in Colombia [55].

Here we have studied simulations from three common types of mechanistic models supporting early sub-exponential growth dynamics and incorporating characteristics of the host contact network, behavior changes, and inhomogeneous mixing. In all models, relatively small departures from crude SIR dynamics led to sub-exponential growth profiles, and in turn a quick decline in effective reproduction number, speaking to the generalizability of our findings. With real outbreak case series data however, it can be difficult to disentangle the mechanism or combination of mechanisms shaping the early epidemic growth profile, especially when case series data are limited to the early epidemic growth phase. In order to assess the contribution of different mechanisms, independent sources of data would be required to quantify the structural characteristics of the contact network [56-58], or the timing and intensity of possible behavior changes. Further, for comparison purposes, it can be particularly useful to enumerate secondary cases from

transmission tree data whenever available, and obtain independent estimates of R_0 [56-58] agnostic of any model form. There is clearly scope for more research work in this area. In the absence of detailed information on the chains of transmission or the biological mechanisms at play however, a phenomenological approach such as that proposed here with the generalized-growth model may be preferable.

In addition to providing a quantitative framework for estimation of the reproduction number based on a phenomenological approach, this study has implications for disease control, particularly our understanding of herd immunity and extinction thresholds [1, 2]. In the simple SIR models, the critical fraction of the population needed to be effectively vaccinated to prevent an epidemic is given by $1 - 1/R_0$, which is in the range 50–90% of the population for most epidemic diseases [1, 54]. However, this fraction may be potentially considerably lower for epidemics rendering sub-exponential growth, where the effective reproduction number naturally declines towards unity irrespective of other intervention measures and before susceptible depletion sets in. For example, the 2014 West-African Ebola outbreak ended with less than 1% of the population registered as cases, which defies expectations from SIR models, and the contribution of large-scale interventions on these low attack rates remains debated [59]. These data-driven observations suggest that more attention should be paid to the shape of the early ascending phase of emerging infectious diseases outbreaks, and the associated uncertainty in the reproduction number estimates should be considered.

A related consequence of sub-exponential growth dynamics, and associated decline in effective reproduction number, is the effect on the extinction threshold. Indeed, it is natural to expect a higher probability of extinction due to stochastic effects for epidemics governed by sub-exponential growth. This may in part explain the small magnitude and a short duration of most Ebola outbreaks since 1976 [60-62], as Ebola showed substantial departure from exponential growth ($0.6 < p < 0.72$). In fact, simulations using an individual-level stochastic model for Ebola with household and community contact network structure are consistent with early sub-exponential growth dynamics, and has a probability of ~40% of spontaneous die-out of an outbreak within the first month of transmission [12]. This

model [12] is also consistent with an effective reproduction number that asymptotically declines toward unity as the virus spreads through the population. From a public health perspective, outbreaks characterized by sub-exponential growth dynamics may provide a greater window of opportunity for implementation of control interventions compared to those following exponential or near exponential growth dynamics [12].

Overall, our results underscore the need to carefully characterize the shape of the epidemic growth phase in order to accurately assess early trends in reproduction number.

Consideration of the sub-exponential growth phenomenon will improve our ability to appropriately model transmission scenarios, assess the potential effects of control interventions, and provide accurate forecasts of epidemic impact. Looking to the future, the development of new mechanistic transmission models is needed to provide a better understanding of the factors shaping early epidemic growth. Such models would allow for systematic evaluation of epidemic outcomes and disease control policies. A recent review of forecasting models for the West-African Ebola epidemic highlighted a range of approaches to investigating disease spread from simple phenomenological models, to compartmental epidemic models, to intricate contact networks [63]. The vast majority of these approaches considered early exponential growth dynamics, an assumption that led to substantial overestimation of Ebola epidemic size and peak timing and intensity. In light of these findings, Chretien et al. [63] stress the need for new mechanistic models that incorporate “dampening approaches” to improve characterization of the force of infection and provide uniform forecasting approaches and evaluation metrics. We believe the present study represents a significant step in this direction.

Data Accessibility

All of the epidemic incidence data employed in this paper is being made publicly available in a supplementary file.

Competing interests

We have no competing interests.

Authors' contributions

GC designed the study. GC and SM contributed to methods. GC carried out simulations, analyzed the data, and wrote the first draft of the manuscript. GC, CV, LS and SM contributed to the writing and revisions of the manuscript, and approved its final version.

Funding

GC acknowledges financial support from the NSF grant 1414374 as part of the joint NSF-NIH-USDA Ecology and Evolution of Infectious Diseases program; UK Biotechnology and Biological Sciences Research Council grant BB/M008894/1, NSF-IIS RAPID award #1518939, and NSF grant 1318788 III: Small: Data Management for Real-Time Data Driven Epidemic simulation, and the Division of International Epidemiology and Population Studies, The Fogarty International Center, US National Institutes of Health.

CV and LS acknowledges financial support from the RAPIDD Program of the Science & Technology Directorate and the Division of International Epidemiology and Population Studies, The Fogarty International Center, US National Institutes of Health. LS also acknowledges generous support from a EC Marie Curie Horizon 2020 fellowship. SM acknowledges the support from the Natural Sciences and Engineering Research Council of Canada (NSERC), and the Mathematics of Information Technology and Complex Systems (Mitacs).

Table 1. Summary of empirical disease datasets and our corresponding estimates of the reproduction number using 4 disease generations derived using the methodology described in the main text.

Disease	Outbreak	Disease generation interval mean (S.D.) ^{&}	Time series temporal resolution	Epidemic peak timing (no. data points)	Early epidemic phase (no. data points)	Reproduction number (95% CI)	Data Source
Pandemic influenza	San Francisco (1918)	3 (1)	Days	33	14	1.67 (1.43,2.08)	[15]
Smallpox	Khulna, Bangladesh (1972)	14 (2)	Weeks	10	7	1.97 (1.61,2.62)	[64]
Plague	Bombay (1905-06)	7 (2)	Weeks	19	7	1.37 (1.19,1.61)	[65]
Measles	London (1948)	14 (2)	Weeks	16	8	1.31 (1.26,1.36)	[66]
Cholera	Aalborg (1853)	5 (1)	Days	39	20	1.87 (1.64,2.23)	[67]
FMD	Uruguay (2001)	5(1)	Days	33	10	1.77 (1.33,2.44)	[68, 69]
HIV/AIDS	Japan (1985-2012) NYC (1982-2002)	4 (1.4) 4 (1.4)	Years	24 12	16 8	1.40 (1.37,1.43) 2.39 (2.37,2.44)	[70] [71]
Ebola	Uganda (2000)	13.1 (6.6)	Weeks	8	7	1.65 (1.42,2.08)	[26, 72]
	Congo (1976)	13.1 (6.6)	Days	22	16	3.79 (2.31,6.39)	[73, 74]
	Gueckedou, Guinea (2014)	19 (11)	Weeks	22	12	1.46 (1.24,1.90)	[75]
	Montserrado, Liberia (2014)	13.1 (6.6)	Weeks	16	8	1.46 (1.24,1.93)	[75]
	Margibi, Liberia (2014)	13.1 (6.6)	Weeks	14	8	1.99 (1.52,2.86)	[75]
	Bomi, Liberia (2014)	13.1 (6.6)	Weeks	10	8	1.05 (1.00,1.20)	[75]
	Grand Bassa, Liberia (2014)	13.1 (6.6)	Weeks	7	8	1.25 (1.03,1.64)	[75]
	Western Area Urban, Sierra Leone (2014)	11.6 (5.6)	Weeks	18	8	1.26 (1.16,1.41)	[75]
	Western Area Rural, Sierra Leone (2014)	11.6 (5.6)	Weeks	19	8	1.79 (1.50,2.26)	[75]
	Bo, Sierra Leone (2014)	11.6 (5.6)	Weeks	15	8	1.50 (1.21,1.99)	[75]
Bombali, Sierra Leone (2014)	Bombali, Sierra Leone (2014)	11.6 (5.6)	Weeks	12	7	1.94 (1.53,2.14)	[75]
	Kenema, Sierra Leone (2014)	11.6 (5.6)	Weeks	12	7	1.24 (1.11,1.41)	[75]
Port Loko, Sierra Leone (2014)	Port Loko, Sierra Leone (2014)	11.6 (5.6)	Weeks	16	7	1.32 (1.18,1.50)	[75]

FMD: foot-and-mouth disease

&Mean and SD for generation interval are given in days except for HIV/AIDS datasets given in years.

Figure captions

Figure 1. Simulated profiles of epidemic growth (A) and corresponding effective reproduction number (B) based on the generalized-growth model during the first 5 disease generation intervals. The parameter p is varied from 0.2 to 1.0 while the growth rate parameter r is fixed at 0.65 per day and $C_0 = 1$. The length of the generation interval is assumed to be fixed at $T_g = 3$ days. When $p = 1$ (exponential growth), the reproduction number is constant at $R_0^{\text{exp}} = e^{rT_g} = 7.03$. When $p < 1$ (sub-exponential growth) the reproduction number gradually declines over time and approaches 1.0 asymptotically. The black circles correspond to R_g^{subexp} calculated using the analytical formula in Equation (6) while the solid lines correspond to the estimates from numerical solutions based on renewal Equation (7).

Figure 2. Simulated profiles of the reproduction number during the first 5 disease generation intervals under the generalized-growth model, for different values of p . The parameter r is fixed at 0.65 per day and $C_0 = 1$. Estimates of the reproduction number are numerically obtained using Equation (7) assuming four different distributions for the generation interval: a) uniform distribution in the range 2–4 days, b) exponential distribution with the mean of 3 days, c) gamma distribution with the mean of 3 days and variance of 1 day, and d) fixed generation interval at 3 days. When $p = 1$ (exponential growth) the reproduction number is invariant irrespective of the shape of the generation interval distribution and is given by $1 + rT_g = 2.95$ when the generation interval is exponentially distributed [20], $e^{rT_g} = 7.03$ when the generation interval is fixed [20], and $(1 + rb)^a = 5.84$ when the generation interval follows a gamma distribution with the shape and scale parameters given by $a = 9$ and $b = 1/3$ [20], respectively. For $p < 1$ (sub-exponential growth), the effective reproduction number approaches 1.0 asymptotically

over disease generations. The reproduction number R_g^{subexp} at fixed generation intervals can be explicitly computed using Equation (6) (black circles).

Figure 3. Simulations of case incidence (A, C, E) and the effective reproduction number R_g (B, D, F) for the first few consecutive disease generations, using a network-based transmission model with household-community structure [12, 29] for different values of the community mixing parameter ($\bar{C}=25, 45$, and 65 households) while keeping the household size H fixed at 5 . The household reproduction number R_{0H} was set at 2.0 and the community reproduction number R_{0c} was set at 0.7 based on a previous Ebola mathematical modeling study [12]. Each simulation started with one infectious individual. Early sub-exponential growth dynamics are observed across all scenarios, which is consistent with a declining trend in the effective reproduction number R_g . For comparison with exponential growth dynamics, the dark solid line illustrates exponential growth in case incidence over disease generations with a fixed reproduction number of 2.7 . The horizontal dashed line at $R_g=1.0$ is shown for reference.

Figure 4. Simulations of case incidence (A) and the effective reproduction number R_g (B) for the first few consecutive disease generations, using model (9) for different values of the decline rate parameter q with $R_0 = 2$ and a large population size ($N = 10^8$). Each simulation started with one infectious individual. When $q = 1$, exponential growth during the early epidemic phase is evident as a straight line fits well for several consecutive disease generations of the incidence curve in semi-logarithmic scale, and the effective reproduction number, R_g , remains invariant at 2 . A concave down curve is indicative of sub-exponential growth with $q > 0$. The horizontal dashed line at $R_g=1.0$ is shown for reference.

Figure 5. Simulations of case incidence (A) and the effective reproduction number R_g (B) for the first few consecutive disease generations, using model (11) for different values of

scaling parameter α , with $R_0 = 2$, $\gamma = 1/5$, and a large population size ($N = 10^8$).

Each simulation started with one infectious individual. When $\alpha = 1$, exponential growth during the early epidemic phase is evident as a straight line fits well for several consecutive disease generations of the incidence curve, and the effective reproduction number, R_g , remains invariant at 2. The horizontal dashed line at $R_g = 1.0$ is shown for reference.

Figure 6. Estimates of the deceleration of growth parameter and the corresponding 95% confidence intervals derived from various infectious disease outbreak datasets of case incidence series by fitting the generalized-growth model to the initial epidemic phase comprising of approximately the first 3 (green), 4 (blue), and 5 (red) disease generation intervals.

Figure 7. Estimates of the effective reproduction number and the corresponding 95% confidence intervals derived from various infectious disease outbreak datasets of case incidence series by fitting the generalized-growth model to the initial epidemic phase comprising of approximately the first 3 (green), 4 (blue), and 5 (red) disease generation intervals. The generation interval is assumed to follow a gamma distribution with the corresponding mean and variance provided in Table 1.

References

- [1] Anderson, R.M. & May, R.M. 1991 *Infectious diseases of humans*. Oxford, Oxford University Press.
- [2] Diekmann, O. & Heesterbeek, J. 2000 *Mathematical epidemiology of infectious diseases: model building, analysis and interpretation*, Wiley.
- [3] Kermack, W.O. & McKendrick, A.G. 1937 Contributions to the mathematical theory of epidemics: IV. Analysis of experimental epidemics of the virus disease mouse ectromelia. *J Hyg (Lond)* **37**, 172-187.
- [4] Ross, R. 1911 *The Prevention of Malaria*. London, John Murray.
- [5] Mollison, D. 1995 *Epidemic models: their structure and relation to data*, Cambridge University Press.
- [6] Bailey, N.T.J. 1975 *The mathematical theory of infectious disease and its applications*. New York, Hafner.
- [7] Colgate, S.A., Stanley, E.A., Hyman, J.M., Layne, S.P. & Qualls, C. 1989 Risk behavior-based model of the cubic growth of acquired immunodeficiency syndrome in the United States. *Proceedings of the National Academy of Sciences of the United States of America* **86**, 4793-4797.
- [8] Szendroi, B. & Csányi, G. 2004 Polynomial epidemics and clustering in contact networks. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **271**, S364-S366.
- [9] May, R.M. & Anderson, R.M. 1987 Transmission dynamics of HIV infection. *Nature* **326**, 137-142. (doi:10.1038/326137a0).
- [10] Chowell, G., Viboud, C., Hyman, J.M. & Simonsen, L. 2015 The Western Africa ebola virus disease epidemic exhibits both global exponential and local polynomial growth rates. *PLoS currents* **7**, (doi:10.1371/currents.outbreaks.8b55f4bad99ac5c5db3663e916803261).
- [11] Viboud, C., Simonsen, L. & Chowell, G. 2016 A generalized-growth model to characterize the early ascending phase of infectious disease outbreaks *Epidemics* **15**, 27-37.
- [12] Kiskowski, M. & Chowell, G. 2015 Modeling household and community transmission of Ebola virus disease: epidemic growth, spatial dynamics and insights for epidemic control. *Virulence* **7**, 63-73. (doi:doi:10.1080/21505594.2015.1076613).
- [13] Chowell, G., Sattenspiel, L., Bansal, S. & Viboud, C. 2016 Mathematical models to characterize early epidemic growth: A Review. *Physics of Life Reviews*. pii: S1571-0645(16)30064-1. (doi:10.1016/j.plrev.2016.07.005).
- [14] Watts, D.J. & Strogatz, S.H. 1998 Collective dynamics of 'small-world' networks. *Nature* **393**, 440-442. (doi:10.1038/30918).
- [15] Chowell, G., Nishiura, H. & Bettencourt, L.M. 2007 Comparative estimation of the reproduction number for pandemic influenza from daily case notification data. *Journal of the Royal Society, Interface / the Royal Society* **4**, 155-166. (doi:R1512W8743042865 [pii]



- 10.1098/rsif.2006.0161).
- [16] Anderson, R.M. & May, R.M. 1982 Directly transmitted infections diseases: control by vaccination. *Science* **215**, 1053-1060.
- [17] van den Driessche, P. & Watmough, J. 2002 Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Mathematical biosciences* **180**, 29-48. (doi:S0025556402001086 [pii]).
- [18] Diekmann, O., Heesterbeek, J.A. & Roberts, M.G. 2010 The construction of next-generation matrices for compartmental epidemic models. *Journal of the Royal Society, Interface / the Royal Society* **7**, 873-885. (doi:rsif.2009.0386 [pii] 10.1098/rsif.2009.0386).
- [19] Nishiura, H. & Chowell, G. 2009 The effective reproduction number as a prelude to statistical estimation of time-dependent epidemic trends. In *Mathematical and statistical estimation approaches in epidemiology* (eds. G. Chowell, J.M. Hyman, L.M.A. Bettencourt & C. Castillo-Chavez), pp. 103-121., Springer (The Netherlands).
- [20] Wallinga, J. & Lipsitch, M. 2007 How generation intervals shape the relationship between growth rates and reproductive numbers. *Proceedings. Biological sciences / The Royal Society* **274**, 599-604.
- [21] Roberts, M.G. & Heesterbeek, J.A. 2007 Model-consistent estimation of the basic reproduction number from the incidence of an emerging infection. *Journal of mathematical biology* **55**, 803-816. (doi:10.1007/s00285-007-0112-8).
- [22] Tolle, J. 2003 Can Growth Be Faster than Exponential, and Just How Slow Is the Logarithm? *The Mathematical Gazette* **87**, 522-525.
- [23] Fraser, C. 2007 Estimating individual and household reproduction numbers in an emerging epidemic. *PloS one* **2**, e758. (doi:10.1371/journal.pone.0000758).
- [24] Barbarossa, M.V., Denes, A., Kiss, G., Nakata, Y., Rost, G. & Vizi, Z. 2015 Transmission Dynamics and Final Epidemic Size of Ebola Virus Disease Outbreaks with Varying Interventions. *PloS one* **10**, e0131398. (doi:10.1371/journal.pone.0131398).
- [25] Drake, J.M., Kaul, R.B., Alexander, L.W., O'Regan, S.M., Kramer, A.M., Pulliam, J.T., Ferrari, M.J. & Park, A.W. 2015 Ebola cases and health system demand in Liberia. *PLoS Biol* **13**, e1002056. (doi:10.1371/journal.pbio.1002056).
- [26] Chowell, G., Hengartner, N.W., Castillo-Chavez, C., Fenimore, P.W. & Hyman, J.M. 2004 The basic reproductive number of Ebola and the effects of public health measures: the cases of Congo and Uganda. *Journal of theoretical biology* **229**, 119-126. (doi:10.1016/j.jtbi.2004.03.006 S0022519304001092 [pii]).
- [27] Finkenstädt, B.F. & Grenfell, B.T. 2000 Time series modelling of childhood diseases: a dynamical systems approach. *Appl. Stat.* **49**, 187-205. (doi:doi:10.1111/1467-9876.00187).
- [28] Grenfell, B.T., Bjornstad, O.N. & Finkenstadt, B.F. 2002 Dynamics of measles epidemics: Scaling noise, determinism, and predictability with the TSIR model. *Ecol Monogr* **72**, 185-202.
- [29] Kiskowski, M. 2014 Three-Scale Network Model for the Early Growth Dynamics of 2014 West Africa Ebola Epidemic. *PLOS Currents Outbreaks* doi: **10.1371/currents.outbreaks.b4690859d91684da963dc40e00f3da81**. (doi:doi:10.1371/currents.outbreaks.c6efe8274dc55274f05cbbc62bbe6070.).

- [30] Fenichel, E.P., Castillo-Chavez, C., Cuddia, M.G., Chowell, G., Parra, P.A., Hickling, G.J., Holloway, G., Horan, R., Morin, B., Perrings, C., et al. 2011 Adaptive human behavior in epidemiological models. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 6306-6311. (doi:10.1073/pnas.1011250108).
- [31] Granich, R.M., Gilks, C.F., Dye, C., De Cock, K.M. & Williams, B.G. 2009 Universal voluntary HIV testing with immediate antiretroviral therapy as a strategy for elimination of HIV transmission: a mathematical model. *Lancet* **373**, 48-57. (doi:10.1016/S0140-6736(08)61697-9).
- [32] Becker, N.G. 2015 *Modeling to inform infectious disease control*, Chapman and Hall/CRC
- [33] Liu, W.M., Hethcote, H.W. & Levin, S.A. 1987 Dynamical behavior of epidemiological models with nonlinear incidence rates. *Journal of mathematical biology* **25**, 359-380.
- [34] Liu, W.M., Levin, S.A. & Iwasa, Y. 1986 Influence of nonlinear incidence rates upon the behavior of SIRS epidemiological models. *Journal of mathematical biology* **23**, 187-204.
- [35] Severo, N. 1969 Generalizations of some stochastic epidemic models. *Mathematical biosciences* **4**, 395-402.
- [36] Ferrari, M.J., Grais, R.F., Bharti, N., Conlan, A.J., Bjornstad, O.N., Wolfson, L.J., Guerin, P.J., Djibo, A. & Grenfell, B.T. 2008 The dynamics of measles in sub-Saharan Africa. *Nature* **451**, 679-684. (doi:10.1038/nature06509).
- [37] Metcalf, C.J., Munayco, C.V., Chowell, G., Grenfell, B.T. & Bjornstad, O.N. Rubella metapopulation dynamics and importance of spatial coupling to the risk of congenital rubella syndrome in Peru. *Journal of the Royal Society, Interface / the Royal Society* **8**, 369-376. (doi:rsif.2010.0320 [pii] 10.1098/rsif.2010.0320).
- [38] Kraemer, M.U., Perkins, T.A., Cummings, D.A., Zakar, R., Hay, S.I., Smith, D.L. & Reiner, R.C., Jr. 2015 Big city, small world: density, contact rates, and transmission of dengue across Pakistan. *Journal of the Royal Society, Interface / the Royal Society* **12**, 20150468. (doi:10.1098/rsif.2015.0468).
- [39] Chowell, G., Ammon, C.E., Hengartner, N.W. & Hyman, J.M. 2006 Transmission dynamics of the great influenza pandemic of 1918 in Geneva, Switzerland: Assessing the effects of hypothetical interventions. *Journal of theoretical biology* **241**, 193-204. (doi:S0022-5193(05)00509-6 [pii] 10.1016/j.jtbi.2005.11.026).
- [40] Carrat, F., Vergu, E., Ferguson, N.M., Lemaitre, M., Cauchemez, S., Leach, S. & Valleron, A.J. 2008 Time lines of infection and disease in human influenza: a review of volunteer challenge studies. *American journal of epidemiology* **167**, 775-785. (doi:10.1093/aje/kwm375).
- [41] Fine, P.E. 2003 The interval between successive cases of an infectious disease. *American journal of epidemiology* **158**, 1039-1047.
- [42] Halloran, M.E., Longini, I.M., Jr., Nizam, A. & Yang, Y. 2002 Containing bioterrorist smallpox. *Science* **298**, 1428-1432. (doi:10.1126/science.1074674 298/5597/1428 [pii]).





- [43] Team, W.H.O.E.R. 2014 Ebola Virus Disease in West Africa - The First 9 Months of the Epidemic and Forward Projections. *The New England journal of medicine* **371**, 1481-1495. (doi:10.1056/NEJMoa1411100).
- [44] Gani, R. & Leach, S. 2004 Epidemiologic determinants for modeling pneumonic plague outbreaks. *Emerging infectious diseases* **10**, 608-614. (doi:10.3201/eid1004.030509).
- [45] Burrows, R. 1968 Excretion of foot-and-mouth disease virus prior to development of lesions. *Veterinary Record* **82**, 387.
- [46] Nishiura, H. 2010 Correcting the actual reproduction number: a simple method to estimate $R(0)$ from early epidemic growth data. *International journal of environmental research and public health* **7**, 291-302. (doi:10.3390/ijerph7010291).
- [47] Gani, R. & Leach, S. 2001 Transmission potential of smallpox in contemporary populations. *Nature* **414**, 748-751. (doi:10.1038/414748a).
- [48] Bacaer, N. 2012 The model of Kermack and McKendrick for the plague epidemic in Bombay and the type reproduction number with seasonality. *Journal of mathematical biology* **64**, 403-422. (doi:10.1007/s00285-011-0417-5).
- [49] Althaus, C.L. 2014 Estimating the reproduction number of Zaire ebolavirus (EBOV) during the 2014 outbreak in West Africa. *PLOS Currents Outbreaks. Edition 1*. doi: 10.1371/currents.outbreaks.91afb5e0f279e7f29e7056095255b288.
- [50] Nishiura, H. & Chowell, G. 2014 Early transmission dynamics of Ebola virus disease (EVD), West Africa, March to August 2014. *Euro surveillance : bulletin Europeen sur les maladies transmissibles = European communicable disease bulletin* **19**.
- [51] Towers, S., Patterson-Lomba, O. & Castillo-Chavez, C. 2014 Temporal Variations in the Effective Reproduction Number of the 2014 West Africa Ebola Outbreak. *PLOS Currents Outbreaks*.
- [52] Lewnard, J.A., Ndeffo Mbah, M.L., Alfaro-Murillo, J.A., Altice, F.L., Bawo, L., Nyenswah, T.G. & Galvani, A.P. 2014 Dynamics and control of Ebola virus transmission in Montserrado, Liberia: a mathematical modelling analysis. *The Lancet. Infectious diseases* **14**, 1189-1195. (doi:10.1016/S1473-3099(14)70995-8).
- [53] Yamin, D., Gertler, S., Ndeffo-Mbah, M.L., Skrip, L.A., Fallah, M., Nyenswah, T.G., Altice, F.L. & Galvani, A.P. 2014 Effect of Ebola Progression on Transmission and Control in Liberia. *Annals of internal medicine*. (doi:10.7326/M14-2255).
- [54] Pandey, A., Atkins, K.E., Medlock, J., Wenzel, N., Townsend, J.P., Childs, J.E., Nyenswah, T.G., Ndeffo-Mbah, M.L. & Galvani, A.P. 2014 Strategies for containing Ebola in West Africa. *Science* **346**, 991-995. (doi:10.1126/science.1260612).
- [55] Chowell, G., Hincapie-Palacio, D., Ospina, J., Pell, B., Tariq, A., Dahal, S., Moghadas, S., Smirnova, A., Simonsen, L. & Viboud, C. 2016 Using Phenomenological Models to Characterize Transmissibility and Forecast Patterns and Final Burden of Zika Epidemics. *PLOS Currents Outbreaks*
pii:ecurrents.outbreaks.f14b2217c902f453d9320a43a35b9583. doi:
10.1371/currents.outbreaks.f14b2217c902f453d9320a43a35b9583. (doi:doi:
10.1371/currents.outbreaks.f14b2217c902f453d9320a43a35b9583.).
- [56] Cleaton, J.M., Viboud, C., Simonsen, L., Hurtado, A.M. & Chowell, G. 2015 Characterizing Ebola Transmission Patterns Based on Internet News Reports. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* **62**, 24-31. (doi:10.1093/cid/civ748).

- [57] Chowell, G., Abdirizak, F., Lee, S., Lee, J., Jung, E., Nishiura, H. & Viboud, C. 2015 Transmission characteristics of MERS and SARS in the healthcare setting: a comparative study. *BMC medicine* **13**, 210. (doi:10.1186/s12916-015-0450-0).
- [58] Maganga, G.D., Kapetshi, J., Berthet, N., Kebela Ilunga, B., Kabange, F., Mbala Kingebeni, P., Mondonge, V., Muyembe, J.J., Bertherat, E., Briand, S., et al. 2014 Ebola virus disease in the Democratic Republic of Congo. *The New England journal of medicine* **371**, 2083-2091. (doi:10.1056/NEJMoa1411099).
- [59] Meltzer, M.I., Atkins, C.Y., Santibanez, S., Knust, B., Petersen, B.W., Ervin, E.D., Nichol, S.T., Damon, I.K. & Washington, M.L. 2014 Estimating the future number of cases in the ebola epidemic --- liberia and sierra leone, 2014--2015. *Morbidity and mortality weekly report. Surveillance summaries* **63**, 1-14.
- [60] House, T. 2014 Epidemiological Dynamics of Ebola Outbreaks. *eLife* **3**, e03908. (doi:10.7554/eLife.03908).
- [61] Althaus, C.L. 2015 Rapid drop in the reproduction number during the Ebola outbreak in the Democratic Republic of Congo. *PeerJ* **3**, e1418. (doi:10.7717/peerj.1418).
- [62] Ajelli, M., Parlamento, S., Bome, D., Kebbi, A., Atzori, A., Frasson, C., Putoto, G., Carraro, D. & Merler, S. 2015 The 2014 Ebola virus disease outbreak in Pujehun, Sierra Leone: epidemiology and impact of interventions. *BMC medicine* **13**, 281. (doi:10.1186/s12916-015-0524-z).
- [63] Chretien, J.P., Riley, S. & George, D.B. 2015 Mathematical modeling of the West Africa Ebola epidemic. *eLife* **4**. (doi:10.7554/eLife.09186).
- [64] Sommer, A. 1974 The 1972 smallpox outbreak in Khulna Municipality, Bangladesh. II. Effectiveness of surveillance and containment in urban epidemic control. *American journal of epidemiology* **99**, 303-313.
- [65] 1907 XXII. Epidemiological observations in Bombay City. *J Hyg (Lond)* **7**, 724-798.
- [66] Measles time-series data. Professor Ben Bolker's Personal data repository at McMaster University. Available from:
<https://ms.mcmaster.ca/~bolker/measdata.html> (
- [67] Det Kongelige Sundhedskollegiums Aarsberetning for 1853. Uddrag fra Aalborg Physikat. (
- [68] Chowell, G., Rivas, A.L., Smith, S.D. & Hyman, J.M. 2006 Identification of case clusters and counties with high infective connectivity in the 2001 epidemic of foot-and-mouth disease in Uruguay. *Am J Vet Res* **67**, 102-113. (doi:10.2460/ajvr.67.1.102).
- [69] Chowell, G., Rivas, A.L., Hengartner, N.W., Hyman, J.M. & Castillo-Chavez, C. 2006 The role of spatial mixing in the spread of foot-and-mouth disease. *Prev Vet Med* **73**, 297-314. (doi:S0167-5877(05)00275-8 [pii]
10.1016/j.prevetmed.2005.10.002).
- [70] HIV/AIDS in Japan, 2013. Infectious Agents Surveillance Report (IASR). Month Vol.35 No.9(No.415). (
- [71] CDC Wonder - AIDS Public Information Dataset U.S. Surveillance. Available from: <http://wonder.cdc.gov/aidsPublic.html> (
- [72] WHO. 2001 World Health Organization (WHO), 2001. Outbreak of Ebola hemorrhagic fever, Uganda, August 2000–January 2001. *Epidemiol. Record* **76**, 41-48.







- [73] Breman JG, Piot P, Johnson KM, White MK, Mbuyi M, Sureau P, Heymann DL, Van Nieuwenhove S, McCormick JB, Ruppol JP, et al. 1978 The epidemiology of Ebola hemorrhagic fever in Zaire, 1976. *Ebola virus haemorrhagic fever*, 103-124.
- [74] Camacho, A., Kucharski, A.J., Funk, S., Breman, J., Piot, P. & Edmunds, W.J. 2014 Potential for large outbreaks of Ebola virus disease. *Epidemics* **9**, 70-78. (doi:10.1016/j.epidem.2014.09.003).
- [75] 2015 Ebola response roadmap - Situation report - 14 October 2015. Available from: <http://apps.who.int/ebola/current-situation/ebola-situation-report-14-october-2015> (accessed on 17 October 2015). (







effective reproduction number




case incidence








Deceleration of growth parameter, ρ



Reproduction number



Supporting Information

Characterizing the reproduction number of epidemics with early sub-exponential growth dynamics¹

Gerardo Chowell^{1,2}, Cécile Viboud², Lone Simonsen^{3,4}, Seyed M. Moghadas⁵

¹ School of Public Health, Georgia State University, Atlanta, GA, USA

² Division of International Epidemiology and Population Studies, Fogarty International Center, National Institutes of Health, Bethesda, MD, USA

³ Department of Public health, University of Copenhagen, Copenhagen, Denmark

⁴ Department of Global Health, School of Public Health and Health Services, George Washington University, Washington DC, USA

⁵ Agent Based Modelling Laboratory, York University, Toronto, Canada.

¹ *Chowell, Viboud, Simonsen, Moghadas. J. R. Soc. Interface, 2016*

Alternative formulation of the generalized-growth model (GGM)

Here we derive an alternative formulation of the generalized-growth model where the biological factor acts on the growth rate r rather than C . The original generalized-growth model is given by [1]:

$$C' = rC^p \quad (2.1)$$

where $C'(t)$ describes the incidence curve over time t , the solution $C(t)$ describes the cumulative number of cases at time t , r is a positive parameter denoting the growth rate (with the units of (people) $^{1-p}$ per time), and $p \in [0,1]$ is a ‘deceleration of growth’ parameter (dimensionless).

We first apply a change of variable $u = \log(C)$ so that $C = e^u$. Then the model becomes:

$$u' = re^{u(p-1)}$$

The solution of this model is given by:

$$u(t) = \begin{cases} rt & p=1 \\ \frac{1}{1-p} \log(r(1-p)t + e^{u_0(1-p)}) & 0 \leq p < 1 \end{cases}$$

where $u_0 = \log(C_0)$. The rate of change of the growth rate in the original generalized-growth model (Equation 1) is given by:

$$u'(t) = \begin{cases} r & p=1 \\ \frac{r}{r(1-p)t + e^{u_0(1-p)}} & 0 \leq p < 1 \end{cases}$$


Hence, the alternative formulation of the generalized-growth model is given by:

$$C'(t) = u'(t)C(t)$$

$$\text{where } u'(t) = \begin{cases} \frac{r}{r(1-p)t + e^{\log(C_0)(1-p)}} & 0 \leq p < 1 \\ r & p=1 \end{cases}$$

The same numerical results are obtained from both formulations of the generalized growth model as shown in Figure S1.

Figure S1. Comparison of the numerical results derived from both formulations of the generalized-growth model for different values of p when $r = 1.5$ and $C_0 = 1$.



Mathematical proof for $R_g^{\text{subexp}} \rightarrow e^{rT_g}$ as $p \rightarrow 1^-$

Using methods of limit, one can easily show that the reproduction number R_g^{subexp} according to disease generations converges to e^{rT_g} as $p \rightarrow 1^-$ as follows:

$$R_g^{\text{subexp}} = \left[1 + \frac{r(1-p)T_g}{r(1-p)gT_g + C_0^{1-p}} \right]^{\frac{p}{1-p}} \quad (2.6)$$

If $p \rightarrow 1^-$ then $R_g^{\text{subexp}} \approx 1^\infty$

Taking the logarithm on both sides we obtain (when $C_0 \neq 0$):

$$\ln(R_g^{\text{subexp}}) = \frac{p}{1-p} \ln \left[1 + \frac{r(1-p)T_g}{r(1-p)gT_g + C_0^{1-p}} \right] \approx \frac{0}{0} \text{ when } p \rightarrow 1^-$$

Applying L'Hospital's Rule, we obtain:

$$y = \frac{1 \left[\frac{-rT_g[1] - \frac{0}{1^2}}{1+0} \right]^0}{-1} = \frac{-rT_g}{-1} = rT_g$$

as $p \rightarrow 1^-$. Then $\lim_{p \rightarrow 1^-} R_g^{\text{subexp}} = e^{rT_g}$

Putting everything together, we have:

$$R_g^{\text{subexp}} = \begin{cases} 1 & \text{If } g \rightarrow \infty \text{ and } p < 1 \\ e^{rT_g} & \text{If } p \rightarrow 1^- \end{cases}$$

Figure S2. Simulated profiles of the effective reproduction number during the first 5 generation intervals derived from case incidence curves of the generalized-growth model with different values of the growth rate (r) and the deceleration of growth parameter (p). The initial number of cases is set to $C(0)=1$. Estimates of the effective reproduction number are generated assuming a fixed generation interval at 3 days.




Figure S3. Simulations of the early epidemic growth phase derived from the SIR model described by Equations (2.10) for different values of the power-law scaling parameter α , $\gamma = \frac{1}{5}$, and (A) $\beta_0 = 0.4$, (B) $\beta_0 = 0.48$, (C) $\beta_0 = 0.56$, and (D) $\beta_0 = 0.6$ with a large population size N set at 10^8 . The epidemic simulations start with one infectious individual. In semi-logarithmic scale, exponential growth is evident if a straight line fits well several consecutive disease generations of the epidemic curve, whereas a strong downward curvature in semi-logarithmic scale is indicative of sub-exponential growth. Our simulations show that case incidence curves display early sub-exponential growth dynamics even for values of α slightly below 1.0.




Figure S4. Examples of the fits provided by the exponential and generalized-growth models to three infectious disease outbreaks (pandemic influenza in San Francisco in 1918, HIV/AIDS epidemic in Japan, and Ebola epidemic in Western Urban, Sierra Leone in 2014).




Figure S5. Comparison of the goodness of fit provided by the exponential and the generalized-growth models across all of the 21 infectious disease outbreaks (see Table 1).




Figure S6. Mean estimates of the effective reproduction number and the deceleration of growth parameter p derived from our sample of infectious disease datasets (Table 1) were significantly correlated for three estimation periods with an initial phase length comprising 3 to 5 generation intervals.




Figure S7. The 1918 influenza pandemic in San Francisco. Estimates and 95% confidence intervals of the effective reproduction number derived from fitting the generalized-growth model to an increasing length of the early epidemic phase comprising of approximately 3-5 disease generation intervals. The generation interval is assumed to be gamma distributed with the mean of 3 days and standard deviation of 1 day. Estimates and 95% confidence intervals for parameters r and p are also shown.




Figure S8. The 2001 foot-and-mouth disease epidemic in Uruguay.

Estimates and 95% confidence intervals of the effective reproduction number derived from fitting the generalized-growth model to an increasing length of the early epidemic phase comprising of approximately 3-5 disease generation intervals. The generation interval is assumed to be gamma distributed with the mean of 5 days and standard deviation of 1 day. Estimates and 95% confidence intervals for parameters r and p are also shown.




Figure S9. The HIV/AIDS epidemic in Japan (1985-2012).

Estimates and 95% confidence intervals of the effective reproduction number derived from fitting the generalized-growth model to an increasing length of the early epidemic phase comprising of approximately 3-5 disease generation intervals. The generation interval is assumed to be gamma distributed with the mean of 4 years and standard deviation of 1.4 years. Estimates and 95% confidence intervals for parameters r and p are also shown.




Figure S10. The 2014-15 Ebola epidemic in Gueckedou, Guinea.

Estimates and 95% confidence intervals of the effective reproduction number derived from fitting the generalized-growth model to an increasing length of the early epidemic phase comprising of approximately 3-5 disease generation intervals. The generation interval is assumed to be gamma distributed with the mean of 19 days and standard deviation of 11 days. Estimates and 95% confidence intervals for parameters r and p are also shown.





Figure S11. The 2014-15 Ebola epidemic in Western Area Urban, Sierra Leone.

Estimates and 95% confidence intervals of the effective reproduction number derived from fitting the generalized-growth model to an increasing length of the early epidemic phase comprising of approximately 3-5 disease generation intervals. The generation interval is assumed to be gamma distributed with the mean of 11.6 days and standard deviation of 5.6 days. Estimates and 95% confidence intervals for parameters r and p are also shown.



References

1. Viboud C, Simonsen L, Chowell G. A generalized-growth model to characterize the early ascending phase of infectious disease outbreaks *Epidemics* **2016**; 15:27–37.